

# MATH 3100 SPRING 2022. LECTURE SUMMARIES

LEONID PETROV

## 1. LECTURE 1. JANUARY 19

Introductory lecture, just saying hello to the students.

## 2. LECTURE 2. NO DEADLINE

### Section 1.1

A recording of an in-class piece, explaining the definition of the probability space  $\Omega$ , events  $\mathcal{F}$ , and probability measure  $P(A)$ .

## 3. LECTURE 3. JANUARY 21

Sections 1.1–1.2.

- Recalling  $(\Omega, \mathcal{F}, P)$ .
- Finite sample spaces, when it is enough to specify  $P(\{\omega\})$  for each singleton  $\omega \in \Omega$ . We only need to have  $P(\{\omega\}) \geq 0$  and  $\sum_{\omega \in \Omega} P(\{\omega\}) = 1$
- Biased coin, 2 flips — example
- Equally likely outcomes,  $P(A) = \frac{\#A}{\#\Omega}$ .
- Example with urns.

## 4. LECTURE 4. JANUARY 24

Section 1.2. Discussing 3 settings, when we sample  $k$  times from  $n$  objects. This is an instance of equally likely outcomes. Recall the factorial:

$$\begin{cases} n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot n, & n \geq 1 \\ 0! = 1. \end{cases}$$

So,  $0! = 1$ ,  $1! = 1$ ,  $2! = 2$ ,  $3! = 6$ ,  $4! = 24$ , and so on.

- Sampling with replacement, order matters:  $\#\Omega_k = n^k$ .
- Sampling without replacement, order matters:  $\#\Omega_k = n(n-1)\dots(n-k+1) = \frac{n!}{(n-k)!}$ .
- Sampling without replacement, order does not matter:

$$\#\Omega_k = \frac{n(n-1)\dots(n-k+1)}{k!} = \frac{n!}{k!(n-k)!} = \binom{n}{k}.$$

The quantity  $\binom{n}{k}$  has a special name, “binomial coefficient”.