

Lectures on Random Matrices (Spring 2025)

Lecture 3: Gaussian and tridiagonal matrices

Leonid Petrov

Wednesday, January 22, 2025*

Contents

1	Recap	2
2	Gaussian ensembles	2
2.1	Definitions	2
2.2	Joint eigenvalue distribution for GOE	3
2.3	Step A. Joint density of matrix entries	4
2.4	Step B. Spectral decomposition	4
2.5	Step C. Jacobian	5
2.6	Step D. Final Form of the density	6
3	Other classical ensembles with explicit eigenvalue densities	7
3.1	Wishart (Laguerre) ensemble	7
3.1.1	Definition via SVD	7
3.1.2	Joint density of eigenvalues	7
3.2	Jacobi (MANOVA/CCA) ensemble	8
3.2.1	Setup	8
3.2.2	Jacobi ensemble	9
3.3	General Pattern and β -Ensembles	9
4	Tridiagonal form for real symmetric matrices	10
5	Tridiagonalization of random matrices	11
5.1	Dumitriu–Edelman tridiagonal model for GOE	12
5.2	Generalization to β -ensembles	13
C	Problems (due 2025-02-22)	13
C.1	Invariance of GOE and GUE	13
C.2	Preimage size for spectral decomposition	13
C.3	Distinct eigenvalues	14

*[Course webpage](#) • [Live simulations](#) • [TeX Source](#) • Updated at 04:22, Wednesday 26th March, 2025

C.4	Testing distinctness of eigenvalues via rank-1 perturbations	14
C.5	Jacobian for GUE	14
C.6	Normalization for GOE	14
C.7	Wishart eigenvalue density	14
C.8	Householder reflection properties	15
C.9	Distribution of the Householder vector in random tridiagonalization	15
C.10	Householder reflection for GUE	15
C.11	Jacobi ensemble is related to two Wisharts	16

1 Recap

We have established the semicircle law for real Wigner random matrices. If W is an $n \times n$ real symmetric matrix with independent entries X_{ij} above the main diagonal (mean zero, variance 1), and mean zero diagonal entries, then the empirical spectral distribution of W/\sqrt{n} converges to the semicircle law as $n \rightarrow \infty$:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i/\sqrt{n}} = \mu_{\text{sc}}, \quad (1.1)$$

where

$$\mu_{\text{sc}}(dx) = \begin{cases} \frac{1}{2\pi} \sqrt{4 - x^2} dx, & \text{if } |x| \leq 2, \\ 0, & \text{otherwise.} \end{cases}$$

The convergence in (1.1) is weakly almost sure. The way we got the result is by expanding $\mathbb{E} \text{Tr}(W^k)$ and counting trees, plus analytic lemmas which ensure that the convergence of expected powers of traces is enough to conclude the convergence (1.1) of the empirical spectral measures.

Today, we are going to focus on Gaussian ensembles. The plan is:

- Definition and spectral density for real symmetric Gaussian matrices (GOE).
- Other random matrix ensembles with explicit eigenvalue densities: Wishart (Laguerre) and Jacobi (MANOVA/CCA) ensembles.
- Tridiagonalization and general beta ensemble.
- (next week, not today) Wigner's semicircle law via tridiagonalization.

2 Gaussian ensembles

2.1 Definitions

Recall that a real Wigner matrix W can be modeled as

$$W = \frac{Y + Y^\top}{\sqrt{2}},$$

where Y is an $n \times n$ matrix with independent entries Y_{ij} , $1 \leq i, j \leq n$, such that Y_{ij} are mean zero, variance 1. Then for $1 \leq i < j \leq n$, we have for the matrix $W = (X_{ij})$:

$$\text{Var}(X_{ii}) = \text{Var}(\sqrt{2}Y_{ii}) = 2, \quad \text{Var}(X_{ij}) = \text{Var}\left(\frac{Y_{ij} + Y_{ji}}{\sqrt{2}}\right) = 1.$$

If, in addition, we assume that Y_{ij} are standard Gaussian $\mathcal{N}(0, 1)$, then the distribution of W is called the *Gaussian Orthogonal Ensemble* (GOE).

For the complex case, we have the *standard complex Gaussian random variable*

$$Z = \frac{1}{\sqrt{2}}(Z^R + \mathbf{i}Z^I), \quad \mathbb{E}(Z) = 0, \quad \text{Var}_{\mathbb{C}}(Z) := \mathbb{E}(|Z|^2) = \frac{\mathbb{E}(|Z^R|^2) + \mathbb{E}(|Z^I|^2)}{2} = 1,$$

where Z^R and Z^I are independent standard Gaussian real random variables $\mathcal{N}(0, 1)$.

If we take Y to be an $n \times n$ matrix with independent entries Y_{ij} , $1 \leq i, j \leq n$ distributed as Z , then the random matrix¹

$$W = \frac{Y + Y^\dagger}{\sqrt{2}}$$

is said to have the *Gaussian Unitary Ensemble* (GUE) distribution. For the GUE matrix $W = (X_{ij})$, we have for $1 \leq i < j \leq n$:

$$\text{Var}_{\mathbb{C}}(X_{ii}) = 1, \quad \text{Var}_{\mathbb{C}}(X_{ij}) = \frac{1}{4} \left[\mathbb{E}(Z_{ij}^R + Z_{ji}^R)^2 + \mathbb{E}(Z_{ij}^I + Z_{ji}^I)^2 \right] = 1.$$

Both GOE and GUE have real eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$. We are going to describe the joint distribution of these eigenvalues. Despite the fact that the map from a matrix to its eigenvalues is quite complicated and nonlinear (you need to solve an equation of degree n), the distribution of eigenvalues in the Gaussian cases is fully explicit.

See Problem C.1 for invariance of GOE/GUE under orthogonal/unitary conjugation (this is where the names “orthogonal” and “unitary” come from).

Remark 2.1. There is a third player in the game, the *Gaussian Symplectic Ensemble* (GSE), which we will mainly ignore in this course due to its less intuitive quaternionic nature.

2.2 Joint eigenvalue distribution for GOE

In this section, we give a derivation of the joint probability density for the GOE.

Theorem 2.2 (GOE Joint Eigenvalue Density). *Let W be an $n \times n$ real symmetric matrix with the GOE distribution (Section 2.1). Then its ordered real eigenvalues $\lambda_1 \leq \dots \leq \lambda_n$ of $W/\sqrt{2}$ have a joint probability density function on \mathbb{R}^n given by:*

$$p(\lambda_1, \dots, \lambda_n) = \frac{1}{Z_n} \prod_{1 \leq i < j \leq n} |\lambda_i - \lambda_j| \exp\left(-\frac{1}{2} \sum_{k=1}^n \lambda_k^2\right),$$

where Z_n is a constant (depending on n but not on λ_i) ensuring the density integrates to 1:

$$Z_n = Z_n^{\text{GOE}} = \frac{(2\pi)^{n/2}}{n!} \prod_{j=0}^{n-1} \frac{\Gamma(1 + (j+1)\beta/2)}{\Gamma(1 + \beta/2)}, \quad \beta = 1.$$

¹ Y^\dagger denotes the transpose of Y combined with complex conjugation.

Remark 2.3. We renormalized the GOE by a factor of $\sqrt{2}$ to make the Gaussian part of the density, $\exp(-\frac{1}{2} \sum_{k=1}^n \lambda_k^2)$, standard. In the GUE case, no normalization is required.

We break the proof into four major steps, considered in Sections 2.3 to 2.6 below.

2.3 Step A. Joint density of matrix entries

Let us label all independent entries of $W/\sqrt{2}$:

$$\{\underbrace{X_{12}, X_{13}, \dots, X_{23}, \dots}_{\text{above diag}}, \underbrace{X_{22}, X_{33}, \dots}_{\text{diag}}\}.$$

There are $\frac{n(n-1)}{2}$ off-diagonal entries with variance $1/2$, and n diagonal entries with variance 1. The joint density of these entries (ignoring normalization for a moment) is proportional to

$$f(x_{12}, x_{13}, \dots, x_{22}, x_{33}, \dots) \propto \exp\left(-\sum_{i < j} x_{ij}^2 - \frac{1}{2} \sum_{i=1}^n x_{ii}^2\right) = \exp\left(-\frac{1}{2} \sum_{i,j=1}^n x_{ij}^2\right), \quad (2.1)$$

where in the right-hand side, we have $x_{ij} = x_{ji}$ for $i \neq j$. We then recognize

$$\sum_{i,j=1}^n x_{ij}^2 = \text{Tr}(W^2) = \sum_{k=1}^n \lambda_k^2.$$

Including the normalization for Gaussians, one arrives at the density on $\mathbb{R}^{n(n+1)/2}$:

$$f(W) dW = \pi^{-\frac{n(n-1)}{4}} (2\pi)^{-\frac{n}{4}} \exp\left(-\frac{1}{2} \text{Tr}(W^2)\right) dW,$$

where dW is the product measure over the $\frac{n(n+1)}{2}$ independent entries.

2.4 Step B. Spectral decomposition

Since W is real symmetric, it can be orthogonally diagonalized:

$$W = Q \Lambda Q^\top, \quad Q \in O(n),$$

where $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$ has the eigenvalues. Then, as we saw before, we have

$$\text{Tr}(W^2) = \text{Tr}(Q \Lambda Q^\top Q \Lambda Q^\top) = \text{Tr}(\Lambda^2) = \sum_{k=1}^n \lambda_k^2.$$

The map from W to (Λ, Q) is not one-to one, but in case W has distinct eigenvalues, the preimage of (Λ, Q) contains 2^n elements. See Problems C.2 and C.3.

It remains to make the change of variables from W to Λ , which involves the Jacobian.

2.5 Step C. Jacobian

We now examine how the measure dW in the space of real symmetric matrices factors into a piece depending on $\{\lambda_i\}$ and a piece depending on Q . Formally,

$$dW = \left| \det \left(\frac{\partial W}{\partial (\Lambda, Q)} \right) \right| d\Lambda dQ,$$

where dQ is the Haar measure² on $O(n)$, and $d\Lambda$ is the Lebesgue measure on \mathbb{R}^n . The Lebesgue measure later needs to be restricted to the “Weyl chamber” $\lambda_1 \leq \dots \leq \lambda_n$ if we want an ordering, this introduces the simple factor $n!$ in the final density.

Lemma 2.4 (Jacobian for Spectral Decomposition). *For real symmetric $W = Q\Lambda Q^\top$, one has*

$$\left| \det \left(\frac{\partial W}{\partial (\Lambda, Q)} \right) \right| = \text{const} \prod_{1 \leq i < j \leq n} |\lambda_i - \lambda_j|,$$

where the constant is independent of the λ_i ’s and depends only on n .

Remark 2.5. Equivalently, one often writes

$$dW = |\Delta(\lambda_1, \dots, \lambda_n)| d\Lambda dQ, \quad \text{where } \Delta(\lambda_1, \dots, \lambda_n) = \prod_{i < j} (\lambda_j - \lambda_i)$$

is the *Vandermonde determinant*.

We prove Lemma 2.4 in the rest of this subsection.

Consider small perturbations of Λ and Q . Write

$$W = Q \Lambda Q^\top, \quad \Lambda = \text{diag}(\lambda_1, \dots, \lambda_n).$$

Let δW be an infinitesimal change in W . We want to see how δW depends on $\delta\Lambda$ and δQ .

Parametrizing δQ . Since $Q \in O(n)$, any small variation of Q can be expressed as

$$Q \exp(B) \approx Q(I + B),$$

where B is an infinitesimal skew-symmetric matrix ($B^\top = -B$). Indeed, $\exp(B)$ must be orthogonal, so $\exp(B)^\top \exp(B) = I$. Thus, we have

$$(I + B)^\top (I + B) = I, \quad \text{or} \quad B^\top + B = 0.$$

Note that $\exp(B)$ is the matrix exponential of B , which is defined by the usual power series. Note also that the dimension of $O(n)$ is $\dim(O(n)) = \frac{n(n-1)}{2}$, which matches the dimension of the space of skew-symmetric matrices.

²Recall that the Haar measure on $O(n)$ is the unique (up to a constant factor) measure that is invariant under group shifts (in this situation, both left and right shifts work). In probabilistic terms, if a random orthogonal matrix Q is Haar-distributed, then QR and RQ are also Haar-distributed for any fixed orthogonal matrix R .

Computing δW . Under an infinitesimal change, say,

$$Q \mapsto Q(I + B), \quad \Lambda \mapsto \Lambda + \delta\Lambda,$$

we have

$$W = Q\Lambda Q^\top \implies Q^\top \delta W Q = \delta\Lambda + B\Lambda - \Lambda B,$$

to first order in small quantities. Here we used the orthogonality of Q and the skew-symmetry of B .

Local structure of the map. We see that the map $W \mapsto (\Lambda, Q)$ in a neighborhood of (Λ, Q) determined by $\delta\Lambda$ and B locally translates by $Q^\top \delta\Lambda Q$, which implies the Lebesgue factor $d\lambda_1 \dots d\lambda_n$ in δW . Indeed, the Lebesgue measure on \mathbb{R}^n is invariant under orthogonal transformations.

The next terms, the commutator $[B, \Lambda]$, has the form (recall that B is infinitesimally small and Λ is diagonal):

$$\begin{aligned} B\Lambda - \Lambda B &= \begin{pmatrix} 0 & b_{12} & \cdots \\ -b_{12} & 0 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 & \cdots \\ 0 & \lambda_2 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} - \begin{pmatrix} \lambda_1 & 0 & \cdots \\ 0 & \lambda_2 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} 0 & b_{12} & \cdots \\ -b_{12} & 0 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} \\ &= \begin{pmatrix} 0 & b_{12}\lambda_2 & \cdots \\ -b_{12}\lambda_1 & 0 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} - \begin{pmatrix} 0 & b_{12}\lambda_1 & \cdots \\ b_{12}\lambda_2 & 0 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix} \\ &= \begin{pmatrix} 0 & b_{12}(\lambda_2 - \lambda_1) & \cdots \\ b_{12}(\lambda_1 - \lambda_2) & 0 & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix}. \end{aligned}$$

Thus, this action locally means that the infinitesimal b_{ij} is multiplied by $\lambda_i - \lambda_j$, for all $1 \leq i < j \leq n$. This is a scalar factor that does not depend on the orthogonal component Q , but only on the eigenvalues. Therefore, this factor is the same in $Q^\top \delta W Q$.

This completes the proof of Lemma 2.4. See also Problem C.5 for the GUE Jacobian.

2.6 Step D. Final Form of the density

Putting Steps A–C together, we find:

$$dW = \text{const} \cdot \prod_{i < j} |\lambda_i - \lambda_j| d\Lambda \underbrace{\left(\text{Haar measure on } O(n) \right)}_{\text{does not depend on } \lambda_i}.$$

Hence, the joint density of $\{\lambda_1, \dots, \lambda_n\}$ is, up to normalization depending only on n , equal to

$$\prod_{i < j} |\lambda_i - \lambda_j| \exp\left(-\frac{1}{2} \sum_{k=1}^n \lambda_k^2\right). \quad (2.2)$$

We leave the computation of the normalization constant in Theorem 2.2 as Problem C.6.

Remark 2.6. We emphasize that in the GOE case, the normalization $W/\sqrt{2}$ for (2.2) is so that the variance is 1 on the diagonal and $\frac{1}{2}$ off the diagonal.

3 Other classical ensembles with explicit eigenvalue densities

Let us briefly discuss other classical ensembles with explicit eigenvalue densities, which are not necessarily Gaussian, but are related to other classical structures like orthogonal polynomials. These ensembles also have a built-in parameter β (and in the cases $\beta = 1, 2, 4$, they have invariance under orthogonal/unitary/symplectic conjugation).

3.1 Wishart (Laguerre) ensemble

In this subsection, we describe another classical family of random matrices whose eigenvalues form a fundamental example of a β -ensemble with a “logarithmic” pairwise interaction. These are called the *Wishart* or *Laguerre* ensembles. Their importance arises in statistics (covariance estimation, principal component analysis), signal processing, and many other areas.

3.1.1 Definition via SVD

Let X be an $n \times m$ random matrix with iid entries drawn from a real/complex/quaternionic normal distribution. We assume $n \leq m$. We can perform the *singular value decomposition* (SVD) of X :

$$X = U \begin{pmatrix} s_1 & & 0 \\ & \ddots & \\ 0 & & s_n \end{pmatrix} V^\dagger,$$

where U, V are orthogonal/unitary/symplectic matrices (depending on β), $s_1, \dots, s_n \geq 0$ are the singular values of X , and \dagger means the corresponding conjugation. For example, in the real case, s_1, \dots, s_n are the square roots of the eigenvalues of XX^\top .

Moreover, let $W = XX^\dagger$; this is called the Wishart random matrix ensemble. We have

$$\lambda_i = s_i^2, \quad i = 1, \dots, n; \quad \lambda_1 \geq \dots \geq \lambda_n \geq 0.$$

These eigenvalues admit a closed-form joint probability density function (pdf) in complete analogy with the GOE/GUE calculations from previous subsections.

3.1.2 Joint density of eigenvalues

Theorem 3.1 (Wishart eigenvalue density). *The ordered eigenvalues $\lambda_1, \dots, \lambda_n \geq 0$ of the $n \times n$ Wishart matrix W have the joint density on $\{\lambda_i \geq 0\}$ proportional to*

$$\prod_{1 \leq i < j \leq n} (\lambda_i - \lambda_j)^\beta \prod_{i=1}^n \lambda_i^{\frac{\beta}{2}(m-n+1)-1} \exp\left(-\frac{\lambda_i}{2}\right),$$

where $\beta = 1, 2, 4$ corresponds to the real, complex, or quaternionic case, respectively.

Idea of proof (sketch). The proof is a variant of the derivation for the joint eigenvalue density in the GOE/GUE case (see Section 2.2). One writes down the joint distribution of all entries of X , changes variables to singular values and orthogonal/unitary transformations, and identifies the Jacobian factor as $\prod_{i < j} |s_i^2 - s_j^2|^\beta = \prod_{i < j} |\lambda_i - \lambda_j|^\beta$. The extra factors in front arise from the powers of λ_i (i.e. from $\prod_i s_i$) and the Gaussian exponential $\exp(-\frac{1}{2} \sum s_i^2)$ when reshaped to $\exp(-\frac{1}{2} \sum \lambda_i)$. \square

Remark 3.2. The exponent of λ_i in the product is often written as $\alpha = \frac{\beta}{2}(m - n + 1) - 1$. One also sees the name *multivariate Gamma distribution* in statistics. For $\beta = 1$ the ensemble is sometimes called the *real Wishart* (or *Laguerre Orthogonal*) ensemble; for $\beta = 2$ it is the *complex Wishart* (or *Laguerre Unitary*) ensemble; and $\beta = 4$ (not discussed in detail here) is the *symplectic version*. In point processes, the case $\beta = 2$ is also referred to as the *Laguerre orthogonal polynomial ensemble*.

3.2 Jacobi (MANOVA/CCA) ensemble

The *Jacobi* (sometimes called *MANOVA* or *CCA*) ensemble arises when one looks at the interaction between two independent rectangular Gaussian matrices that share the same number of columns. Statistically, this corresponds to questions of canonical correlations or multivariate Beta distributions. In random matrix theory, it appears as yet another fundamental example of a β -ensemble with an explicit eigenvalue density.

3.2.1 Setup

Let X be an $n \times t$ real (or complex) matrix and Y be a $k \times t$ matrix, with $n \leq k \leq t$. Assume X and Y have iid Gaussian entries (real or complex) of mean 0 and variance 1 and are independent of each other.

Definition 3.3 (Projectors and canonical correlations). Denote by

$$P_X = X^\top (X X^\top)^{-1} X \quad (\text{or } X^\dagger (X X^\dagger)^{-1} X),$$

the orthogonal (unitary) projector onto the row span of X . Similarly, define

$$P_Y = Y^\top (Y Y^\top)^{-1} Y.$$

These are $t \times t$ projection matrices of ranks n and k , respectively, embedded in a space of dimension t . One checks that P_X and P_Y commute if and only if the row spaces of X and Y are aligned in a certain way. The *canonical correlations* between these two subspaces are the singular values of $P_X P_Y$. Equivalently, the *squared* canonical correlations are the nonzero eigenvalues of $P_X P_Y$.

Since $\text{rank}(P_X P_Y) \leq \min(n, k)$, there are at most $\min(n, k)$ nonzero eigenvalues of $P_X P_Y$. In fact, generically (when the subspaces are in “general position”), there are exactly $\min(n, k)$ nonzero eigenvalues.

Example 3.4. For $n = k = 1$, we have

$$P_X P_Y = \frac{\langle X, Y \rangle}{\langle X, X \rangle \langle Y, Y \rangle} X^\top Y,$$

which is a rank one matrix with the only nonzero singular eigenvalue $\langle X, Y \rangle$. Therefore, the singular value is exactly the sample correlation coefficient between X and Y .

3.2.2 Jacobi ensemble

Theorem 3.5 (Jacobi/MANOVA/CCA Distribution). *Let X and Y be as above, each having iid (real or complex) Gaussian entries of size $n \times t$ and $k \times t$, respectively, with $n \leq k \leq t$. Assume further that X and Y are independent of each other (this is the null hypothesis in statistics).*

Then the nonzero eigenvalues $\lambda_1, \dots, \lambda_n$ of the matrix $P_X P_Y$ lie in the interval $[0, 1]$ and have the joint density function of the form

$$\prod_{i < j} |\lambda_i - \lambda_j|^\beta \prod_{i=1}^n \lambda_i^{\frac{\beta}{2}(k-n+1)-1} (1 - \lambda_i)^{\frac{\beta}{2}(t-n-k+1)-1},$$

up to a normalization constant that depends on n, k, t (but not on $\{\lambda_i\}$). Here again $\beta = 1$ for the real case and $\beta = 2$ for the complex case.

This distribution is called the *Jacobi* (or *MANOVA*, or *CCA*) ensemble, and it is also sometimes called the *multivariate Beta distribution*. In point processes, the $\beta = 2$ case is often referred to as the *Jacobi orthogonal polynomial ensemble*.

Remark 3.6. The derivation is again parallel to that in the GOE/GUE context, but one now keeps track of the row spaces and the relevant rectangular dimensions. The matrix $(X X^\top)$ (or $(X X^\dagger)$) is invertible with high probability whenever $n \leq t$ and X is in general position. The distribution above reflects the geometry of overlapping projectors in a higher-dimensional space \mathbb{R}^t (or \mathbb{C}^t).

3.3 General Pattern and β -Ensembles

We have now seen three classical examples:

- *Wigner (Gaussian) ensembles* (real/complex/quaternionic),
- *Wishart/Laguerre ensembles* $W = X X^\top$,
- *Jacobi/MANOVA/CCA ensembles*.

Their eigenvalue densities (ordered or unordered) always display the same building blocks:

$$\prod_{1 \leq i < j \leq n} |\lambda_i - \lambda_j|^\beta \times \prod_{i=1}^n V(\lambda_i),$$

where β indicates the real ($\beta = 1$), complex ($\beta = 2$), or symplectic ($\beta = 4$) symmetry class, and $V(\lambda)$ is a single-variable potential function. Such distributions are often referred to as β -ensembles or *log-gases*, reflecting that the factor $\prod_{i < j} |\lambda_i - \lambda_j|^\beta$ can be interpreted as the Boltzmann weight for charges with a logarithmic pairwise repulsion.

Remark 3.7. Beyond these three classical families, there are many other *matrix models* and *discrete distributions* whose eigenvalues produce similar log-gas structures but with different potentials $V(\lambda)$. These share many of the same techniques and phenomena (e.g. local eigenvalue statistics, largest-eigenvalue asymptotics, etc.) that appear throughout modern random matrix theory.

Remark 3.8. For $\beta = 2$, the connection to orthogonal polynomials suggests discrete models of log-gases, which are powered by most known orthogonal polynomials in one variable from the (q-)Askey scheme [KS96]. For example, the model of (uniformly random) lozenge tilings of the hexagon is connected to Hahn orthogonal polynomials [Gor21] whose orthogonality weight is the classical hypergeometric distribution from probability theory.

4 Tridiagonal form for real symmetric matrices

Any real symmetric matrix can be orthogonally transformed into a tridiagonal matrix. This fact is standard in numerical linear algebra (the “Householder reduction”) and also central in random matrix theory—notably in the Dumitriu–Edelman approach [DE02] for Gaussian ensembles.

Theorem 4.1. *Any real symmetric matrix $W \in \mathbb{R}^{n \times n}$ can be represented as*

$$W = Q^\top T Q, \quad Q \in O(n),$$

where T is real symmetric tridiagonal. Concretely, T has nonzero entries only on the main diagonal and the first super-/sub-diagonals:

$$T = \begin{pmatrix} d_1 & \alpha_1 & 0 & \cdots & 0 \\ \alpha_1 & d_2 & \alpha_2 & \cdots & 0 \\ 0 & \alpha_2 & d_3 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & \alpha_{n-1} \\ 0 & 0 & \cdots & \alpha_{n-1} & d_n \end{pmatrix}.$$

Definition 4.2 (Householder reflection). A *Householder reflection* in \mathbb{R}^n is a matrix H of the form

$$H = I - 2 \frac{v v^\top}{\|v\|^2}, \quad v \in \mathbb{R}^n \text{ nonzero column vector.}$$

One checks that $H^\top = H$, $H^2 = I$, and H is orthogonal (i.e. $H^\top H = I$). Geometrically, H is the reflection across the hyperplane orthogonal to v .

Proof of Theorem 4.1. Let $A \in \mathbb{R}^{n \times n}$ be a symmetric matrix. We will show how to orthogonally conjugate A into a tridiagonal matrix T .

Step 1: Zeroing out subdiagonal entries in the first column. Write A in block form as

$$A = \begin{pmatrix} a_{11} & r^\top \\ r & B \end{pmatrix},$$

where $r \in \mathbb{R}^{n-1}$ is the rest of the first column below a_{11} , and B is $(n-1) \times (n-1)$. We seek an orthogonal matrix H_1 acting on \mathbb{R}^{n-1} (and in the full space \mathbb{R}^n it preserves the first basis vector e_1 and its orthogonal complement) that “annihilates” the part of this first column below the subdiagonal. Specifically, H_1 is a Householder reflection chosen so that H_1 when acting in the $(n-1)$ -dimensional subspace spanned by r zeroes out all but the first entry of r . In the ambient

space \mathbb{R}^n , H_1 has a block form, so that it does not touch the 11-entry of the matrix A . Since A is symmetric, conjugating A by H_1 also zeroes out the corresponding superdiagonal entries in the first row. Concretely,

$$H_1 A H_1^\top = \begin{pmatrix} d_1 & \alpha_1 & 0 & \cdots & 0 \\ \alpha_1 & * & * & \cdots & * \\ 0 & * & * & \cdots & * \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & * & * & \cdots & * \end{pmatrix}.$$

This is always possible because Householder reflections can exchange any two given unit vectors. Note also that $\alpha_1 = \|r\|$.

Step 2: Inductive reduction on the trailing principal submatrix. Next, we restrict attention to rows 2 through n and columns 2 through n . Let H_2 be a second Householder reflection that acts as the identity on the first row and column, and zeroes out the subdiagonal entries of the *second* column (viewed within that trailing $(n-1) \times (n-1)$ block). Conjugate again:

$$H_2 (H_1 A H_1^\top) H_2^\top = (H_2 H_1) A (H_1^\top H_2^\top).$$

Now the first two columns (and rows) are in the desired form.

Step 3: Repeat for columns (and rows) 3, 4, By repeating this procedure for each successive column (and row, by symmetry), we eventually force all off-diagonal entries outside the main and first super-/subdiagonals to be zero. After $n-2$ steps, the resulting matrix

$$T = Q^\top A Q, \quad Q = H_1 H_2 \cdots H_{n-2},$$

is *tridiagonal*, and Q is orthogonal because it is a product of orthogonal (Householder) transformations.

Since each H_k is orthogonal, none of these transformations change the eigenvalues of A . Thus T has the same spectrum as A . This completes the tridiagonalization argument. \square

Remark 4.3. This Householder procedure is also used in practical numerical methods for eigenvalue computations: once a real symmetric matrix is reduced to tridiagonal form, specialized algorithms (such as the QR algorithm) can then be applied more efficiently. Overall, computations with tridiagonal matrices are much simpler and with better numerical stability than with general dense matrices.

5 Tridiagonalization of random matrices

Here we discuss the tridiagonal form of the GOE random matrices, and extend it to the general beta case.

5.1 Dumitriu–Edelman tridiagonal model for GOE

Theorem 5.1. *Let W be an $n \times n$ GOE matrix (real symmetric) with variances chosen so that each off-diagonal entry has variance $1/2$ and each diagonal entry has variance 1. Then there exists an orthogonal matrix Q such that*

$$W = Q^\top T Q,$$

where T is a real symmetric tridiagonal matrix of the special form

$$T = \begin{pmatrix} d_1 & \alpha_1 & 0 & \cdots \\ \alpha_1 & d_2 & \alpha_2 & \ddots \\ 0 & \alpha_2 & d_3 & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix},$$

and the random variables $\{d_i, \alpha_j\}_{1 \leq i \leq n, 1 \leq j \leq n-1}$ are mutually independent, with

$$d_i \sim \mathcal{N}(0, 1), \quad \alpha_j = \sqrt{\frac{\chi_{n-j}^2}{2}},$$

where χ_ν^2 is a chi-square distribution with ν degrees of freedom.

Remark 5.2 (Chi-square distributions). The *chi-square distribution* with ν degrees of freedom, denoted by χ_ν^2 , is a fundamental distribution in statistics and probability theory. It arises naturally as the distribution of the sum of the squares of ν independent standard normal random variables. Formally, if Z_1, Z_2, \dots, Z_ν are independent random variables with $Z_i \sim \mathcal{N}(0, 1)$, then the random variable

$$Q = \sum_{i=1}^{\nu} Z_i^2$$

follows a chi-square distribution with ν degrees of freedom, i.e., $Q \sim \chi_\nu^2$. In the context of the Dumitriu–Edelman tridiagonal model (Theorem 5.1), the subdiagonal entries α_j are defined as $\alpha_j = \sqrt{\frac{\chi_{n-j}^2}{2}}$. One can call this a *chi random variable*, as this is a square root of a chi-square variable.

The parameter ν does not need to be an integer, and the chi-square distribution is well defined for any positive real ν , by continuation of the density formula.

Idea of proof of Theorem 5.1. This construction is essentially a specialized version of the Householder reduction in Section 4, set up so that each step matches precisely the distributions $\alpha_j \sim \sqrt{\frac{\chi_{n-j}^2}{2}}$ and $d_i \sim \mathcal{N}(0, 1)$. One uses the rotational invariance of Gaussian matrices to ensure at each step that the “residual vector” is isotropic (i.e., its distribution is invariant under orthogonal transformations). The norm of that vector yields the χ^2 -type variables. \square

Thus, to study the eigenvalues of a GOE matrix W , one can equivalently study the (much sparser) random tridiagonal matrix T .

5.2 Generalization to β -ensembles

The tridiagonal GOE construction (Theorem 5.1) extends to a whole family of ensembles, parametrized by $\beta > 0$. In particular, for $\beta = 1, 2, 4$ we get the classical Orthogonal, Unitary, and Symplectic (GOE/GUE/GSE) ensembles, respectively. The general β case is known as the β -ensemble; outside of the classical cases $\beta = 1, 2, 4$, there is no matrix ensemble interpretation with iid entries, but the tridiagonal form model still works.

We saw that the β -ensembles arise naturally as *log-gases* in physics, with density proportional to

$$\exp\left(-\sum_{i=1}^n V(\lambda_i)\right) \prod_{1 \leq i < j \leq n} |\lambda_i - \lambda_j|^\beta$$

for some potential V . The simplest choice, $V(\lambda) = \frac{1}{2}\lambda^2$, corresponds to Gaussian β -ensembles, which in the classical cases reproduce GOE/GUE/GSE.

Remark 5.3 (Tridiagonal Construction for General β). A breakthrough [DE02] showed that the Gaussian β -ensembles (for *any* $\beta > 0$) can be represented as eigenvalues of real symmetric *tridiagonal* matrices whose entries are independent (but not identically distributed), and have Gaussian and chi distributions:

- The diagonal entries are iid standard normal random variables $\mathcal{N}(0, 1)$.
- The subdiagonal entries are $\alpha_j = \sqrt{\frac{\chi_{(n-j)\beta}^2}{2}}$, where χ_ν^2 is a chi-square distribution with ν degrees of freedom. Here we use the fact that the parameter ν in the chi-square distribution does not need to be an integer.
- The superdiagonal entries are determined by symmetry.

In the next lecture, we will see how the tridiagonal form allows to prove the Wigner's semicircle law for the Gaussian β -ensembles.

C Problems (due 2025-02-22)

C.1 Invariance of GOE and GUE

Show that the distribution of the GOE and GUE is invariant under, respectively, orthogonal and unitary conjugation. For GOE, this means that if W is a random GOE matrix and Q is a fixed orthogonal matrix of order n , then the distribution of QWQ^\top is the same as the distribution of W . (Similarly for GUE.)

Hint: write the joint density of all entries of GOE/GUE (for instance, GOE is determined by $n(n+1)/2$ real random independent variables) in a coordinate-free way.

C.2 Preimage size for spectral decomposition

Show that for a real symmetric matrix W with distinct eigenvalues, if $W = Q\Lambda Q^\top$ is its spectral decomposition where Q is orthogonal and $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$ is diagonal with $(\lambda_1 \geq \dots \geq \lambda_n)$, then there are exactly 2^n different choices of Q that give the same matrix W .

C.3 Distinct eigenvalues

Show that under GOE and GUE, almost surely, all eigenvalues are distinct.

C.4 Testing distinctness of eigenvalues via rank-1 perturbations

Suppose λ is an eigenvalue of a fixed matrix W with multiplicity ℓ . Consider the rank-1 perturbation

$$W_\varepsilon = W + \alpha u u^\top, \quad \alpha \sim \mathcal{N}(0, \varepsilon),$$

where $u \in \mathbb{R}^n$ is fixed. Prove that with probability one (in α), the eigenvalue λ *splits* into ℓ distinct eigenvalues of W_ε .

Hint: Write the characteristic polynomial of W_ε as $\det(W_\varepsilon - \mu I)$. Show that the infinitesimal change in α moves the roots in a non-degenerate way, splitting a repeated root.

C.5 Jacobian for GUE

Arguing similarly to Section 2.5, show that the Jacobian for the spectral decomposition of a complex Hermitian matrix is proportional to

$$\prod_{1 \leq i < j \leq n} |\lambda_i - \lambda_j|^2.$$

In particular, make sure you understand where the factor 2 comes from in the complex case.

C.6 Normalization for GOE

Compute the n -dimensional integral (in the ordered on unordered form):

$$\begin{aligned} \int_{\lambda_1 < \dots < \lambda_n} \prod_{i < j} (\lambda_i - \lambda_j) \exp\left(-\frac{1}{2} \sum_{k=1}^n \lambda_k^2\right) d\lambda_1 \cdots d\lambda_n. \\ = \frac{1}{n!} \int_{\mathbb{R}^n} \prod_{i < j} |\lambda_i - \lambda_j| \exp\left(-\frac{1}{2} \sum_{k=1}^n \lambda_k^2\right) d\lambda_1 \cdots d\lambda_n. \end{aligned}$$

Hint: The following identity might be useful:

$$\int_{-\infty}^{\infty} x^{2m} e^{-x^2/2} dx = 2^{m+1/2} \Gamma\left(m + \frac{1}{2}\right).$$

C.7 Wishart eigenvalue density

Prove Theorem 3.1 (in the real case $\beta = 1$) by using the singular value decomposition of X and the properties of the Wishart ensemble.

C.8 Householder reflection properties

Show that the Householder reflection $H = I - 2vv^\top/\|v\|^2$ has the following properties:

1. H is orthogonal, i.e., $H^\top H = I$.
2. H is symmetric, i.e., $H^\top = H$.
3. H is idempotent, i.e., $H^2 = I$.
4. H is a reflection across the hyperplane orthogonal to v .

C.9 Distribution of the Householder vector in random tridiagonalization

Consider the first step of the Householder tridiagonalization of a GOE matrix W . Denote the first column by $x \in \mathbb{R}^n$, and let

$$v = x + \alpha e_1, \quad \alpha = \pm \|x\|.$$

Then the first Householder reflection is given by

$$H_1 = I - 2 \frac{vv^\top}{\langle v, v \rangle}.$$

Prove that:

1. $\|v\|^2$ follows a χ_ν^2 distribution with ν degrees of freedom (determine ν in terms of n).
2. The direction $v/\|v\|$ is uniformly distributed on the unit sphere \mathbb{S}^{n-1} and is independent of $\|v\|$.

Hint: View x as a Gaussian vector in \mathbb{R}^n , using the fact that the first column of a GOE matrix (including its diagonal entry) is an isotropic normal vector (up to small adjustments for the diagonal). Orthogonal invariance of the underlying distribution ensures the direction is uniform on \mathbb{S}^{n-1} .

C.10 Householder reflection for GUE

Modify the tridiagonalization procedure which was discussed for the GOE case, and show that the GUE random matrix can be transformed (by a unitary conjugation) into

$$\begin{pmatrix} \mathcal{N}(0, 1) & \chi_{2(n-1)}/\sqrt{2} & 0 & 0 & \cdots \\ \chi_{2(n-1)}/\sqrt{2} & \mathcal{N}(0, 1) & \chi_{2(n-2)}/\sqrt{2} & 0 & \cdots \\ 0 & \chi_{2(n-2)}/\sqrt{2} & \mathcal{N}(0, 1) & \chi_{2(n-3)}/\sqrt{2} & \cdots \\ 0 & 0 & \chi_{2(n-3)}/\sqrt{2} & \mathcal{N}(0, 1) & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

(this matrix is symmetric, and in the entries, we list the distributions).

C.11 Jacobi ensemble is related to two Wisharts

Let X be an $n \times m$ and Y be a $k \times m$ real Gaussian matrices with iid $\mathcal{N}(0, 1)$ entries, independent of each other, and assume $n \leq k \leq m$. Consider the matrix

$$(X X^\top + Y Y^\top)^{-1} (X X^\top) \in \mathbb{R}^{n \times n}.$$

1. Prove that it is well-defined (invertible denominator) with probability 1, and that it is symmetric and diagonalizable in \mathbb{R}^n .
2. Show that its eigenvalues lie in $[0, 1]$ and follow a Jacobi (MANOVA) distribution of parameters $\beta = 1$ and (n, k, m) .
3. Identify explicitly how these parameters match the shape parameters in the standard multivariate Beta / Jacobi pdf

$$\prod_{i < j} |\lambda_i - \lambda_j| \prod_{i=1}^n \lambda_i^\alpha (1 - \lambda_i)^\gamma,$$

with appropriate α, γ in terms of n, k, m .

Hint: Use that $X X^\top$ and $Y Y^\top$ are (independent) Wishart matrices. Rewrite

$$(X X^\top + Y Y^\top)^{-1} X X^\top$$

via block-inversion or projector-based arguments to see it is related to the product of two orthogonal projectors in \mathbb{R}^m . The Jacobi distribution then emerges from the overlapping subspace geometry.

References

- [DE02] I. Dumitriu and A. Edelman, *Matrix models for beta ensembles*, Journal of Mathematical Physics **43** (2002), no. 11, 5830–5847. arXiv:math-ph/0206043. [↑10](#), [13](#)
- [Gor21] V. Gorin, *Lectures on random lozenge tilings*, Cambridge Studies in Advanced Mathematics. Cambridge University Press (2021). [↑10](#)
- [KS96] R. Koekoek and R.F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue*, Technical report, Delft University of Technology and Free University of Amsterdam (1996). arXiv:math/9602214 [math.CA], report no. OP-SF 20 Feb 1996. Updated version available at <https://fa.ewi.tudelft.nl/~koekoek/documents/as98.pdf>. [↑10](#)

L. PETROV, UNIVERSITY OF VIRGINIA, DEPARTMENT OF MATHEMATICS, 141 CABELL DRIVE, KERCHOF HALL, P.O. BOX 400137, CHARLOTTESVILLE, VA 22904, USA
E-mail: lenia.petrov@gmail.com