

# Lectures on Random Matrices (Spring 2025)

## Lecture 6: Double contour integral kernel. Steepest descent and local statistics

Leonid Petrov

February 12, 2025\*

### Notes for the lecturer

- GUE det structure
- Formulate Cauchy–Binet and Andreief
- Recall that  $\rho_n = P_n$  and it is  $(\det[\psi_i(x_j)]_{n \times n})^2$ , then reproduce the proofs here.
- Recall the Christoffel–Darboux formula:

$$K_n(x, y) = \frac{e^{-\frac{x^2+y^2}{4}}}{\sqrt{2\pi}h_{n-1}} \frac{p_n(x)p_{n-1}(y) - p_{n-1}(x)p_n(y)}{x - y},$$

here  $h_{n-1} = \sqrt{2\pi}(n-1)!$ .

## 1 Recap: Determinantal structure of the GUE

Last time, we proved the following result:

**Theorem 1.1.** *The GUE correlation functions are given by*

$$\rho_k(x_1, \dots, x_k) = \det \left[ K_n(x_i, x_j) \right]_{i,j=1}^k,$$

with the correlation kernel

$$K_n(x, y) = \sum_{j=0}^{n-1} \psi_j(x) \psi_j(y).$$

Here

$$\psi_j(x) = \frac{1}{\sqrt{h_j}} p_j(x) e^{-x^2/4},$$

where  $p_j(x)$  are the monic Hermite polynomials, and  $h_j$  are the normalization constants so that  $\psi_j(x)$  are orthonormal in  $L^2(\mathbb{R})$ .

---

\*[Course webpage](#) • [Live simulations](#) • [TeX Source](#) • Updated at 03:53, Sunday 9<sup>th</sup> February, 2025

For this theorem, we need Cauchy–Binet summation formula and Andreief identity (which is essentially the same as Cauchy–Binet, but when summation is replaced by integration). Having these, we can write

$$\begin{aligned}
\rho_k(x_1, \dots, x_k) &= \frac{n!}{(n-k)!} \int_{\mathbb{R}^{n-k}} p(x_1, \dots, x_n) dx_{k+1} \cdots dx_n \\
&= \frac{1}{(n-k)! \widehat{Z}_{n,2}} \sum_{\substack{\sigma, \tau \in S_n \\ \sigma(k+1)=\tau(k+1), \dots, \sigma(n)=\tau(n)}} \text{sgn}(\sigma) \text{sgn}(\tau) \prod_{i=1}^k \psi_{\sigma(i)-1}(x_i) \psi_{\tau(i)-1}(x_i) \\
&= \text{const}_n \sum_{I \subseteq [n], |I|=k} \sum_{\sigma', \tau' \in S(I)} \text{sgn}(\sigma') \text{sgn}(\tau') \prod_{i=1}^k \psi_{\sigma'(i)-1}(x_i) \psi_{\tau'(i)-1}(x_i) \\
&= \text{const}_n \sum_{I \subseteq [n], |I|=k} \det [\psi_{i_\alpha}(x_j)]_{\alpha, j=1}^k \det [\psi_{i_\alpha}(x_j)]_{\alpha, j=1}^k,
\end{aligned}$$

where  $I = \{i_1, \dots, i_k\}$  is a subset of  $[n]$  of size  $k$ , and  $S(I)$  is the set of permutations of  $I$ . The last sum of products of two determinants is written by the Cauchy–Binet formula as

$$\text{const}_n \cdot \det \left[ \sum_{j=0}^{n-1} \psi_j(x_\alpha) \psi_j(x_\beta) \right]_{\alpha, \beta=1}^k,$$

and finally the constant is equal to 1 by Andreief identity.

## 2 Double Contour Integral Representation for the GUE Kernel

### 2.1 One contour integral representation for Hermite polynomials

Recall that the GUE kernel is defined by

$$K_N(x, y) = \sum_{n=0}^{N-1} \psi_n(x) \psi_n(y),$$

with the orthonormal functions

$$\psi_n(x) = \frac{1}{\sqrt{h_n}} p_n(x) e^{-x^2/4},$$

where the (monic, probabilists') Hermite polynomials are given by

$$p_n(x) = (-1)^n e^{x^2/2} \frac{d^n}{dx^n} e^{-x^2/2}. \tag{2.1}$$

Note that the monic Hermite polynomials are uniquely defined by the orthogonality property. We are not proving (2.1) here, it is an exercise.

**Lemma 2.1** (Generator function for Hermite polynomials). *We have*

$$\exp\left(xt - \frac{t^2}{2}\right) = \sum_{n \geq 0} p_n(x) \frac{t^n}{n!}.$$

*The series converges for all  $t$  since the left-hand side is an entire function of  $t$ .*

*Proof.* Write the generating function as

$$\sum_{n \geq 0} p_n(x) \frac{t^n}{n!} = \sum_{n \geq 0} \frac{(-1)^n t^n}{n!} e^{x^2/2} \frac{d^n}{dx^n} e^{-x^2/2}.$$

Since the factor  $e^{x^2/2}$  does not depend on  $n$ , we can factor it out:

$$\sum_{n \geq 0} p_n(x) \frac{t^n}{n!} = e^{x^2/2} \sum_{n \geq 0} \frac{(-t)^n}{n!} \frac{d^n}{dx^n} e^{-x^2/2}.$$

Now, recall Taylor's theorem: for any analytic function  $f$  we have

$$f(x - t) = \sum_{n \geq 0} \frac{(-t)^n}{n!} f^{(n)}(x).$$

Applying this with  $f(x) = e^{-x^2/2}$ , we deduce that

$$\sum_{n \geq 0} \frac{(-t)^n}{n!} \frac{d^n}{dx^n} e^{-x^2/2} = e^{-(x-t)^2/2}.$$

Thus, our generating function becomes

$$\sum_{n \geq 0} p_n(x) \frac{t^n}{n!} = e^{x^2/2} e^{-(x-t)^2/2},$$

as desired. □

By Cauchy's integral formula we can write using Lemma 2.1:

$$p_n(x) = \frac{n!}{2\pi i} \oint_C \frac{\exp\left(xt - \frac{t^2}{2}\right)}{t^{n+1}} dt, \quad (2.2)$$

where the contour  $C$  is a simple closed curve encircling the origin. Indeed, here we use the complex analysis property

$$\frac{1}{2\pi i} \oint_C \frac{1}{z^{k+1}} dz = \begin{cases} 1, & \text{if } k = 0, \\ 0, & \text{if } k \neq 0, \end{cases}$$

so (2.2) is simply a complex analysis version of the operation of extracting the coefficient of  $t^n$  in the Taylor expansion.

Therefore,

$$\psi_n(x) = \frac{1}{\sqrt{h_n}} p_n(x) e^{-x^2/4} = \frac{e^{-x^2/4}}{\sqrt{h_n}} \frac{n!}{2\pi i} \oint_C \frac{\exp\left(xt - \frac{t^2}{2}\right)}{t^{n+1}} dt.$$

## 2.2 Another contour integral representation for Hermite polynomials

We start with the Fourier transform identity

$$\int_{-\infty}^{\infty} \exp\left(-\frac{t^2}{2} + i t x\right) dt = \sqrt{2\pi} e^{-x^2/2}.$$

Differentiating both sides  $n$  times with respect to  $x$  yields

$$\frac{d^n}{dx^n} \left( e^{-x^2/2} \right) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (i t)^n e^{-t^2/2 + i t x} dt.$$

Recalling the definition

$$p_n(x) = (-1)^n e^{x^2/2} \frac{d^n}{dx^n} \left( e^{-x^2/2} \right),$$

we obtain

$$p_n(x) = \frac{(-1)^n e^{x^2/2}}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (i t)^n e^{-t^2/2 + i t x} dt.$$

Next, perform the change of variable

$$s = i t, \quad \text{so that} \quad t = -i s, \quad dt = -i ds.$$

Under this substitution the factors transform as follows:

$$(i t)^n = s^n,$$

and the exponent becomes

$$-\frac{t^2}{2} + i t x = -\frac{(-i s)^2}{2} + i (-i s) x = \frac{s^2}{2} + s x.$$

Thus, the integral transforms into

$$\int_{-\infty}^{\infty} (i t)^n e^{-t^2/2 + i t x} dt = -i \int_{-i\infty}^{i\infty} s^n e^{s^2/2 + s x} ds.$$

Substituting back we have

$$p_n(x) = \frac{(-1)^n e^{x^2/2}}{\sqrt{2\pi}} (-i) \int_{-i\infty}^{i\infty} s^n e^{s^2/2 + s x} ds.$$

That is,

$$p_n(x) = \frac{i (-1)^{n+1} e^{x^2/2}}{\sqrt{2\pi}} \int_{-i\infty}^{i\infty} s^n e^{s^2/2 + s x} ds.$$

Finally, change the sign of  $s$ , and we get:

$$p_n(x) = \frac{i e^{x^2/2}}{\sqrt{2\pi}} \int_{-i\infty}^{i\infty} s^n e^{s^2/2 - s x} ds.$$

Therefore,

$$\psi_n(x) = \frac{i e^{x^2/4}}{\sqrt{2\pi} h_n} \int_{-i\infty}^{i\infty} s^n e^{s^2/2 - s x} ds.$$

### 2.3 Normalization of Hermite polynomials

**Lemma 2.2.** *We have*

$$h_n = \int_{-\infty}^{\infty} p_n(x)^2 e^{-x^2/2} dx = n! \sqrt{2\pi}.$$

*Proof.* Multiply the generating function

$$\exp\left(xt - \frac{t^2}{2}\right) = \sum_{n \geq 0} p_n(x) \frac{t^n}{n!}$$

with a second copy (with parameter  $s$ ):

$$\exp\left(xs - \frac{s^2}{2}\right) = \sum_{m \geq 0} p_m(x) \frac{s^m}{m!}.$$

Then,

$$\exp\left(xt - \frac{t^2}{2}\right) \exp\left(xs - \frac{s^2}{2}\right) = \sum_{n, m \geq 0} p_n(x) p_m(x) \frac{t^n s^m}{n! m!}.$$

Integrate both sides against  $e^{-x^2/2} dx$ . Using the orthogonality

$$\int_{-\infty}^{\infty} p_n(x) p_m(x) e^{-x^2/2} dx = h_n \delta_{nm},$$

the right-hand side becomes

$$\sum_{n \geq 0} \frac{h_n}{(n!)^2} (ts)^n.$$

On the left-hand side, we have

$$\int_{-\infty}^{\infty} e^{-x^2/2} \exp\left(x(t+s) - \frac{t^2+s^2}{2}\right) dx.$$

Completing the square in  $x$  or recalling the standard Gaussian integral yields

$$\sqrt{2\pi} \exp\left(\frac{(t+s)^2 - (t^2+s^2)}{2}\right) = \sqrt{2\pi} \exp(ts).$$

Thus, we obtain

$$\sqrt{2\pi} \exp(ts) = \sum_{n \geq 0} \frac{h_n}{(n!)^2} (ts)^n.$$

Expanding the left side as

$$\sqrt{2\pi} \sum_{n \geq 0} \frac{(ts)^n}{n!},$$

and comparing coefficients, we conclude that

$$\frac{h_n}{(n!)^2} = \frac{\sqrt{2\pi}}{n!} \implies h_n = n! \sqrt{2\pi}.$$

This completes the proof. □

## 2.4 Double contour integral representation for the GUE kernel

We can sum up the kernel (essentially, this is another proof of the Christoffel–Darboux formula):

$$\begin{aligned}
K_n(x, y) &= \sum_{k=0}^{n-1} \psi_k(x) \psi_k(y) \\
&= \frac{e^{\frac{x^2-y^2}{4}}}{(2\pi)^2} \oint_C dt \int_{-i\infty}^{i\infty} ds \exp \left\{ -\frac{t^2}{2} + xt + \frac{s^2}{2} - ys \right\} \underbrace{\sum_{k=0}^{n-1} s^k t^{-k-1}}_{\frac{1-(s/t)^n}{t-s}}. \tag{2.3}
\end{aligned}$$

Here we used the two contour integral representations for Hermite polynomials, and the explicit norm (Lemma 2.2). At this point, the  $t$  contour is a small circle around 0, and the  $s$  contour is a vertical line in the complex plane. Their mutual position can be arbitrary at this point — the  $s$  contour goes along the imaginary line. Indeed, the fraction  $\frac{1-(s/t)^n}{t-s}$  does not have a singularity at  $s = t$  due to the cancellation.

Let us now move the  $s$  contour to be to the left of the  $t$  contour, as in Figure 1. On the new contours, we have  $|s| > |t|$ . Now we can add the summands  $s^k t^{-k-1}$  for all  $k \leq -1$  into the sum in (2.3). Indeed, for  $|s| > |t|$ , the series in  $k$  converges, while the summand  $s^k t^{-k-1}$  has zero residue at 0 and thus adding the summands does not change the value of the integral.




Figure 1: Integration contours for the GUE kernel (2.4).

With this extension of the sum, formula (2.3) becomes

$$K_n(x, y) = \frac{e^{(y^2-x^2)/4}}{(2\pi)^2} \oint_C dt \int_{-i\infty}^{i\infty} ds \frac{\exp \left\{ \frac{s^2}{2} - sy - \frac{t^2}{2} + tx \right\}}{s - t} \left( \frac{s}{t} \right)^n. \tag{2.4}$$

**Remark 2.3.** The  $s$  contour passes to the right of the  $t$  contour, but it might as well pass to the left of it. Indeed, one can deform the  $s$  contour to the left while picking the residue at  $s = t$ :

$$\text{Res}_{s=t} \frac{\exp \left\{ \frac{s^2}{2} - sy - \frac{t^2}{2} + tx \right\}}{s - t} \left( \frac{s}{t} \right)^n = -e^{t(x-y)}.$$

This function is entire in  $t$ , and its integral over the  $t$  contour is zero. Therefore, there is no difference where the  $s$  contour passes with respect to the  $t$  contour.

## 2.5 Extensions

Many other versions of the GUE / unitary invariant ensembles admit determinantal structure:

1. The GUE corners process [JN06]
2. The Dyson Brownian motion (e.g., add a GUE to a diagonal matrix) [NF98]
3. GUE corners plus a fixed matrix [FF14]
4. Corners invariant ensembles with fixed eigenvalues  $UDU^\dagger$ , where  $D$  is a fixed diagonal matrix and  $U$  is Haar distributed on the unitary group [Met13]

We will discuss the corners process structure in the next [Lecture 7](#).

## F Problems (due 2025-03-12)

### References

- [FF14] P. Ferrari and R. Frings, *Perturbed GUE minor process and Warren's process with drifts*, J. Stat. Phys **154** (2014), no. 1-2, 356–377. arXiv:1212.5534 [math-ph]. [↑7](#)
- [JN06] K. Johansson and E. Nordenstam, *Eigenvalues of GUE minors*, Electron. J. Probab. **11** (2006), no. 50, 1342–1371. arXiv:math/0606760 [math.PR]. [↑7](#)
- [Met13] A. Metcalfe, *Universality properties of Gelfand-Tsetlin patterns*, Probab. Theory Relat. Fields **155** (2013), no. 1-2, 303–346. arXiv:1105.1272 [math.PR]. [↑7](#)
- [NF98] T. Nagao and P.J. Forrester, *Multilevel dynamical correlation functions for Dyson's Brownian motion model of random matrices*, Physics Letters A **247** (1998), no. 1-2, 42–46. [↑7](#)

L. PETROV, UNIVERSITY OF VIRGINIA, DEPARTMENT OF MATHEMATICS, 141 CABELL DRIVE, KERCHOF HALL, P.O. BOX 400137, CHARLOTTESVILLE, VA 22904, USA  
E-mail: [lenia.petrov@gmail.com](mailto:lenia.petrov@gmail.com)