

Studying the Kosterlitz-Thouless Transition of the two dimensional XY model using Monte Carlo methods

Lennart Voorgang

Bachelorarbeit in Physik
angefertigt am Institute for Advanced Simulation
des
Forschungszentrum Jülich

vorgelegt der
Mathematisch-Naturwissenschaftlichen Fakultät
der
Rheinischen Friedrich-Wilhelms-Universität
Bonn

August 2025

I hereby declare that this thesis was formulated by myself and that no sources or tools other than those cited were used.

Bonn,
Date

.....
Signature

1. Assessor: Prof. Dr. Stefan Krieg
2. Assessor: TBD

Contents

1. Introduction	1
2. Theory	3
2.1. Phase Transitions in Thermodynamics	3
2.2. XY Model	3
2.2.1. Definition	3
2.2.2. Observables	3
2.2.3. Kosterlitz–Thouless Transition	3
2.2.4. Vortices	3
2.3. Numerical Methods	3
2.3.1. Metropolis Algorithm	3
2.3.2. Wolff Cluster Algorithm	4
2.3.3. Bootstrapping	4
A. Source Code	A
List of Figures	I
Bibliography	III

1. Introduction

In physics there are a variety of problems for which it is either very hard or impossible to find analytical solutions. These problems often involve many particle systems and/or integrals of high dimensionality. With ever increasing compute power available to researchers, it becomes more and more feasible to study such system using Monte Carlo methods. Monte Carlo methods is an umbrella term for numerical methods which employ random numbers.

The perhaps most well-known application of Monte Carlo methods in physics is the Ising model. It describes the total magnetization of a lattice as the superposition of all spins $\sigma = \pm 1$ on the lattice. The Ising model features a phase transition at a critical temperature where the system transitions from an ordered low-temperature to a unordered high-temperature state.

One can now generalize the problem from a discrete \mathbb{Z}_2 symmetry to a continuous $U(2)$ symmetry. This is called the XY model and describes the spins as two-dimensional vectors on the unit circle. Unlike the Ising model, the XY model does not have a 2nd order phase transition. It does however have something we call KT phase transition where below a critical temperature T_C metastable states can exist. These metastable states are closely related to vortex/anti-vortex pairs on the lattice. The discovery of the KT phase transition was awarded with the Nobel prize in 2016.

In this bachelor thesis I will be employing Monte Carlo methods, namely the Metropolis and Wolff algorithms, to study the two dimensional XY model.

2. Theory

2.1. Phase Transitions in Thermodynamics

Test

2.2. XY Model

2.2.1. Definition

The XY model describes spins as two-dimensional vectors on the unit circle

$$\sigma_i = \begin{pmatrix} \cos \theta_i \\ \sin \theta_i \end{pmatrix} \quad (2.1)$$

parametrized by the angle $\theta_i \in [0, 2\pi)$.

2.2.2. Observables

2.2.3. Kosterlitz–Thouless Transition

2.2.4. Vortices

Test

2.3. Numerical Methods

2.3.1. Metropolis Algorithm

The overall procedure for the lattice is as follows:

1. Calculate E and M for the initial lattice configuration.
2. Perform a lattice sweep by iterating over all lattice sites. For every site i do:

2. Theory

- a) Propose a new angle $\theta_i \in [0, 2\pi)$ from a uniform distribution.
- b) Calculate $\Delta H = \Delta E$ and ΔM for the proposed new state.
- c) Accept or reject state with probability $P = \min(1, \exp(-\beta\Delta H))$.
3. Update the observables $E += \Delta E$ and $M += \Delta M$ and add them to the result set.
4. Repeat from 2. for a total of N sweeps.

2.3.2. Wolff Cluster Algorithm

2.3.3. Bootstrapping

Since the Metropolis-Hastings algorithm scales with the number of lattice sites, it is computationally impractical to run the Metropolis-Hastings algorithm for long timescales. Given that we have an observable whose iid samples follow a Gaussian distribution and we already have “enough” iid samples which cover enough of the possible configuration space, we can then use bootstrapping to generate more samples (?).

1. Collect B intermediate means by repeating the following:
 - a) Take A random samples from the blocked samples obtained in ?? with replacement.
 - b) Calculate the mean of those samples and add the result to the set of intermediate means.
2. Calculate the final mean and the sample standard deviation of those B intermediate means.

A. Source Code

The program requires a *gcc*¹ or *Intel® oneAPI DPC++/C++ Compiler*² installation with support for *C++26*. The build process was tested with version 15.1.1 and 2025.0.4 respectively. The source code for the simulation as well as the sources for this report can be obtained from GitHub:

```
https://github.com/lennartvrg/physik690-Bachelorarbeit
```

As the program makes use of submodules, it is necessary to clone with

```
git clone --recurse-submodules [...]
```

to automatically initialize all submodules.

Dependencies The program requires several third party libraries to compile. The following packages names are those used in the *Ubuntu launchpad*³.

```
sudo apt-get install meson libpqxx-dev libsimde-dev  
↪ libboost-all-dev libfftw3-dev libtomlplusplus-dev  
↪ libflatbuffers-dev libsqlite3-dev libtbb-dev
```

The build process was tested with the versions available in *Ubuntu 25.04* on July 22, 2025. Build errors might occur when using older versions. In particular for *libpqxx-dev* at least version 7.10.x is required.

Building First create a build directory in the repository root via `mkdir buildDir`. Then, the build files can be generated either for *gcc* via

```
CC=gcc CXX=g++ meson setup --reconfigure  
↪ --buildtype=release buildDir/ .
```

or for the *Intel® oneAPI DPC++/C++ Compiler* via

```
CC=icx CXX=icpx meson setup --reconfigure  
↪ --buildtype=release buildDir/ .
```

¹<https://gcc.gnu.org/>

²<https://www.intel.com/content/www/us/en/developer/tools/oneapi/dpc-compiler.html>

³<https://launchpad.net/ubuntu>

List of Figures

Bibliography