Exercise 6.5 Schwarzschild spacetime in isotropic coordinates

```
{t, r, \theta, \varphi}::Coordinate.
{a,b,c,d,e,f,g,h#}::Indices(values={t, r, \theta, \varphi}, position=independent).

\partial{#}::PartialDerivative.

g^{a b}::InverseMetric.  # essential when using complete (gab, $g^{a b}$)

Gamma := \Gamma^{a}_{b c} -> 1/2 g^{a d} (   \partial_{b}{g_{d c}}
                                           + \partial_{c}{g_{b d}}
                                           - \partial_{d}{g_{b c}}).

Rabcd := R^{a}_{b c d} ->   \partial_{c}{\Gamma^{a}_{b d}}
                          - \partial_{d}{\Gamma^{a}_{b c}}
                          + \Gamma^{e}_{b d} \Gamma^{a}_{c e}
                          - \Gamma^{e}_{b c} \Gamma^{a}_{d e}.

Rab := R_{a b} -> R^{c}_{a c b}.

gab := { g_{t t}            = -((2*r-m)/(2*r+m))**2,
         g_{r r}            = (1+m/(2*r))**4,
         g_{\theta\theta}   = r**2 (1+m/(2*r))**4,
         g_{\varphi\varphi} = r**2 \sin(\theta)**2 (1+m/(2*r))**4}.  # cdb(ex-0605.101,gab)

complete   (gab, $g^{a b}$)                                          # cdb(ex-0605.102,gab)

substitute (Rabcd, Gamma)
substitute (Rab, Rabcd)

evaluate   (Gamma, gab, rhsonly=True)                                # cdb(ex-0605.103,Gamma)
evaluate   (Rabcd, gab, rhsonly=True)                                # cdb(ex-0605.104,Rabcd)
evaluate   (Rab,   gab, rhsonly=True)                                # cdb(ex-0605.105,Rab)
```

$$\left[g_{tt}=-\left(\left(2r-m\right)\left(2r+m\right)^{-1}\right)^{2},\ g_{rr}=\left(1+\frac{1}{2}mr^{-1}\right)^{4},\ g_{\theta\theta}=r^{2}\left(1+\frac{1}{2}mr^{-1}\right)^{4},\ g_{\varphi\varphi}=r^{2}(\sin\theta)^{2}\left(1+\frac{1}{2}mr^{-1}\right)^{4}\right] \tag{ex-0605.101}$$

$$\left[g_{tt}=-\left(\left(2r-m\right)\left(2r+m\right)^{-1}\right)^{2},\ g_{rr}=\left(1+\frac{1}{2}mr^{-1}\right)^{4},\ g_{\theta\theta}=r^{2}\left(1+\frac{1}{2}mr^{-1}\right)^{4},\ g_{\varphi\varphi}=r^{2}(\sin\theta)^{2}\left(1+\frac{1}{2}mr^{-1}\right)^{4},\ g^{tt}= -(m+2r)^{2}(-m+2r)^{-2},\ g^{rr}=\left(\frac{1}{2}mr^{-1}+1\right)^{-4},\ g^{\theta\theta}=\left(r^{2}\left(\frac{1}{2}mr^{-1}+1\right)^{4}\right)^{-1},\ g^{\varphi\varphi}=\left(r^{2}\left(\frac{1}{2}mr^{-1}+1\right)^{4}(\sin\theta)^{2}\right)^{-1}\right] \tag{ex-0605.102}$$

$$\Gamma^{a}{}_{bc}\to\Box_{cb}{}^{a}\begin{cases}\Box_{\varphi r}{}^{\varphi}=\left(-m+2r\right)\left(r\left(m+2r\right)\right)^{-1}\\ \Box_{\varphi\theta}{}^{\varphi}=(\tan\theta)^{-1}\\ \Box_{\theta r}{}^{\theta}=\left(-m+2r\right)\left(r\left(m+2r\right)\right)^{-1}\\ \Box_{rr}{}^{r}=-2m(r\left(m+2r\right))^{-1}\\ \Box_{tr}{}^{t}=4m\left(-m^{2}+4r^{2}\right)^{-1}\\ \Box_{r\varphi}{}^{\varphi}=\left(-m+2r\right)\left(r\left(m+2r\right)\right)^{-1}\\ \Box_{\theta\varphi}{}^{\varphi}=(\tan\theta)^{-1}\\ \Box_{r\theta}{}^{\theta}=\left(-m+2r\right)\left(r\left(m+2r\right)\right)^{-1}\\ \Box_{rt}{}^{t}=4m\left(-m^{2}+4r^{2}\right)^{-1}\\ \Box_{\varphi\varphi}{}^{r}=r\left(m-2r\right)(\sin\theta)^{2}(m+2r)^{-1}\\ \Box_{\varphi\varphi}{}^{\theta}=-\frac{1}{2}\sin\left(2\theta\right)\\ \Box_{\theta\theta}{}^{r}=r\left(m-2r\right)\left(m+2r\right)^{-1}\\ \Box_{tt}{}^{r}=-64mr^{4}\left(m-2r\right)\left(m+2r\right)^{-7}\end{cases} \tag{ex-0605.103}$$

$$R^{a}{}_{bcd} \to \square_{db}{}^{a}{}_{c} \begin{cases} \square_{tt}{}^{r}{}_{r} = -128m^3r^3(m+2r)^{-8} + 512m^2r^4(m+2r)^{-8} - 512mr^5(m+2r)^{-8} \\ \square_{\theta\theta}{}^{r}{}_{r} = -4mr\big(m^2+4mr+4r^2\big)^{-1} \\ \square_{\varphi\varphi}{}^{\theta}{}_{\theta} = 8mr(\sin\theta)^2(m+2r)^{-2} \\ \square_{\varphi\varphi}{}^{r}{}_{r} = -4mr(\sin\theta)^2\big(m^2+4mr+4r^2\big)^{-1} \\ \square_{tr}{}^{t}{}_{r} = -8m\big(r\left(m^2+4mr+4r^2\right)\big)^{-1} \\ \square_{\theta r}{}^{\theta}{}_{r} = 4m\big(r\left(m^2+4mr+4r^2\right)\big)^{-1} \\ \square_{\varphi\theta}{}^{\varphi}{}_{\theta} = (m-2r)^2(m+2r)^{-2} - 1 \\ \square_{\varphi r}{}^{\varphi}{}_{r} = 4m\big(r\left(m^2+4mr+4r^2\right)\big)^{-1} \\ \square_{rt}{}^{r}{}_{t} = 128m^3r^3(m+2r)^{-8} - 512m^2r^4(m+2r)^{-8} + 512mr^5(m+2r)^{-8} \\ \square_{r\theta}{}^{r}{}_{\theta} = 4mr\big(m^2+4mr+4r^2\big)^{-1} \\ \square_{\theta\varphi}{}^{\theta}{}_{\varphi} = (m-2r)^2(\sin\theta)^2(m+2r)^{-2} - (\sin\theta)^2 \\ \square_{r\varphi}{}^{r}{}_{\varphi} = 4mr(\sin\theta)^2\big(m^2+4mr+4r^2\big)^{-1} \\ \square_{rr}{}^{t}{}_{t} = 8m\big(r\left(m^2+4mr+4r^2\right)\big)^{-1} \\ \square_{rr}{}^{\theta}{}_{\theta} = -4m\big(r\left(m^2+4mr+4r^2\right)\big)^{-1} \\ \square_{\theta\theta}{}^{\varphi}{}_{\varphi} = 8mr(m+2r)^{-2} \\ \square_{rr}{}^{\varphi}{}_{\varphi} = -4m\big(r\left(m^2+4mr+4r^2\right)\big)^{-1} \\ \square_{\varphi\varphi}{}^{t}{}_{t} = -4mr(\sin\theta)^2(m+2r)^{-2} \\ \square_{\theta\theta}{}^{t}{}_{t} = -4mr(m+2r)^{-2} \\ \square_{tt}{}^{\varphi}{}_{\varphi} = 64mr^3(m-2r)^2(m+2r)^{-8} \\ \square_{tt}{}^{\theta}{}_{\theta} = 64mr^3(m-2r)^2(m+2r)^{-8} \\ \square_{t\varphi}{}^{t}{}_{\varphi} = 4mr(\sin\theta)^2(m+2r)^{-2} \\ \square_{t\theta}{}^{t}{}_{\theta} = 4mr(m+2r)^{-2} \\ \square_{\varphi t}{}^{\varphi}{}_{t} = -64mr^3(m-2r)^2(m+2r)^{-8} \\ \square_{\theta t}{}^{\theta}{}_{t} = -64mr^3(m-2r)^2(m+2r)^{-8} \end{cases} \tag{ex-0605.104}$$

$$R_{ab} \to 0 \tag{ex-0605.105}$$