M11482 PROBLEM SET 2

Leo Alcock, Luis Alvernaz M1182-teachers@esp.mit.edu

Problems marked with * are more difficult; we think...

problem numba Problem statement

1: Open balls are open Given a metric space (X, d), we defined the open ball, $N_r(p)$ about a point $p \in X$ of radius r to be

$$N_r(p) = \{ q \in X | d(p, q) < r \}$$

We defined a set $S \subset X$ to be open if for every element in S an open ball around that point is contained in S. Show that open balls are open and conclude after that we can equivalently define a set S to be open if it is the union of a collection of open balls.

- 2: Open/Closed relationship We defined a closed set to be a subset which contains all of it's limit points. Show that a set $S \subset X$ is closed if and only if the complement of it (in X) is open (and vice versa).
- 3: Topology? Show that an arbitrary union of open sets $\{U_{\alpha}\}$ is open in X and that finite intersections of open sets are open. Conclude by this and problem 2 that arbitrary intersections of closed sets are closed and finite unions of closed sets are closed.¹
- 4: Circle, diamond, and square metric With $X = \mathbb{R}^2$, recall the following three metrics defined in class:

$$d_1(p,q) := \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

$$d_2(p,q) := |x_1 - x_2| + |y_1 - y_2|$$

$$d_3(p,q) := \max(|x_1 - x_2|, |y_1 - y_2|)$$

With these three metrics, we get three different metric spaces: (X, d_1) , (X, d_2) , and (X, d_3) . Show that a set is open in one of these spaces if and only if it is open in another. hint: Show given $N_{r,d_i}(p)$ there exists r' > 0 such that $N_{r',d_j}(p) \subset N_{r,d_i}(p)$.

Date: July 16, 2017.

¹Good exercise to find an arbitrary intersection of open sets which isn't open and an arbitrary union of closed sets which isn't closed

5*: Closure Given an arbitrary subset $S \subset X$ of some metric space (X, d), define $\bar{S} = \bigcap_{C \in \mathscr{C}_S} C$ where \mathscr{C}_S is the collection of all closed sets containing S. Show that \bar{S} is the union of S and all of S's limit points.

Can you define a corresponding notion to closure for open sets?