

Galaxy Clusters & AGN Feedback

The Clusters Hiding in Plain Sight (CHiPS) Survey


KIPAC Tea talk

12-05-2020

Taweewat Somboonpanyakul | Massachusetts Institute of Technology

Advisor: Michael McDonald


In collaboration with: A. Nobel, M. Bayliss, M. Gaspari, A. Stark, B. Stadler



Galaxy Clusters

Typical Properties

- Most massive gravitational bounded objects in the universe
- Consist of many galaxy members
- Host a central BCG and giant dark matter halo
- Contain extremely hot gas (10^7 K) inside a halo (ICM)
 - Emitting X-ray radiation



Two type of Clusters


Relaxed Clusters vs Disturbed Clusters

- E.g., **Abell 1835**
 - Mostly cool core
(cool/denser in a core)
 - Has some cooling ($\sim 100 \text{ M}_\odot \text{ yr}^{-1}$)
 - Host a radio-loud AGN



Credit: A. Morandi

Two type of Clusters



Disturbed Cluster

- E.g., **El Gordo** •
- Mostly non-cool core •
- Has almost no cooling ($<10 M_{\odot} \text{ yr}^{-1}$) •
- Host a radio-quiet AGN •
- Highly disturbed morphology for the ICM •

Two type of Clusters

Relaxed Cluster

- E.g., **Abell 1835**
 - Mostly cool core
(cool/denser in a core)
 - Has some cooling ($\sim 100 \text{ M}_\odot \text{ yr}^{-1}$)
 - Host a radio-loud AGN


Disturbed Cluster

- E.g., **El Gordo**
 - Mostly non-cool core
 - Has almost no cooling ($< 10 \text{ M}_\odot \text{ yr}^{-1}$)
 - Host a radio-quiet AGN
 - Highly disturbed morphology for the ICM

These are what massive clusters look like.

But they are not what clusters should look like.

Suppression of Star Formation



The Phoenix Cluster

The most famous exception to the rule




- Extremely high star formation rate (~ $600 \text{ M}_\odot \text{ yr}^{-1}$)
- A strong cool core (core X-ray density $> 0.1 \text{ cm}^{-3}$)
- Relaxed morphology for the ICM

How rare is the Phoenix Cluster?

The Clusters Hiding in Plain Sight (CHiPS) Survey

Goal: To find Phoenix-like clusters from X-ray all-sky surveys


The Tale of the Phoenix Discovery



How many of these so-called X-ray point sources are in fact clusters?

ROSAT ALL-SKY SURVEY Bright Sources

Aitoff Projection
Galactic II Coordinate System




www.eso.org/9-Jul-98


Energy range: 0.1 - 2.4 keV
Number of RASS-II sources: 18811
Hardness ratio: -1.0 | -0.4 | -0.2 | 0.2 | 0.6 | 1.0 (soft -> hard : magenta - red - yellow - green - cyan)

Matching with other all-sky surveys

(X-ray=ROSAT, Mid-IR=WISE, Near-IR=2MASS, Radio=NVSS/SUMSS)




Optical Follow-up




● SDSS ● PAN-STARRS ● PISCO

Cluster Richness vs Redshift



Cluster Richness vs Redshift



CHIPS1911+4455

Phoenix meets El Gordo

The Phoenix Cluster

The most famous exception

- Extremely high star formation rate
- A strong cool core
- Relaxed morphology for the ICM

El Gordo

The most massive cluster

- Very low star formation rate
- Non cool-core clusters
- Highly disturbed morphology for the ICM

CHIPS1911+4455

- Extremely high star formation rate
- A strong cool core
- ~~Relaxed morphology for the ICM~~
- ~~Very low star formation rate~~
- ~~Non cool-core clusters~~
- Highly disturbed morphology for the ICM

BUT, this is not what massive clusters should look like.

100 kpc


CHIPS1911+4455

Phoenix meet El Gordo ($z=0.485$)

Nordic Optical Telescope
for optical spectra




- **Extremely high star formation rate**
- A strong cool core
- Highly disturbed morphology for the ICM



SFR [OII] = $120 \pm 15 \text{ M}_\odot \text{ yr}^{-1}$
SFR [WISE4] = $143 \pm 30 \text{ M}_\odot \text{ yr}^{-1}$

Chandra X-ray Image

- Observed in 2019 with ACIS-I for 30.5 ks
- This image is adaptively smoothed.
- It shows a cool-core in the center
- Also shows highly asymmetries on large scale.



CHIPS1911+4455

Phoenix meet El Gordo

- Extremely high star formation rate
- A strong cool core
- Highly disturbed morphology for the ICM






100 kpc

Hubble Image

- Observed in 2019 during Cycle 27 Mid-Cycle
- With F550M (1 orbit) and F110W (1 orbit) using ACS/WFC
- F550M: blue continuum and bright [OII] doublets
- F110W: red continuum for elliptical galaxies

100 kpc



What we learned from CHiPS



- The rarity of these extreme cases told us that CCA might be a mechanism for the feedback
- The survey found a new mode of cooling with **CHIPS1911+4455** (massive starburst in a merging cluster)

Pretty rare ($2\pm1\%$)



Contributions

AGN Feedback in Galaxy Clusters



Finding more extreme clusters will help us unlock mysteries of AGN feedback.