

ARTICLE

Model-based optimization of antibody galactosylation in CHO cell culture

Pavlos Kotidis^{1*} | Philip Jedrzejewski^{1,2,3*} | Si Nga Sou^{1,2,3†} | Christopher Sellick^{4‡} | Karen Polizzi^{2,3} | Ioscani Jimenez del Val⁵ | Cleo Kontoravdi¹

¹Department of Chemical Engineering, Imperial College London, United Kingdom

²Department of Life Sciences, Imperial College London, United Kingdom

³Centre for Synthetic Biology and Innovation, Imperial College London, United Kingdom

⁴Cell Culture and Fermentation Sciences BioPharmaceutical Development, MedImmune, Granta Park, Cambridge, United Kingdom

⁵School of Chemical & Bioprocess Engineering, University College Dublin, Ireland

Correspondence

Cleo Kontoravdi, Department of Chemical Engineering, Imperial College London SW7 2AZ, United Kingdom.
Email: cleo.kontoravdi@imperial.ac.uk

Present address

[†]MedImmune, Granta Park, Cambridge, United Kingdom.

[‡]Kymab Ltd., Babraham Research Campus, Cambridge, United Kingdom.

Abstract

Exerting control over the glycan moieties of antibody therapeutics is highly desirable from a product safety and batch-to-batch consistency perspective. Strategies to improve antibody productivity may compromise quality, while interventions for improving glycoform distribution can adversely affect cell growth and productivity. Process design therefore needs to consider the trade-off between preserving cellular health and productivity while enhancing antibody quality. In this work, we present a modeling platform that quantifies the impact of glycosylation precursor feeding – specifically that of galactose and uridine – on cellular growth, metabolism as well as antibody productivity and glycoform distribution. The platform has been parameterized using an initial training data set yielding an accuracy of $\pm 5\%$ with respect to glycoform distribution. It was then used to design an optimized feeding strategy that enhances the final concentration of galactosylated antibody in the supernatant by over 90% compared with the control without compromising the integral of viable cell density or final antibody titer. This work supports the implementation of Quality by Design towards higher-performing bioprocesses.

KEYWORDS

antibody glycosylation, Chinese hamster ovary (CHO) cells, galactosylation, mathematical modeling, nucleotide sugars, process optimization

1 | INTRODUCTION

Monoclonal antibodies (mAbs) are the most successful biopharmaceutical products of the last decade (Urquhart, 2018; Walsh, 2014). Chinese hamster ovary (CHO) cells are the workhorse of recombinant protein production in the biopharmaceutical industry, owing to their ability to perform complex posttranslational modifications and secrete proteins that are compatible with and bioactive in humans (Lai, Yang, & Ng, 2013; Yusufi et al., 2017). CHO cells are usually grown in fed-batch cultures to replenish the depleted nutrients, prolong the cell culture period, and maximize protein productivity.

Most commercial mAbs are N-linked glycosylated in their constant fragment (Fc). Earlier studies have shown that N-linked glycosylation is strongly dependent on bioprocess conditions, such as temperature (Sou et al., 2015), nutrient availability (Liu et al., 2014), and catabolite accumulation (Yang & Butler, 2000). Bioprocess conditions affect the monosaccharide composition and relative abundance of the oligosaccharides, which in turn, can alter the safety and potency of the mAb. The presence of terminal galactose residues on mAb Fc glycans is known to increase complement-dependent cytotoxicity (Hodoniczky, Zheng, & James, 2005) and antibody-dependent cytotoxicity (Thomann, Reckermann, Reusch, Prasser, & Tejada, 2016) and is, therefore, a desirable attribute in oncological products. In addition, galactosylation is a major source of

*These authors are considered as co-first author.

mAb structural variability during scale-up and technology transfer (Gramer et al., 2011). Manufacturing variability can, therefore, affect product structure consistency and potentially impact efficacy (Goetze et al., 2011; Shinkawa et al., 2003; Yu et al., 2012). Several efforts have been undertaken to increase galactosylation mainly by feeding galactose and uridine, the metabolic precursors of uridine diphosphate galactose (UDPGal), to the cell culture (Grainger & James, 2013; Gramer et al., 2011; Wong et al., 2010). UDPGal is the nucleotide sugar donor (NSD) required for galactosylation as the cosubstrate of the β -1,4-galactosyltransferase (GalT). However, the benefits of precursor addition towards galactosylation can be offset by metabolic perturbations, compromised cell growth, and reduced product titer (Grainger & James, 2013).

Mathematical models are useful tools that have been widely used to describe cell metabolism (Kontoravdi et al., 2010; Nolan & Lee, 2011) and N-linked glycosylation (Jimenez del Val, Nagy, & Kontoravdi, 2011; Kontoravdi & del Val, 2018; Krambeck & Betenbaugh, 2005; Spahn et al., 2016). Metabolic flux analysis and flux balance analysis are usually conducted to unveil the most active metabolic pathways in cells during culture. Glycosylation flux analysis has also been proposed to calculate the intracellular fluxes and distributions of glycans (Hutter et al., 2017; Kremkow & Lee, 2018). In an effort to provide a holistic model for mAb production, Jedrzejewski et al. (2014) linked the extracellular concentrations of nutrients to glycosylation, using a model framework that included the nucleotide and NSD synthesis pathways as the channel between cell metabolism and glycosylation. The proposed concept opened a new path for monitoring mAb production and glycan distribution through mathematical models (Kaveh et al., 2013).

After the Quality by Design (QbD) paradigm, where the end-product quality is rationally built into the manufacturing process, a mathematical model that could describe the perturbations affected by different feeding strategies on cell metabolism and mAb glycosylation would be a valuable tool for process and design space optimization. In order to bridge this gap, herein we propose a combined experimental and computational strategy that captures the effects of feeding strategies on cell metabolism, intracellular NSD concentrations, and mAb glycan distribution. The proposed mechanistic model is able to capture the metabolic alterations caused by the addition of uridine and galactose at different concentrations. The model has been trained with a set of experimental data, and its predictive capabilities have been validated against an independent feeding experiment that was designed using model-based optimization. The proposed model sets the groundwork for in silico glycan optimization by means of metabolic control, speeding up process development, and reducing glycoform variability.

2 | MATERIALS AND METHODS

2.1 | Cell culture

A CHO cell line (kindly donated by MedImmune, Cambridge, UK), producing an IgG antibody, was used in this study. The cells were maintained in suspension culture in CD CHO medium (Life

Technologies, Paisley, UK) at 36.5°C and 5% CO₂ on an orbital shaking platform rotating at 150 rpm and were passaged every 3 days. Fifty micrometer methionine sulfoximine was added for the first two passages postrevival. The experiments were conducted in 500 ml vented Erlenmeyer flasks with a working volume of 100 ml at a seeding density of 2×10^5 cells·ml⁻¹. Cell concentration was determined using a Neubauer ruling haemocytometer, and viability was estimated by the trypan blue dye exclusion method using light microscopy. All cultures were supplemented with 10% v/v CD EfficientFeed C™ C AGT™ Nutrient supplement (Life Technologies) on all even culture days starting on Day 2. Cell cultures were supplemented with 1 µM manganese(II) chloride solution at seeding as well as varying amounts of D-(+)-galactose (6.5 mM to 50 mM in total culture volume) and uridine (0.76 mM to 20 mM in total culture volume; all Sigma-Aldrich, Dorset, UK). Cultures were terminated at Day 12 or when cell viability dropped below 60%. In total, six different feeding strategies were followed: (a) no galactose and uridine addition (control experiment), (b) addition of 10 mM galactose on Days 4 and 8 (10G), (c) addition of 10 mM galactose and 5 mM uridine on Days 4 and 8 (10G5U), (d) addition of 10 mM galactose and 20 mM uridine on Days 4 and 8 (10G20U), (e) addition of 50 mM galactose and 5 mM uridine on Days 4 and 8 (50G5U) and (f) addition of 6.5 mM galactose and 0.76 mM uridine on Day 4, addition of 9.3 mM galactose and 1.34 mM uridine on Day 6, addition of 9 mM galactose and 2.8 mM uridine on Day 8 and addition of 8.7 mM galactose and 10 mM uridine on Day 10 (independent experiment).

2.2 | Metabolite and antibody quantification

Antibody concentration in cell culture supernatant was determined using the BLItz® system and the Dip and Read™ Protein A Biosensors (all Pall ForteBio, Portsmouth, UK). Extracellular glucose, glutamine, glutamate, ammonia, and lactate concentrations were measured using the BioProfile 400 (NOVA Biomedical, MA). Extracellular galactose concentration was determined using the Amplex® Red Galactose/Galactose Oxidase Assay Kit (Life Technologies) according to the manufacturer's instructions. Intracellular glutamine concentration was determined using the Glutamine EnzyChrom™ Assay Kit (Universal Biologicals, Cambridge, UK). Extracellular uridine concentrations were measured using an optimized high-performance anion-exchange chromatography (HPAEC) method as described in del Val, Kyriakopoulos, Polizzi, and Kontoravdi (2013).

2.3 | Nucleotide and NSD quantification

Cell culture samples containing 2×10^6 viable cells were centrifuged at 800 rpm. In a washing step the resulting cell pellet was resuspended in 2 ml of ice-cold 0.9% w/v sodium chloride and then immediately centrifuged at 800 rpm to remove supernatant residues. The cell pellet was resuspended in 400 µl of ice-cold 50% v/v aqueous acetonitrile (Sigma-Aldrich, Gillingham, UK). The solution was incubated on ice for 10 minutes before centrifugation

(4°C, 10,000g, 5 min). The nucleotide and nucleotide sugar containing supernatant was dried using a SpeedVac (Savant Inc. Laboratory, MI) and resuspended in 150 µl water. The final supernatant sample was filtered using a 0.22 µm syringe filter and stored at -80°C prior to the analysis.

The HPAEC method described in del Val et al. (2013) was also used for intracellular nucleotide, nucleoside, and nucleotide sugar quantification. Briefly, elutions were performed with gradients of 1.5 M sodium acetate solution (Sigma-Aldrich) in 3 mM potassium hydroxide (Sigma-Aldrich) with a maximum ion concentration of 1 M sodium acetate using a CarboPac PA1 column (Dionex, Bannockburn, IL). A modified method was used for the quantification of GDP-Fuc and GDP-Man to prevent species coelution with intracellular tryptophan. Elutions were performed with gradients of 1.5 M sodium acetate solution in 3 mM potassium hydroxide. Elutions of the sodium acetate solution were carried out using the following gradients: t_0 min = 20%, t_1 min = 66%, t_6 min = 66%, t_{23} min = 20%, and t_{30} min = 20%. Detection was performed at 262.1 nm.

2.4 | Antibody glycan analysis

Secreted antibody was purified from cell culture supernatant with a Protein A IgG Purification Kit (Thermo Fisher Scientific, Horsham, UK) and concentrated using Vivaspin® 2 ultracentrifugation spin columns 50 kDa (Generon, Maidenhead, UK). Further glycan processing and glycan quantification was kindly performed by MedImmune. The workflow involved the denaturation of purified and concentrated IgG samples, PNGase digestion, and glycan labeling. Separation/quantification was performed using the LabChip® platform (PerkinElmer, Seer Green, UK) on the basis of capillary electrophoresis.

2.5 | Mathematical modeling

Model construction, simulation, parameter estimation, and optimization were conducted in gPROMS v.5.0.1 (Process System Enterprise Ltd, London, U.K., www.psenterprise.com/gproms). Unless otherwise stated, the model parameters were estimated using the maximum likelihood optimization formulation in gPROMS that maximizes the probability that the mathematical model will predict the experimental measurements by assigning values to the physical and variance model parameters. The same solver was used to design the independent feeding strategy used for validation. The model framework comprises three submodels: an unstructured cell growth, death, and metabolism model, a model describing the NSD synthesis and an N-linked glycosylation model (Jimenez del val et al., 2011; del Val et al. 2016). The cell culture model estimates the specific cell growth and specific mAb productivity that are used as inputs for the NSD synthesis model. The latter calculates the NSD concentration and the fluxes of NSDs from the cytosol to the Golgi environment, which are fed to the N-linked glycosylation model to estimate the distribution of mAb glycoforms.

2.6 | Statistical analysis

The MedCalc Statistical Software v.18.2.1 (MedCalc Software bvba, Ostend, Belgium) was used for the estimation of the Concordance Correlation Coefficient (CCC; Lin, 1989) for a 95% confidence interval, to quantify the degree of agreement of the model framework with experimental results, alongside with the R^2 . CCC and R^2 were used to quantify the agreement of two sets of variables, which in our case were the observed and the modeling results. The Pearson Correlation Coefficient (PCC) was, in addition, calculated and presented for completeness, even though it is limited to quantifying the correlation of two variable sets and lacks the ability to describe their agreement. OriginPro 2017 (OriginLab, Northampton, MA) was used for the calculation of PCC for a 95% confidence interval.

3 | MODEL CONSTRUCTION AND PARAMETER ESTIMATION

3.1 | Cell culture model

The cell culture model framework was adapted from the previous work on fed-batch hybridoma cell cultures (Kontoravdi et al., 2010). The model is comprised of three modules: a CHO cell growth and death model, a CHO cell metabolism model, and a mAb production model. The control, 10G, 10G5U, 10G20U, and 50G5U experiments were used for model construction and parameter estimation.

3.1.1 | CHO cell growth and death

CHO cell growth description was on the basis of modified Monod equations that were defined by the limiting substrates and inhibiting metabolites. The equations of the model framework that describe CHO cell growth and death are listed below:

$$\frac{dV}{dt} = F_{in} - F_{out} \quad (1)$$

$$\frac{d(VX_v)}{dt} = (\mu - \mu_{death})VX_v - F_{out}X_v \quad (2)$$

Equation (1) describes the material balance for the cell culture volume V (L) according to the inlet F_{in} ($L \cdot h^{-1}$) and the outlet F_{out} ($L \cdot h^{-1}$) flow rates. Equation (2) describes the material balance for the cell density X_v ($cell \cdot L^{-1}$), where μ (h^{-1}) and μ_{death} (h^{-1}) indicate the specific cell growth and death rate, respectively.

$$\mu = \mu_{max} f_{lim} f_{inh} \quad (3)$$

$$f_{lim} = \frac{[Glc]}{[Glc] + K_{Glc}} \frac{[Asn]}{[Asn] + K_{Asn}} \quad (4)$$

$$f_{inh} = \frac{K_{I_{Amm}}}{[Amm] + K_{I_{Amm}}} \frac{K_{I_{Lac}}}{[Lac] + K_{I_{Lac}}} \frac{K_{I_{Urd}}}{[Urd] + K_{I_{Urd}}} \quad (5)$$

Equations (3–5) describe the specific cell growth rate, where $\mu_{\max}$ (h^{-1}) is the maximum specific cell growth rate, $f_{\text{lim}}(-)$ is the substrate limiting factor and $f_{\text{inh}}(-)$ is the metabolite inhibiting factor. Glucose and asparagine were regarded as the limiting substrates, while lactate, ammonia, and uridine were considered to inhibit growth. The latter two were also regarded as toxic for cell growth. K_{Glc} (mM) and K_{Asn} (mM) are the Monod constants for glucose and asparagine, respectively. $K_{\text{I,amm}}$ (mM), $K_{\text{I,lac}}$ (mM), and $K_{\text{I,urd}}$ (mM) are the inhibiting constants for the ammonia, lactate, and uridine, respectively. $[\text{Glc}]$, $[\text{Asn}]$, $[\text{Amm}]$, $[\text{Lac}]$, and $[\text{Urd}]$ are the extracellular concentrations (mM) of glucose, asparagine, ammonia, lactate, and uridine, respectively.

$$\mu_{\text{death}} = \mu_{\text{death,max}} \left(\frac{[\text{Amm}]}{[\text{Amm}] + K_{\text{d,Amm}}} + \frac{[\text{Urd}]}{[\text{Urd}] + K_{\text{d,Urd}}} \right) \quad (6)$$

Equation (6) describes the specific cell death rate, where $\mu_{\text{death,max}}$ (h^{-1}) is the maximum specific cell death rate and $K_{\text{d,Amm}}$ (mM) and $K_{\text{d,Urd}}$ (mM) are the ammonia and uridine constants for cell death, respectively.

3.1.2 | CHO cell metabolism

Cell metabolism was described by the material balances of substrates, products, and amino acids that were calculated according to Equation (7):

$$\frac{d(V[\text{Metabolite}])}{dt} = F_{\text{in}}[\text{Metabolite}_{\text{feed}}] - F_{\text{out}}[\text{Metabolite}] + q_{\text{metabolite}} V X_v \quad (7)$$

where, $[\text{Metabolite}_{\text{feed}}]$ (mM) is the concentration of each metabolite in the feed, $[\text{Metabolite}]$ (mM) is the extracellular concentration of each metabolite in the culture and $q_{\text{metabolite}}$ ($\text{mmol} \cdot \text{cell}^{-1} \cdot \text{h}^{-1}$) is the specific production/consumption rate of each metabolite and is on the basis of the intracellular metabolic links between the metabolites as discussed below. The sign convention used in the model for the specific rates is positive (+) for the secreted and negative (−) for the consumed metabolites, respectively. The specific production/consumption rates for each metabolite are listed below. In Equations (8–16), $Y_{\text{X,metabolite}}$ ($\text{cell} \cdot \text{mmol}^{-1}$) describes the yield of cell biomass on the metabolite and is assumed to remain constant for each metabolite during the cell culture process to avoid overparameterization.

$$q_{\text{Glc}} = \left(-\frac{\mu}{Y_{\text{X,Glc}}} - m_{\text{Glc}} \right) \cdot \left(\frac{K_{\text{G,Gal}}}{K_{\text{G,Gal}} + [\text{Gal}]} \right)^{n_{\text{Gal}}} \quad (8)$$

$$n_{\text{Gal}} = 1 - f_{\text{Gal}} \frac{q_{\text{Gal}}}{q_{\text{Glc}}} \quad (9)$$

Equations (8) and (9) describe the specific consumption rate of glucose. Moreover, the specific consumption rate of glucose was found to depend on the respective galactose rate, as described in Equation (9). The link between the two metabolites is expected as galactose feeds into the glycolytic pathway through its enzymatic conversion into glucose 1-phosphate. The $\frac{K_{\text{G,Gal}}}{K_{\text{G,Gal}} + [\text{Gal}]}$ term in Equation (8) is used as a regulator when galactose is fed in the

culture. In the absence of galactose the above term is equal to 1 and does not affect the glucose specific rate. m_{Glc} ($\text{mmol} \cdot \text{cell}^{-1} \cdot \text{h}^{-1}$) is the maintenance coefficient of glucose for other metabolic pathways of the cell, $K_{\text{G,Gal}}$ (mM) is the regulating concentration of galactose, and n_{Gal} (−) and f_{Gal} (−) are factors that participate in regulating the specific uptake rate of glucose. n_{Gal} accounts for the fraction of the galactose-specific consumption rate and the respective glucose consumption rate. In Equation (9) when the galactose consumption rate increases, n_{Gal} decreases and consequently from Equation (8) q_{Glc} increases. The majority of glucose transporters (GLUTs) in mammalian cells show affinity to both glucose and galactose and, therefore, a regulation of the glucose and galactose flux from the GLUTs is possible (Wieczorke, Dlugai, Krampe, & Boles, 2003; Zhao & Keating, 2007).

$$q_{\text{Gln}} = \frac{\mu}{Y_{\text{X,Gln}}} + q_{\text{Amm}} Y_{\text{Gln/Amm}} \quad (10)$$

where, $Y_{\text{Gln/Amm}}$ ($\text{mmol}_{\text{Gln}} \cdot \text{mmol}_{\text{Amm}}^{-1}$) is the yield of glutamine from ammonia.

$$q_{\text{Lac}} = \left(\frac{\mu}{Y_{\text{X,Lac}}} - Y_{\text{Lac/Glc}} q_{\text{Glc}} \right) \frac{(\text{Lac}_{\text{max1}} - [\text{Lac}])}{\text{Lac}_{\text{max1}}} + m_{\text{lac}} \frac{\text{Lac}_{\text{max2}} - [\text{Lac}]}{\text{Lac}_{\text{max2}}} \quad (11)$$

where, $Y_{\text{Lac/Glc}}$ ($\text{mmol}_{\text{Lac}} \cdot \text{mmol}_{\text{Glu}}^{-1}$) is the yield of lactate from glucose, Lac_{max1} (mM) and Lac_{max2} (mM) are kinetic constants for the lactate consumption that occurs during the stationary phase (Zagari, Jordan, Stettler, Broly, & Wurm, 2013) and m_{lac} ($\text{mmol} \cdot \text{cell}^{-1} \cdot \text{h}^{-1}$) is the maintenance coefficient of lactate for other metabolic pathways of the cell.

$$q_{\text{Amm}} = \frac{\mu}{Y_{\text{X,Amm}}} - Y_{\text{Amm/Urd}} q_{\text{Urd}} \quad (12)$$

where, $Y_{\text{Amm/Urd}}$ ($\text{mmol}_{\text{Amm}} \cdot \text{mmol}_{\text{Urd}}^{-1}$) is the yield of ammonia from uridine.

$$q_{\text{Glu}} = -\frac{\mu}{Y_{\text{X,Glu}}} \quad (13)$$

$$q_{\text{Gal}} = -\frac{\mu}{Y_{\text{X,Gal}}} \frac{[\text{Gal}]}{[\text{Gal}] + K_{\text{G,Gal}}} \quad (14)$$

where, $K_{\text{G,Gal}}$ (mM) is a Monod constant. The $\frac{[\text{Gal}]}{[\text{Gal}] + K_{\text{G,Gal}}}$ term functions as a regulator of the specific consumption rate of galactose.

$$q_{\text{Urd}} = \frac{\mu}{Y_{\text{X,Urd}}} \frac{[\text{Urd}]}{[\text{Urd}] + K_{\text{Urd}}} \quad (15)$$

where, K_{Urd} (mM) is a Monod constant that regulates the effect uridine concentration on the specific consumption rate of uridine, with the same way that was previously described for galactose.

$$q_{\text{Asn}} = -\frac{\mu}{Y_{\text{X,Asn}}} - Y_{\text{Asn/Asp}} q_{\text{Asp}} \quad (16)$$

where, $Y_{Asn/Asp}$ ($\text{mmol}_{Asn} \cdot \text{mmol}_{Asp}^{-1}$) is the yield of asparagine from aspartate.

$$q_{Asp} = -\frac{\mu}{Y_{X/Asp}} - Y_{Asp/Asn} q_{Asn} \quad (17)$$

where, $Y_{Asp/Asn}$ ($\text{mmol}_{Asp} \cdot \text{mmol}_{Asn}^{-1}$) is the yield of aspartate from asparagine.

3.1.3 | mAb production

mAb synthesis was described with the material balance of the extracellular mAb concentration and the respective specific production rate:

$$\frac{d(V[mAb])}{dt} = -F_{out}[mAb] + q_{mAb} V X_v \quad (18)$$

$$q_{mAb} = Y_{mAbX} \mu + m_{mAb} \quad (19)$$

where, $[mAb]$ ($\text{pg} \cdot \text{l}^{-1}$) is the extracellular mAb concentration, q_{mAb} ($\text{pg} \cdot \text{cell}^{-1} \cdot \text{h}^{-1}$) is the specific productivity of mAbs, Y_{mAbX} ($\text{pg} \cdot \text{cell}^{-1}$) is the yield of mAbs from cell growth, and m_{mAb} ($\text{pg} \cdot \text{cell}^{-1} \cdot \text{h}^{-1}$) is a nongrowth associated term.

3.2 | NSD metabolic model

The NSD metabolic model uses the specific production rate of mAbs and the specific cell growth rate as inputs. Throughout the NSD model, the nucleotides are assumed to be in excess, as supported by

our experimental measurements (data not shown) and, therefore, were not included in the rate calculations. The rates are on the basis of simple Michaelis-Menten saturation kinetics and the substrates used are the intracellular NSD concentrations, the extracellular concentrations of glucose, galactose, and uridine and the intracellular concentration of glutamine. Apart from glutamine, the intracellular concentrations of the metabolites were not estimated in order to avoid oversimplifications and overparameterization. The intracellular concentration of glutamine was linked to the respective extracellular through this simple equation:

$$[Gln_{intra}] = f_{Gln}[Gln] \quad (20)$$

where, $[Gln_{intra}]$ (mM) is the intracellular concentration of glutamine and f_{Gln} is the secretion factor of glutamine to the extracellular environment.

The rates were calculated according to the proposed NSD metabolic pathway (Figure 1) and the kinetic mechanisms that have been reported for each enzyme (Kanehisa & Goto, 2000; Schomburg et al., 2002).

The rates ($\text{mmol}_{NSD} \cdot \text{L}_{cell}^{-1} \cdot \text{h}^{-1}$) as shown in Figure 1 are listed below:

$$r_1 = V_{max1} \frac{[Gln_{intra}]}{K_{M1Gln} + [Gln_{intra}]} \quad (22)$$

$$r_{1sink} = V_{max1sink} \frac{[UDPGlcNAc]}{(K_{M1sink} + [UDPGlcNAc]) \left(1 + \frac{[CMPNeu5Ac]}{K_{I1sink}}\right)} \quad (23)$$




FIGURE 1 The simplified NSD metabolic pathway. The reactions are indicated by solid lines and the transport rates by dotted lines. The transport of extracellular metabolites is not included in model formulation. NSD: nucleotide sugar donor

$$r_2 = V_{\max 2} \frac{[\text{Glc}]}{K_{M2\text{Glc}} + [\text{Glc}]} \quad (24)$$

$$r_{2b} = V_{\max 2b} \frac{[\text{UDPGal}]}{K_{M2b\text{UDPGal}} \left(1 + \frac{[\text{UDPGlcNac}]}{K_{I2A}} + \frac{[\text{UDPGalNac}]}{K_{I2B}} + \frac{[\text{UDPGlc}]}{K_{I2C}} + \frac{[\text{UDPGal}]}{K_{I2D}} \right) + [\text{UDPGal}]} \quad (25)$$

$$r_3 = V_{\max 3} \frac{[\text{Glc}]}{K_{M3\text{Glc}} + [\text{Glc}]} \quad (26)$$

$$r_4 = V_{\max 4} \frac{[\text{UDPGlcNac}]}{K_{M4\text{UDPGlcNac}} + [\text{UDPGlcNac}]} \quad (27)$$

$$r_5 = V_{\max 5} \frac{[\text{UDPGlcNac}]}{K_{M5\text{UDPGlcNac}} \left(1 + \frac{[\text{CMPNeu5Ac}]}{K_{I5}} \right) + [\text{UDPGlcNac}]} \quad (28)$$

$$r_6 = V_{\max 6} \frac{[\text{UDPGlc}]}{K_{M6\text{UDPGlc}} \left(1 + \frac{[\text{UDPGlcNac}]}{K_{I6A}} + \frac{[\text{UDPGalNac}]}{K_{I6B}} + \frac{[\text{UDPGal}]}{K_{I6C}} \right) + [\text{UDPGlc}]} \quad (29)$$

$$r_{\delta\text{sink}} = V_{\max \delta\text{sink}} \frac{[\text{UDPGal}]}{K_{\delta\text{sink}} \left(1 + \frac{[\text{UDPGlc}]}{K_{I\delta\text{sink}}} \right) + [\text{UDPGal}]} \frac{[\text{Gal}]}{[\text{Gal}] + K_{\text{regulator}}} \quad (30)$$

$$r_7 = V_{\max 7} \frac{[\text{GDPMan}]}{\left(K_{M7\text{GDPMan}} + [\text{GDPMan}] \right) \left(1 + \frac{[\text{GDPFuc}]}{K_{I7}} \right)} \quad (31)$$

$$r_{7\text{sink}} = V_{\max 7\text{sink}} \frac{[\text{GDPFuc}]}{K_{M7\text{sink}} + [\text{GDPFuc}]} \quad (32)$$

where, $V_{\max, i}$ ($\text{mmol}_{\text{NSD}} \cdot \text{L}_{\text{cell}}^{-1} \cdot \text{h}^{-1}$) is the maximum turnover rate of reaction i , $[\text{NSD}]$ (mM) are the cytosolic NSD concentrations, $K_{M\text{NSD}}$ (mM) are the saturation constants of the NSD in the i reaction, K_{Ii} (mM) is the inhibition constant in the i reaction and $K_{\text{regulator}}$ (mM) is a fixed parameter that deactivates the regulating $\frac{[\text{Gal}]}{[\text{Gal}] + K_{\text{regulator}}}$ term when no galactose is fed. The r_{sink} terms (not shown in Figure 1) describe the flux of NSD used in reactions that are not included in the chosen metabolic framework and were found to affect GDP-Fuc, UDPGal, and UDPGlcNac concentration.

The contribution of uridine and galactose to NSDs concentration was described with separate reaction rates ($\text{mmol}_{\text{NSD}} \cdot \text{L}_{\text{cell}}^{-1} \cdot \text{h}^{-1}$) that are described below:

$$r_{\text{Urd}} = V_{\max, \text{Urd}} \frac{[\text{Urd}]}{K_{M, \text{Urd}} + [\text{Urd}]} \quad (33)$$

where, i stands only for reactions 1, 2, 4, and 6.

$$r_{\delta\text{Gal}} = V_{\max \delta\text{Gal}} \frac{[\text{Gal}]}{K_{M\delta\text{Gal}} \left(1 + \frac{[\text{UDPGal}]}{K_{I\delta\text{D}}} + \frac{[\text{Gal}]}{K_{I\delta\text{E}}} + \frac{[\text{Urd}]}{K_{I\delta\text{F}}} \right) + [\text{Gal}]} \quad (34)$$

The material balance of each NSD is derived from Figure 1 and follows the general form:

$$\frac{d([\text{NSD}])}{dt} = \sum_i v_i r_i - F_{\text{outNSD}} \quad (35)$$

where, v_i is the stoichiometric coefficient for reaction i , r_i is any of the reactions depicted in Figure 1 and F_{outNSD} ($\text{mmol}_{\text{NSD}} \cdot \text{L}^{-1} \cdot \text{h}^{-1}$) describes the flux of each cytosolic NSD into the Golgi apparatus and represents the transport rate of each NSD consumed for the glycosylation of host cell proteins (HCPs), glycolipids, and mAb product, as shown in Equation (36):

$$F_{\text{outNSD}} = \frac{[\text{NSD}]}{K_{\text{TPNSD}} + [\text{NSD}]} \left(\frac{N_{\text{HCP/LipidsNSD}} \mu}{V_{\text{cell}}} + \frac{N_{\text{precursorNSD}} q_{\text{mAb}}}{V_{\text{cell}}} + r_{\text{NSD}_k}^{\text{mAb, glyc}} \right) \quad (36)$$

where, K_{TPNSD} (mM) is the transport protein saturation constant. The first sum term of Equation (36) describes the flux of NSD used for HCPs and glycolipids synthesis, where V_{cell} (L) is the cellular volume and $N_{\text{HCP/LipidsNSD}}$ ($\text{mmol}_{\text{NSD}} \cdot \text{cell}^{-1}$) is the NSD stoichiometric coefficient for HCPs and glycolipids synthesis, as described by del Val, Polizzi et al. (2016). The second term describes the NSD demand for the formation of the precursor oligosaccharide ($\text{Glc}_3\text{Man}_9\text{GlcNac}_2$), where $N_{\text{precursorNSD}}$ ($\text{mmol}_{\text{NSD}} \cdot \text{cell}^{-1}$) is the stoichiometric coefficient of the NSDs that participate in precursor oligosaccharide synthesis (del Val, Polizzi et al. 2016). The third term of Equation (36) describes the fluxes of NSDs in the latter stages of N-linked glycosylation and are described by Equation (37) (Jimenez del val et al., 2011):

$$r_{\text{NSD}_k}^{\text{mAb, glyc}} = \left(\frac{V_{\text{Golgi}}}{V_{\text{cell}}} \right) \int_{z=0}^z \sum_{j=1}^{N.R.} v_{k,j} \cdot r_j(z) dz \quad (37)$$

where, $r_{\text{NSD}_k}^{\text{glyc}}$ ($\text{mmol}_{\text{NSD}} \cdot \text{L}^{-1} \cdot \text{h}^{-1}$) is the rate of the NSD k that is used for mAb glycosylation, k can be any NSD, V_{Golgi} (L^{-1}) is the Golgi volume, $v_{k,j}$ ($\text{mmol}_{\text{NSD}} \cdot \text{mmol}_{\text{OS}}^{-1}$) is the stoichiometric coefficient of NSD k required as a cosubstrate in reaction j , z is the normalized Golgi length and $r_j(z)$ ($\text{mmol}_{\text{OS}} \cdot \text{L}^{-1} \cdot \text{h}^{-1}$) is the rate of glycosylation reaction j along the length of the Golgi that is calculated in the N-linked glycosylation model.

However, the use of Equation (37) in the current modeling framework substantially increases simulation time because the numerical solution requires convergence between the fluxes calculated by Equation (36) and the integral over z . Equation (38) is proposed to reduce the simulation time while maintaining a mechanistic relationship between the NSD and glycosylation submodels. NSD flux for mAb glycosylation is assumed to be a function of oligosaccharide's concentration in the end of the normalized Golgi length ($z=1$) as the residence time of an oligosaccharide molecule in the Golgi (22 min, (Jimenez del val et al., 2011)) is significantly lower than the time scale within which experimentally observed changes in NSD concentrations occur.

$$r_{NSD_k}^{mAb, glyc} = Vel_{Golgi} \left(\frac{V_{Golgi}}{V_{cell}} \right) \sum_{i=1}^{N_{OS}} \nu_{k,i} \cdot OS_i(1) \quad (38)$$

where, Vel_{Golgi} (Golgi length $\cdot \text{min}^{-1}$) is the linear velocity with which the molecules travel through the Golgi apparatus, $\nu_{k,i}$ ($\text{mmol}_{NSD} \cdot \text{m}^{-1} \cdot \text{mol}_{OS}^{-1}$) is the stoichiometric coefficient of NSD k required to produce the oligosaccharide i and $OS_i(1)$ ($\text{mmol}_{OS} \cdot \text{l}^{-1}$) is the concentration of the oligosaccharide i in the end of the normalized Golgi length as calculated by the N-linked glycosylation model.

3.3 | N-linked glycosylation model

The estimated NSD concentrations are fed to the N-linked mAb glycosylation model that was adapted from Jimenez del val et al., 2011, in order to estimate the mAb glycoform distribution. The adapted model relates the cytosolic concentration of NSDs with the glycan distribution, considering that the concentration of NSDs inside the Golgi apparatus is 20 times greater than the cytosolic concentration (Waldman & Rudnick, 1990). The measured NSD concentration is the total intracellular NSD concentration that accounts both the concentration of NSDs in the cytosolic and Golgi environment, respectively. As proposed by del Val, Polizzi, and Kontoravdi (2016), the current model assumes a constant Golgi volume and a variable linear velocity through Golgi (Vel_{Golgi}), to describe the different rates at which secretory cargo (the mAb) traverses the Golgi apparatus (Presley et al., 1997).

3.4 | Parameter estimation

3.4.1 | Culture and NSD model

The CHO cell culture and NSD metabolic model include 102 unknown parameters, from which 85 parameters were estimated by fitting the model equations to the available experimental data (close to 900 data points in total), using the Parameter Estimation entity in gPROMS v.5.0.1. The remaining 17 parameters were assigned fixed values according to the literature. In order to obtain satisfactory parameter confidence intervals, the model was separated in modular sections and parameter estimation was carried out sequentially as described below: The parameters of the CHO cell culture model and the NSD metabolic model that were not related to galactose and uridine were estimated only on the basis of the control experiment. Using the values defined in the first step, the parameters that were introduced because of galactose and uridine feeding were estimated on the basis of the four feeding experiments. Therefore, no parameter differs between the control and the feeding experiments. Subsequently and as a final step, 18 parameters of the N-linked glycosylation model were estimated as further described in section 3.4.2.

The standard deviation of each experimentally measured variable was used to calculate a variance model for that measurement, which was used in the estimation of related model parameters. In all kinetic

rate expressions, both maximum turnover rates and Monod saturation constants were unknown, making the problem of parameter estimation structurally unidentifiable because the linear dependence between the rate and the substrate concentration below saturation only gives us an estimate of their ratio. Therefore, the majority of Monod constants were hypothesized not to differ more than one order of magnitude from the metabolite or NSD concentration used in each rate, in order to estimate more realistic values for both K_m and V_{max} . However, the estimated values are not intended to accurately represent these kinetic parameters owing to the lack of structural and numerical identifiability.

3.4.2 | N-linked Glycosylation model

The method used for estimating the unknown parameters of the glycosylation model was on the basis of a previously presented method by del Val et al. (2016). However, a slightly modified method was used for the estimation of the GalT distribution (no sialylation was observed). The enzyme parameters were estimated by obtaining the minimum amount of enzyme required to achieve the experimental Galactosylation Index (GI) that is described by Equation (37):

$$GI = \frac{0.5 \Delta[\text{monogalactosylated OS}] + 1 \Delta[\text{digalactosylated OS}]}{\Delta[\text{mAb}]} \quad (39)$$

where, the numerator describes the secreted concentration (mM) of galactosylated mAb glycoforms that were secreted during the chosen interval, and the denominator describes the total secreted mAb during that interval. The calculation of the secreted mAb glycoforms in each interval was on the basis of Fan et al. (2015).

In addition to the estimation of two dissociation constants ($K_{d,i}$) for GalT, the dissociation constants for FucT and GnTI were also estimated, owing to the high concentrations of fucosylated glycans and the low concentrations of Man5 glycans that were observed experimentally. The data from the interval that presented the highest specific mAb productivity was used for $K_{d,i}$ FucT and $K_{d,i}$ GnTI estimation. Considering also the high accumulation of UDPGal, that was observed experimentally during the feeding strategies, the estimation of two $K_{d,k}$ GalT was found to be of great importance. The interval that presented the highest galactosylation was chosen for $\bar{\epsilon}(X, Y) \leq (3/2)(k-1)n$ estimation. The remaining parameters were set to their nominal values (Jimenez del val et al., 2011).

4 | RESULTS

4.1 | Model fitting

The model was simulated for all feeding strategies that were used for model construction and parameter estimation to evaluate the fitting of the model to the training data.

TABLE 1 Correlation coefficients to evaluate the agreement of experimental measurements and model results

Concordance correlation coefficients				
Experiment	CCC _{Xv}	CCC _{mAb}	CCC _{UDPGal}	CCC _{UDPGlcNAc}
Control	0.986	0.998	0.965	0.973
10G	0.892	0.958	0.897	0.919
10G5U	0.960	0.990	0.942	0.968
10G20U	0.741	0.974	0.982	0.941
50G5U	0.971	0.986	0.995	0.990
Predictive experiment	0.959	0.968	-	-
Pearson correlation coefficients				
Experiment	PCC _{Xv}	PCC _{mAb}	PCC _{UDPGal}	PCC _{UDPGlcNAc}
Control	0.990	0.999	0.971	0.985
10G	0.971	0.970	0.930	0.983
10G5U	0.978	0.994	0.949	0.982
10G20U	0.946	0.988	0.989	0.965
50G5U	0.975	0.991	0.997	0.992
Predictive experiment	0.972	0.996	-	-
R squared (R ²)				
Experiment	R ² _{Xv}	R ² _{mAb}	R ² _{UDPGal}	R ² _{UDPGlcNAc}
Control	0.992	0.999	0.998	0.968
10G	0.979	0.968	0.963	0.922
10G5U	0.984	0.995	0.961	0.976
10G20U	0.970	0.988	0.998	0.952
50G5U	0.980	0.992	0.990	0.985
Predictive experiment	0.984	0.984	-	-

Note. mAb: monoclonal antibody; UDPGal: uridine diphosphate galactose.

4.2 | CHO cell culture model simulation

The simulation results present very good agreement with the experimental measurements for all feeding strategies as can be seen in the obtained CCC and R² values in Table 1. Selected estimated kinetic parameters that were used for model simulation are depicted in Table 2 (all parameter values can be found in Table S1). Cell density, ammonia, lactate, and glutamine extra-cellular concentrations match closely with the experimental measurements and indicate that the model framework construction and parameter estimation were successfully conducted as shown in Figure 2 and Figure S1. Importantly, the model captures the metabolic shift from lactate production to consumption in the end of the exponential growth phase and the ammonia increase because of uridine addition. The model is also able to describe the inhibitory impact of galactose feeding on glucose consumption. Finally, product titer is in quantitative agreement for all the feeding experiments indicating that mAb specific production rate follows a linear relationship with the specific growth rate as assumed in Equation (19).

TABLE 2 Selected estimated parameters of the CHO cell culture and NSD metabolic models

Estimated parameter	Value	Units	95% confidence interval
μ_{max}	6.50×10^{-2}	h^{-1}	7.44×10^{-4}
$\mu_{death,max}$	1.50×10^{-2}	h^{-1}	2.56×10^{-3}
K_{Glc}	14.04	mM	5.05×10^{-1}
K_{Asn}	2.62	mM	7.18×10^{-2}
K_{Amm}	3.17	mM	2.49×10^{-1}
K_{Lac}	1×10^3	mM	N/A
K_{Lurd}	41.09	mM	2.42
$K_{d,Amm}$	14.28	mM	3.00
$K_{d,Urd}$	27.86	mM	3.71
$Y_{mAb,X}$	3.39	(pg·cell ⁻¹)	3.21×10^{-1}
m_{mAb}	4.10×10^{-1}	(pg·cell ⁻¹ ·h ⁻¹)	6.57×10^{-3}
V_{max1}	9.22×10^{-1}	(mmol _{NSD} ·L _{cell} ⁻¹ ·h ⁻¹)	1.24×10^{-2}
V_{max2}	1.70×10^{-2}	(mmol _{NSD} ·L _{cell} ⁻¹ ·h ⁻¹)	9.78×10^{-4}
V_{max2b}	59.48	(mmol _{NSD} ·L _{cell} ⁻¹ ·h ⁻¹)	6.10
V_{max3}	5.50×10^{-2}	(mmol _{NSD} ·L _{cell} ⁻¹ ·h ⁻¹)	2.60×10^{-3}
V_{max4}	2.65×10^{-2}	(mmol _{NSD} ·L _{cell} ⁻¹ ·h ⁻¹)	2.12×10^{-3}
V_{max5}	1×10^{-4}	(mmol _{NSD} ·L _{cell} ⁻¹ ·h ⁻¹)	7.80×10^{-6}
V_{max6}	5.13	(mmol _{NSD} ·L _{cell} ⁻¹ ·h ⁻¹)	1.95×10^{-1}
V_{max7}	4.60	(mmol _{NSD} ·L _{cell} ⁻¹ ·h ⁻¹)	4.78×10^{-1}

Note. CHO: Chinese hamster ovary; mAb: monoclonal antibody; NSD: nucleotide sugar donor.

The CCCs, R², and the PCCs were calculated for cell density and mAb concentration in order to quantify the agreement of the model results with the experimental measurements. CCC is used to correlate the values of the model results and the experimental measurements and PCC to correlate the trends of the simulated and experimental curves. The calculated CCC and PCC coefficients and R² are presented in Table 1.

As shown in Table 1, all agreement coefficients for cell density and mAb concentration are above 0.9 with the exception of cell density in the 10G experiment, where CCC is just below 0.9 and the 10G20U experiment, for which the model is unable to capture the steep death phase. The calculated PCCs are higher than the respective CCCs, as expected, and show that the model correctly captures the trend of the examined variables versus time.

4.3 | NSD synthesis model simulation

Simulation results for UDPGal, UDPGlc, UDPGlcNAc, and UDPGalNAc concentrations are presented in Figure 3. The model captured the increase of UDPGal concentration because of galactose feeding when no uridine was added. Moreover, the increase of UDPGal concentration in the experiments that included uridine addition was well described in the simulations. Higher concentrations of UDPGlcNAc, owing to uridine addition, were successfully calculated whereas GDP-Man and GDP-Fuc




FIGURE 2 Model simulations (red line) and experimental measurements (black squares) for the viable cell density, extracellular mAb, glucose, and lactate concentrations. Row a presents the fitting of the model to the control experiment, row b the fitting of the model to the 10G experiment, row c the fitting of the model to the 10G5U experiment, row d the fitting of the model to the 10G20U experiment and row e the fitting of the model to the 50G5U experiment [Color figure can be viewed at wileyonlinelibrary.com]

estimated concentrations were in the same order of the magnitude as the experimental measurements (Figure S2). The significant increase of UDPGalNAc, that is not used as a sugar donor in N-linked glycosylation and UDPGlc, was also described. Regarding the concentration of UDPGlc in the 10G (B) experiment, the model indicated a slight increase of the concentration over time while the experimental results followed a fluctuating trend leading to a 50–75% difference between the simulation and experimental results in the stationary and death phase of the cell

culture. However, the concentration of UDPGlc is fluctuating in a narrow range when compared with the other feeding experiments and, therefore, this discrepancy is not regarded as significant. The estimated maximum turnover rates for the reactions between NSDs as depicted in Figure 1 are displayed in Table 2.

The calculated CCCs, R^2 and PCCs of UDPGal and UDPGlcNAc concentrations shown in Table 1 are above 0.9 for all experiments with the exception of the agreement coefficient of UDPGal




FIGURE 3 Model simulations (red line) and experimental measurements (black squares) for UDPGal, UDPGlc, UDPGlcNAc, and UDPGalNAc concentration. Row a presents the fitting of the model to the control experiment, row b the fitting of the model to the 10G experiment, row c the fitting of the model to the 10G5U experiment, row d the fitting of the model to the 10G20U experiment and row e the fitting of the model to the 50G5U experiment. UDPGal: uridine diphosphate galactose [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1002/biot.25960)]

concentration in the 10G experiment that is 0.897. The high values of the coefficients and R^2 indicate that the NSD model can successfully describe the NSD synthesis pathway and capture the significant alterations caused by precursor feeding.

4.4 | Glycoform distribution

Figure 4 compares the simulated and experimental distributions of G0, G0F, G1F, G2, G2F, and Man5 mAb glycoforms on Days 7, 9, 11, and 12 of cell culture. The majority of glycoform

distributions that were calculated by the N-linked glycosylation model were within $\pm 5\%$ of the corresponding experimental measurements. Only the distribution of the G1F glycoform on Day 7 in the 50 G5U experiment presented a 6.8% underestimation compared with experimental results. Experimentally, specific production rate of galactosylated mAb (Table S3) was reduced alongside with the decreasing cell viability (Figure S3) post galactose and uridine feeding.

The simulation results presented a general underestimation of the experimental measurements probably because of insufficient




FIGURE 4 Model simulation of glycan distribution for all feeding experiments, including the control. Experimental data for the 10G (288 h), 10G20U (216, 264, and 288 h), and 50G5U (288 h) experiments is missing as the respective cell cultures were terminated earlier because of the reduced (below 70%) cell viability

estimated enzyme availability. Enzyme distribution parameters were estimated only according to the control experiment to reduce overfitting. Therefore, the increased availability of GlcNAc and galactose was not depicted in enzyme concentrations and resulted in slight underestimations for some distributions in the feeding experiments. The estimated enzyme distribution parameters and dissociation constants are presented in Table S2. The remaining parameters of the glycosylation model have been assigned to their

nominal values (Jimenez del val et al., 2011). Glycoform distributions could not be subjected to the statistical analysis, as the number of measurements (4) was relatively small.

4.5 | Model predictive capability

A model-based dynamic feeding strategy aimed to maximize galactosylation was implemented experimentally and the data




FIGURE 5 Model prediction and experimental measurements for the optimization experiment used to validate the model. The graph presents the model simulation (red line) and experimental data (black squares) for (a) cell density and (b) mAb concentration while the arrows indicate the days that galactose and uridine were added. Graph (c) depicts the glycan distribution in the optimization experiment, comparing the experimental data with the model predictions for unchanged GalT concentration and for a 15% increase in the concentration of GalT.

Experimental conditions: $V_{\text{feed}} = 10\% V_{\text{culture}}$; $[\text{Gal}]_{\text{feed,day2}} = 0 \text{ mM}$, $[\text{Gal}]_{\text{feed,day4}} = 6.53 \text{ mM}$, $[\text{Gal}]_{\text{feed,day6}} = 9.30 \text{ mM}$, $[\text{Gal}]_{\text{feed,day8}} = 8.98 \text{ mM}$ and $[\text{Gal}]_{\text{feed,day10}} = 8.69 \text{ mM}$; $[\text{Urd}]_{\text{feed,day2}} = 0 \text{ mM}$, $[\text{Urd}]_{\text{feed,day4}} = 0.76 \text{ mM}$, $[\text{Urd}]_{\text{feed,day6}} = 1.34 \text{ mM}$, $[\text{Urd}]_{\text{feed,day8}} = 2.81 \text{ mM}$ and $[\text{Urd}]_{\text{feed,day10}} = 10 \text{ mM}$. GalT: β -1,4-galactosyltransferase; mAb: monoclonal antibody [Color figure can be viewed at wileyonlinelibrary.com]

was used to verify the predictive capabilities of the developed mathematical model. The dynamic optimization was carried out in the relevant function within the gPROMS modeling environment. The objective function we sought to maximize was the concentration of galactosylated mAb species ($\text{mg} \cdot \text{l}^{-1}$), as described in Equation (40).

$$\text{galactosylation} = 1 \times [\text{monogalactosylated mAb}] + 2 \times [\text{digalactosylated mAb}] \quad (40)$$

There were three control variables: the feed flow rate and the concentrations of galactose and uridine in the feed. The problem formulation allowed for the feed to be added or not (binary decision)




FIGURE 6 (a) Experimental degree of galactosylation (%) in the control (blue bars) and the optimization (red bars) experiment. (b) Experimentally measured concentration of galactosylated species ($\text{mg}\cdot\text{L}^{-1}$) in the control (blue bars) and the optimization (red bars) experiment [Color figure can be viewed at wileyonlinelibrary.com]

on alternate days starting on Day 2. The concentration of galactose and uridine in the feed was allowed to range between 0 and 500 mM and 0 and 100 mM, respectively. The resulting feeding strategy includes four pulse additions of varying feed concentrations (details shown in Figure 5). Figure 5 compares model predictions with the measured data for that experiment. Cell density and titer concentration predictions are in good agreement with the experimental measurements.

The model captures the timing of the different growth phases and the impact of galactose and uridine feeding on cell density with sufficient accuracy. Despite the addition of these two precursors, which can negatively affect growth, the cell density remains unaffected compared with the control thanks to the inclusion of these dependencies in the mathematical formulation used for optimization. Glycan distributions measured on Days 7, 9, 11, and 12 are then compared with those predicted by the model (Figure 5). Model accuracy is within $\pm 3.5\%$ with the exception of Day 12 measurement of GOF that is overestimated by 5.3%. The experimentally measured galactosylated-mAb concentration at Day 12 of cell culture exceeds the respective model prediction by 33% because of the unexpected higher total mAb concentration and G1F distribution observed. We hypothesize that the underestimation of the galactosylation percentage is because of intracellular regulation of glycosyltransferase enzymes and/or NSD transporter proteins, which are not included in the model. Indeed, if the concentration of GalT used for model simulation is increased by only 15%, the simulation results are in significantly better agreement with the experimental findings, particularly for GOF and G1F species (Figure 5). The CCCs, R^2 , and PCCs of cell density and mAb concentration presented in Table 1, are above 0.95 indicating that the model successfully predicts cell density and mAb concentration over the culture period. The proposed experimental/modeling platform can, therefore, be used to design optimal NSD precursor feeding strategies that control glycosylation.

To this end, Figure 6 illustrates the impact of the optimized feeding strategy on the percentage and the concentration of the

galactosylated mAb. In both cases, the optimized feeding strategy presents higher galactosylation compared with the control experiment. Unlike the control experiment where the galactosylation percentage decreases through time, the optimized feeding strategy leads to an increasing galactosylation percentage and galactosylated mAb concentration during the cell culture period.

5 | DISCUSSION

A mechanistic mathematical model to describe cell metabolism, mAb production, NSD synthesis, and N-linked glycosylation has been proposed. The model was trained to accumulate data from a wide range of culture conditions in order to capture the responses of all described pathways to different feeding strategies of galactose and uridine addition. Then, the model's predictive capability was successfully validated with an independent experiment. The predictive experiment included different additions of galactose and uridine in a range of concentrations and on different culture days compared with the experiments that were used for model training. The model results for the predictive experiment showed very good agreement with the experimental measurements. The agreement was quantified by calculating the corresponding CCCs and R^2 , which were generally over 0.95 (Table 1). Moreover, the majority of the predicted glycoform distributions deviated up to $\pm 3.5\%$ from the measured data (Figure 4). The high values of all CCCs, R^2 , and PCCs indicate that model successfully reproduces the experimental measurements and can, therefore, be potentially used as a predictive and optimization tool. To the authors' knowledge, this is one of the first studies to dynamically capture variations in mAb glycoform distribution and the first to accurately reproduce and predict the impact of different NSD precursor feeding strategies.

The model suggests that a relationship between the glucose flux through the glycolytic pathway and galactose supplementation exists and is regulated from the fraction of galactose and glucose

consumption rates. Experimental results indicate that galactose addition results in a decreased glucose consumption rate, an effect that has been previously observed in CHO cells (Gramer et al., 2011; Wong et al., 2010). In addition, a simple model for mAb production was used to avoid overparameterization. The results of mAb concentration tracked closely the experimental measurements indicating that mAb production rate is linearly correlated to specific cell growth rate.

The proposed NSD synthesis model was used to calculate the concentrations of the sugar donors that participate in N-linked glycosylation, in particular, UDPGal and UDPGlcNAc, that are extensively affected by the addition of galactose and uridine. Jedrzejewski et al. (2014) attempted to describe a detailed NSD metabolic pathway for hybridoma cells but this led to an extensive and overparameterized reaction network with 34 material balances and 60 reactions. The reduced metabolic network that was used in this modeling approach includes only 7 material balances and 16 reactions, including the effects of galactose and uridine in NSD pools. Monod type equations were used to describe the NSD synthesis reaction rates, including the competitive or noncompetitive inhibitions from other NSDs.

However, the NSD model presents issues of structural unidentifiability and, therefore, the calculated values of the parameters are not a representation of their actual values, specifically with respect to K_{MNSD} . Moreover, the model is limited to a relatively small number of metabolites and amino acids, which, however, are routinely monitored industrially, and could be further expanded to account the majority of amino acids that participate in cell metabolism and the metabolic intermediates of the TCA cycle. As previously mentioned, the occasional deviations in glycan distributions are attributed to the limitations set by the chosen parameter estimation method that does not include the effect of galactose and uridine feeding on the expression of glycosyltransferases and transport proteins of nucleotide sugars (Clark, Griffiths, Bailey, & Harcum, 2005).

Unstructured mathematical models are usually limited to the cell line and experimental data that are used for model construction and parameter estimation. Although the proposed modeling framework can adapt to different feeding concentrations of galactose and uridine, of its applicability to other cell lines, and products remain to be tested. As a first step, a parameter sensitivity analysis can be performed to select the most significant parameters to be re-estimated on the basis of a relatively smaller set of experimental data for a different system.

Despite these drawbacks, the model presented herein successfully captures the impact of galactose and uridine feeding on the secreted mAb glycoform distribution through the use of mechanistic relationships in cell metabolism and NSD synthesis. It captures a wide range of dynamic feeding regimes and was successfully used to design an operating strategy that enhances the percentage of galactosylation up to 64% and the concentration of galactosylated mAb up to 93% with no impact on other key performance indicators, in contrast to the experiments used for the model development in which we saw a significant decrease in viable cell density and, consequently, antibody concentration as a result of galactose and uridine feeding. Therefore, the proposed modeling platform can be a

vehicle for applying the QbD paradigm to optimize the design space for the production of the desired glycan distribution.

ACKNOWLEDGMENTS

P. K. gratefully acknowledges his funding from the PhD scholarship scheme of the Department of Chemical Engineering, Imperial College London. P. M. J. and S. N. S. thank the Engineering and Physical Sciences Research Council and the Biotechnology and Biological Sciences Research Council, respectively, for their studentships. The authors thank MedImmune for their funding, providing the cell line used in this study, as well as access to their analytical facilities.

ORCID

Karen Polizzi  <http://orcid.org/0000-0001-5435-2667>

Ioscani Jimenez del Val  <http://orcid.org/0000-0001-9289-0191>

Cleo Kontoravdi  <http://orcid.org/0000-0003-0213-4830>

REFERENCES

- Clark, K. J. R., Griffiths, J., Bailey, K. M., & Harcum, S. W. (2005). Gene-expression profiles for five key glycosylation genes for galactose-fed CHO cells expressing recombinant IL-4/13 cytokine trap. *Biotechnology and Bioengineering*, 90(5), 568–577.
- Fan, Y., Jimenez Del Val, I., Müller, C., Lund, A. M., Sen, J. W., Rasmussen, S. K., ... Andersen, M. R. (2015). A multi-pronged investigation into the effect of glucose starvation and culture duration on fed-batch CHO cell culture. *Biotechnology and Bioengineering*, 112(10), 2172–2184.
- Goetze, A. M., Liu, Y. D., Zhang, Z., Shah, B., Lee, E., Bondarenko, P. V., & Flynn, G. C. (2011). High-mannose glycans on the Fc region of therapeutic IgG antibodies increase serum clearance in humans. *Glycobiology*, 21(7), 949–959.
- Grainger, R. K., & James, D. C. (2013). CHO cell line specific prediction and control of recombinant monoclonal antibody N-glycosylation. *Biotechnology and Bioengineering*, 110(11), 2970–2983.
- Gramer, M. J., Eckblad, J. J., Donahue, R., Brown, J., Shultz, C., Vickerman, K., ... van Berkel, P. H. C. (2011). Modulation of antibody galactosylation through feeding of uridine, manganese chloride, and galactose. *Biotechnology and Bioengineering*, 108(7), 1591–1602.
- Hodoniczky, J., Zheng, Y. Z., & James, D. C. (2005). Control of recombinant monoclonal antibody effector functions by Fc N-glycan remodeling in vitro. *Biotechnology Progress*, 21(6), 1644–1652.
- Hutter, S., Villiger, T. K., Brühlmann, D., Stettler, M., Broly, H., Soos, M., & Gunawan, R. (2017). Glycosylation flux analysis reveals dynamic changes of intracellular glycosylation flux distribution in Chinese hamster ovary fed-batch cultures. *Metabolic Engineering*, 43, 9–20.
- Jedrzejewski, P., del Val, I., Constantinou, A., Dell, A., Haslam, S., Polizzi, K., & Kontoravdi, C. (2014). Towards controlling the glycoform: A model framework linking extracellular metabolites to antibody glycosylation. *International Journal of Molecular Sciences*, 15(3), 4492–4522.
- Kanehisa, M., & Goto, S. (2000). KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Research*, 28(1), 27–30.
- Kaveh, O., Hengameh, A., Johannes, G., Murray, M. Y., Raymond, L. L., Jen, S., & Hector, B. M. (2013). Novel dynamic model to predict the glycosylation pattern of monoclonal antibodies from extracellular cell culture conditions. *IFAC Proceedings Volumes*, 46(31), 30–35.
- Kontoravdi, C., et al. (2010). Systematic development of predictive mathematical models for animal cell cultures. *Computers & Chemical Engineering*, 34, 1192–1198.

- Kontoravdi, C., & Jimenez del val, I. (2018). Computational tools for predicting and controlling the glycosylation of biopharmaceuticals. *Current Opinion in Chemical Engineering*, 22, 89–97.
- Krambeck, F. J., & Betenbaugh, M. J. (2005). A mathematical model of N-linked glycosylation. *Biotechnology and Bioengineering*, 92(6), 711–728.
- Kremkow, B. G., & Lee, K. H. (2018). Glyco-mapper: A Chinese hamster ovary (CHO) genome-specific glycosylation prediction tool. *Metabolic Engineering*, 47, 134–142.
- Lai, T., Yang, Y., & Ng, S. (2013). Advances in mammalian cell line development technologies for recombinant protein production. *Pharmaceuticals*, 6(5), 579–603.
- Lin, L. I. K. (1989). A concordance correlation coefficient to evaluate reproducibility. *Biometrics*, 45(1), 255–268.
- Liu, B., Spearman, M., Doering, J., Lattová, E., Perreault, H., & Butler, M. (2014). The availability of glucose to CHO cells affects the intracellular lipid-linked oligosaccharide distribution, site occupancy and the N-glycosylation profile of a monoclonal antibody. *Journal of Biotechnology*, 170, 17–27.
- Nolan, R. P., & Lee, K. (2011). Dynamic model of CHO cell metabolism. *Metabolic Engineering*, 13(1), 108–124.
- Presley, J. F., Cole, N. B., Schroer, T. A., Hirschberg, K., Zaal, K. J. M., & Lippincott-Schwartz, J. (1997). ER-to-Golgi transport visualized in living cells. *Nature*, 389, 81–85.
- Schomburg, I., et al. (2002). BRENDA, enzyme data and metabolic information. *Nucleic Acids Research*, 30(1), 47–49.
- Shinkawa, T., Nakamura, K., Yamane, N., Shoji-Hosaka, E., Kanda, Y., Sakurada, M., ... Shitara, K. (2003). The absence of fucose but not the presence of galactose or bisecting N-acetylglucosamine of human IgG1 complex-type oligosaccharides shows the critical role of enhancing antibody-dependent cellular cytotoxicity. *Journal of Biological Chemistry*, 278(5), 3466–3473.
- Sou, S. N., Sellick, C., Lee, K., Mason, A., Kyriakopoulos, S., Polizzi, K. M., & Kontoravdi, C. (2015). How does mild hypothermia affect monoclonal antibody glycosylation? *Biotechnology and Bioengineering*, 112(6), 1165–1176.
- Spahn, P. N., Hansen, A. H., Hansen, H. G., Arnsdorf, J., Kildegaard, H. F., & Lewis, N. E. (2016). A Markov chain model for N-linked protein glycosylation – towards a low-parameter tool for model-driven glycoengineering. *Metabolic Engineering*, 33, 52–66.
- Thomann, M., Reckermann, K., Reusch, D., Prasser, J., & Tejada, M. L. (2016). Fc-galactosylation modulates antibody-dependent cellular cytotoxicity of therapeutic antibodies. *Molecular Immunology*, 73, 69–75.
- Urquhart, L. (2018). Market watch: Top drugs and companies by sales in 2017. *Nature Reviews. Drug Discovery*, 17(4), 232–232.
- Jimenez del val, I., Nagy, J. M., & Kontoravdi, C. (2011). A dynamic mathematical model for monoclonal antibody N-linked glycosylation and nucleotide sugar donor transport within a maturing Golgi apparatus. *Biotechnology Progress*, 27(6), 1730–1743.
- del Val, I. J., Kyriakopoulos, S., Polizzi, K. M., & Kontoravdi, C. (2013). An optimized method for extraction and quantification of nucleotides and nucleotide sugars from mammalian cells. *Analytical Biochemistry*, 443(2), 172–180.
- Jimenez del val, I., Fan, Y., & Weilguny, D. (2016). Dynamics of immature mAb glycoform secretion during CHO cell culture: An integrated modelling framework. *Biotechnology Journal*, 11(5), 610–623.
- del Val, I. J., Polizzi, K. M., & Kontoravdi, C. (2016). A theoretical estimate for nucleotide sugar demand towards Chinese hamster ovary cellular glycosylation. *Scientific Reports*, 6, 28547.
- Waldman, B. C., & Rudnick, G. (1990). UDP-GlcNAc transport across the Golgi membrane: Electroneutral exchange for dianionic UMP. *Biochemistry*, 29(1), 44–52.
- Walsh, G. (2014). Biopharmaceutical benchmarks 2014. *Nature Biotechnology*, 32, 992–1000.
- Wieczorke, R., Dlugai, S., Krampe, S., & Boles, E. (2003). Characterisation of mammalian GLUT glucose transporters in a heterologous yeast expression system. *Cellular Physiology and Biochemistry*, 13(3), 123–134.
- Wong, N. S. C., Wati, L., Nissom, P. M., Feng, H. T., Lee, M. M., & Yap, M. G. S. (2010). An investigation of intracellular glycosylation activities in CHO cells: Effects of nucleotide sugar precursor feeding. *Biotechnology and Bioengineering*, 107(2), 321–336.
- Yang, M., & Butler, M. (2000). Effects of ammonia on CHO cell growth, erythropoietin production, and glycosylation. *Biotechnology and Bioengineering*, 68(4), 370–380.
- Yu, M., Brown, D., Reed, C., Chung, S., Lutman, J., Stefanich, E., ... Bayer, R. (2012). Production, characterization and pharmacokinetic properties of antibodies with N-linked mannose-5 glycans. *mAbs*, 4(4), 475–487.
- Yusufi, F. N. K., Lakshmanan, M., Ho, Y. S., Loo, B. L. W., Ariyaratne, P., Yang, Y., ... Lee, D. Y. (2017). Mammalian systems biotechnology reveals global cellular adaptations in a recombinant CHO cell line. *Cell Systems*, 4(5), 530–542. e536
- Zagari, F., Jordan, M., Stettler, M., Broly, H., & Wurm, F. M. (2013). Lactate metabolism shift in CHO cell culture: The role of mitochondrial oxidative activity. *New Biotechnology*, 30(2), 238–245.
- Zhao, F.-Q., & Keating, A. (2007). Functional properties and genomics of glucose transporters. *Current Genomics*, 8(2), 113–128.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Kotidis P, Jedrzejewski P, Sou SN, et al. Model-based optimization of antibody galactosylation in CHO cell culture. *Biotechnology and Bioengineering*. 2019;116: 1612–1626. <https://doi.org/10.1002/bit.26960>