

Open Quantum System

Lectures:

Last time:

- ▶ Daniel Manzano, A short introduction to the Lindblad master equation (*all*)
- ▶ Breuer and Petruccione, The Theory of Open Quantum Systems (*ch. 3 - 4.3*)
- ▶ Daniel A. Lidar, Notes on the Theory of Open Quantum Systems (*up to ch. 12*)

Today:

- ▶ Buča, B., Tindall, J. & Jaksch, D. Non-stationary coherent quantum many-body dynamics through dissipation
- ▶ Victor V. Albert & Liang Jiang, Symmetries and conserved quantities in Lindblad master equations
- ▶ Cameron Booker, Berislav Buča, Dieter Jaksch, Non-stationarity and Dissipative Time Crystals: Spectral Properties and Finite-Size Effects
- ▶ Victor V. Albert. Lindbladians with multiple steady states: theory and applications. A Dissertation Presented to the Faculty of the Graduate School of Yale University

November 20, 2023

Dynamical symmetries

Diagonal Lindblad

$$\mathcal{L}\rho = -i[H_S + H_{LS}, \rho] + \sum_{\mu} \left(2L_{\mu}\rho L_{\mu}^{\dagger} - \{L_{\mu}^{\dagger}L_{\mu}, \rho\} \right) \quad \text{with } L_{\mu} \equiv L_{\nu}(\omega)$$

Dynamical symmetries

Diagonal Lindblad

$$\mathcal{L}\rho = -i[H_S + H_{LS}, \rho] + \sum_{\mu} \left(2L_{\mu}\rho L_{\mu}^{\dagger} - \{L_{\mu}^{\dagger}L_{\mu}, \rho\} \right) \quad \text{with } L_{\mu} \equiv L_{\nu}(\omega)$$

Dynamical symmetry

$$\forall \mu \equiv (\nu, \omega), \quad \underbrace{[L_{\mu}, A]}_{L_{\nu}(\omega)} = 0 \quad \text{with } L_{\nu}(\omega) := \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_{\nu} |\epsilon'\rangle\langle\epsilon'|$$

Dynamical symmetries

Diagonal Lindblad

$$\mathcal{L}\rho = -i[H_S + H_{LS}, \rho] + \sum_{\mu} \left(2L_{\mu}\rho L_{\mu}^{\dagger} - \{L_{\mu}^{\dagger}L_{\mu}, \rho\} \right) \quad \text{with } L_{\mu} \equiv L_{\nu}(\omega)$$

Dynamical symmetry

$$\forall \mu \equiv (\nu, \omega), \quad \underbrace{[L_{\mu}, A]}_{L_{\nu}(\omega)} = 0 \quad \text{with } L_{\nu}(\omega) := \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_{\nu} |\epsilon'\rangle\langle\epsilon'|$$

Remembering

$$H_{\text{int}} \equiv \sum_{\alpha} S_{\alpha} \otimes B_{\alpha} \quad \text{and} \quad L_{\nu} = \sum_i u_{i,\nu} S_i$$

Dynamical symmetries

Commutator

Re-writing commutator - $[L_\nu(\omega), A] = 0 \quad \forall (\nu, \omega)$

Dynamical symmetries

Commutator

Re-writing commutator - $[L_\nu(\omega), A] = 0 \quad \forall (\nu, \omega)$

Giving for a specific ω and ν

$$\left[\sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'|, A \right] = 0 \quad \forall (\nu, \omega) \quad (1)$$

$$\left[\sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_\nu^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0 \quad \forall (\nu, \omega) \quad (2)$$

Dynamical symmetries

Commutator

Re-writing commutator - $[L_\nu(\omega), A] = 0 \quad \forall (\nu, \omega)$

Giving for a specific ω and ν

$$\left[\sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'|, A \right] = 0 \quad \forall (\nu, \omega) \quad (1)$$


$$\left[\sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_\nu^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0 \quad \forall (\nu, \omega) \quad (2)$$

Question: what can one say on A and/or L_ν ?

Dynamical symmetries

Assumption (a) - for every ω , a unique pair (ϵ, ϵ') exists.

Note - assumption **not** the same as non-degeneracy



Dynamical symmetries


Assumption (a) - for every ω , a unique pair (ϵ, ϵ') exists.

Note - assumption **not** the same as non-degeneracy

Then, for a chosen (ν, ϵ) , one writes

$$\left[|\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'|, A \right] = 0 \quad \forall (\nu, (\epsilon', \epsilon)) \quad (3)$$

$$\left[|\epsilon'\rangle\langle\epsilon'| L_\nu^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0 \quad \forall (\nu, (\epsilon', \epsilon)) \quad (4)$$



Dynamical symmetries


Assumption (a) - for every ω , a unique pair (ϵ, ϵ') exists.

Note - assumption **not** the same as non-degeneracy

Then, for a chosen (ν, ϵ) , one writes

$$\left[|\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'|, A \right] = 0 \quad \forall (\nu, (\epsilon', \epsilon)) \quad (3)$$

$$\left[|\epsilon'\rangle\langle\epsilon'| L_\nu^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0 \quad \forall (\nu, (\epsilon', \epsilon)) \quad (4)$$



The states $\{|\epsilon\rangle\}$ - the chosen orthonormal eigenkets of the Hamiltonian H_S .
Clearly, the ket-bra's span the Liouville space -

$$\mathfrak{D}(\mathcal{H}) = \text{span}(\{|\epsilon\rangle\langle\epsilon'|\}) \quad \forall (\nu, (\epsilon', \epsilon))$$

Assumption $\textcircled{\mathfrak{a}}$

Using the completeness relation,

$$L_\nu = \sum_{\varepsilon, \varepsilon'} |\varepsilon\rangle\langle\varepsilon| L_\nu |\varepsilon'\rangle\langle\varepsilon'|$$

For all $(\nu, (\varepsilon, \varepsilon'))$, one writes

Assumption \textcircled{a}

Using the completeness relation,

$$L_\nu = \sum_{\varepsilon, \varepsilon'} |\varepsilon\rangle\langle\varepsilon| L_\nu |\varepsilon'\rangle\langle\varepsilon'|$$

For all $(\nu, (\varepsilon, \varepsilon'))$, one writes

$$\begin{aligned} L_\nu A - A L_\nu &= \sum_{\varepsilon, \varepsilon'} |\varepsilon\rangle\langle\varepsilon| L_\nu |\varepsilon'\rangle\langle\varepsilon'| A - \sum_{\varepsilon, \varepsilon'} A |\varepsilon\rangle\langle\varepsilon| L_\nu |\varepsilon'\rangle\langle\varepsilon'| \\ &= \sum_{\varepsilon} \sum_{\varepsilon'} \underbrace{\left(|\varepsilon\rangle\langle\varepsilon| L_\nu |\varepsilon'\rangle\langle\varepsilon'| A - A |\varepsilon\rangle\langle\varepsilon| L_\nu |\varepsilon'\rangle\langle\varepsilon'| \right)}_{0 \text{ for all } (\nu, (\varepsilon, \varepsilon'))} = 0 \end{aligned} \quad (5)$$

Assumption \textcircled{a}

Using the completeness relation,

$$L_\nu = \sum_{\epsilon, \epsilon'} |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'|$$

For all $(\nu, (\epsilon, \epsilon'))$, one writes

$$\begin{aligned} L_\nu A - A L_\nu &= \sum_{\epsilon, \epsilon'} |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'| A - \sum_{\epsilon, \epsilon'} A |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'| \\ &= \sum_{\epsilon} \sum_{\epsilon'} \underbrace{\left(|\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'| A - A |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'| \right)}_{0 \text{ for all } (\nu, (\epsilon, \epsilon'))} = 0 \end{aligned} \quad (5)$$

The same process can be repeated for L_ν^\dagger .

Meaning that if such dynamical symmetry exists,

$$[L_\nu, A] = [L_\nu^\dagger, A] = 0$$

Assumption \textcircled{a}

Can one show that $A = \lambda \mathbb{I}$?

Assumption ①

Can one show that $A = \lambda \mathbb{I}$?

Let the most general form of A :

$$A = \sum_{kl} a_{kl} |k\rangle\langle l|$$

Assumption ①

Can one show that $A = \lambda \mathbb{I}$?

Let the most general form of A :

$$A = \sum_{kl} a_{kl} |k\rangle\langle l|$$

One wants to show that $a_{kl} = 0$ for $k \neq l$. Suppose the contrary - there exists $a_{kl} \neq 0$ for $k \neq l$

Assumption \textcircled{a}

Can one show that $A = \lambda \mathbb{I}$?

Let the most general form of A:

$$A = \sum_{kl} a_{kl} |k\rangle\langle l|$$

One wants to show that $a_{kl} = 0$ for $k \neq l$. Suppose the contrary - there exists $a_{kl} \neq 0$ for $k \neq l$

Then, the commutator relation will be given by

$$\left[|m\rangle\langle m| L_\nu |n\rangle\langle n|, \sum_{kl} a_{kl} |k\rangle\langle l| \right] = 0 \quad (6)$$

$$\sum_{kl} a_{kl} |m\rangle\langle m| L_\nu |n\rangle\langle n| |k\rangle\langle l| - \sum_{kl} a_{kl} |k\rangle\langle l| |m\rangle\langle m| L_\nu |n\rangle\langle n| \quad (7)$$

Assumption $\textcircled{\mathfrak{a}}$

$$\sum_{kl} a_{kl} |m\rangle\langle m| L_{\nu} |n\rangle\langle n| |k\rangle\langle l| - \sum_{kl} a_{kl} |k\rangle\langle l| |m\rangle\langle m| L_{\nu} |n\rangle\langle n| =$$

$$\sum_l a_{nl} |m\rangle\langle m| L_{\nu} |n\rangle\langle l| - \sum_k a_{km} |k\rangle\langle m| L_{\nu} |n\rangle\langle n|$$

Assumption $\textcircled{\mathfrak{a}}$

$$\sum_{kl} a_{kl} |m\rangle\langle m| L_\nu |n\rangle\langle n| |k\rangle\langle l| - \sum_{kl} a_{kl} |k\rangle\langle l| |m\rangle\langle m| L_\nu |n\rangle\langle n| =$$

$$\sum_l a_{nl} |m\rangle\langle m| L_\nu |n\rangle\langle l| - \sum_k a_{km} |k\rangle\langle m| L_\nu |n\rangle\langle n|$$

$$\xrightarrow{|m\rangle} a_{nm} |m\rangle\langle m| L_\nu |n\rangle = 0 \quad (8)$$

$$\xrightarrow{\langle n|} a_{nm} \langle m| L_\nu |n\rangle\langle n| = 0 \quad (9)$$

Assumption \textcircled{a}

$$\sum_{kl} a_{kl} |m\rangle\langle m| L_\nu |n\rangle\langle n| |k\rangle\langle l| - \sum_{kl} a_{kl} |k\rangle\langle l| |m\rangle\langle m| L_\nu |n\rangle\langle n| =$$

$$\sum_l a_{nl} |m\rangle\langle m| L_\nu |n\rangle\langle l| - \sum_k a_{km} |k\rangle\langle m| L_\nu |n\rangle\langle n|$$

$$\xrightarrow{|m\rangle} a_{nm} |m\rangle\langle m| L_\nu |n\rangle = 0 \quad (8)$$

$$\xrightarrow{\langle n|} a_{nm} \langle m| L_\nu |n\rangle\langle n| = 0 \quad (9)$$

► Must hold for all ν

Assumption $\textcircled{\mathfrak{a}}$

$$\sum_{kl} a_{kl} |m\rangle\langle m| L_\nu |n\rangle\langle n| |k\rangle\langle l| - \sum_{kl} a_{kl} |k\rangle\langle l| |m\rangle\langle m| L_\nu |n\rangle\langle n| =$$

$$\sum_l a_{nl} |m\rangle\langle m| L_\nu |n\rangle\langle l| - \sum_k a_{km} |k\rangle\langle m| L_\nu |n\rangle\langle n|$$

$$\xrightarrow{|m\rangle} a_{nm} |m\rangle\langle m| L_\nu |n\rangle = 0 \quad (8)$$

$$\xrightarrow{\langle n|} a_{nm} \langle m| L_\nu |n\rangle\langle n| = 0 \quad (9)$$

- ▶ Must hold for all ν
- ▶ L_ν was defined to be $H_{\text{int}} \simeq \sum_\chi L_\chi \otimes B_\chi$, meaning that

$$L_\nu \in \text{span}(|\epsilon\rangle\langle\epsilon'|)$$

Assumption (a)

$$\sum_{kl} a_{kl} |m\rangle\langle m| L_\nu |n\rangle\langle n| |k\rangle\langle l| - \sum_{kl} a_{kl} |k\rangle\langle l| |m\rangle\langle m| L_\nu |n\rangle\langle n| =$$

$$\sum_l a_{nl} |m\rangle\langle m| L_\nu |n\rangle\langle l| - \sum_k a_{km} |k\rangle\langle m| L_\nu |n\rangle\langle n|$$

$$\xrightarrow{|m\rangle} a_{nm} |m\rangle\langle m| L_\nu |n\rangle = 0 \quad (8)$$

$$\xrightarrow{\langle n|} a_{nm} \langle m| L_\nu |n\rangle\langle n| = 0 \quad (9)$$

- ▶ Must hold for all ν
- ▶ L_ν was defined to be $H_{\text{int}} \simeq \sum_\chi L_\chi \otimes B_\chi$, meaning that

$$L_\nu \in \text{span}(|\epsilon\rangle\langle\epsilon'|)$$
- ▶ Meaning that $a_{nm} = 0 \implies$ contradiction

Assumption $\textcircled{\mathfrak{a}}$

One can show that $A = \lambda \mathbb{I}$, by showing that $a_{jj} = a_{kk} \equiv d_k$.

Assumption \textcircled{a}

One can show that $A = \lambda \mathbb{I}$, by showing that $a_{jj} = a_{kk} \equiv d_k$. Since $A \sim \text{diag.}$,

$$A = \sum_k d_k |k\rangle\langle k|$$

Assumption ①

One can show that $A = \lambda \mathbb{I}$, by showing that $a_{jj} = a_{kk} \equiv d_k$. Since $A \sim \text{diag.}$,

$$A = \sum_k d_k |k\rangle\langle k|$$

Using the same expansion, and commutator relations -

$$d_\epsilon |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'| - d_{\epsilon'} |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'| = 0$$

Meaning that diagonal elements are the same \implies

$$A = \lambda \mathbb{I}$$

Assumption ①

One can show that $A = \lambda \mathbb{I}$, by showing that $a_{jj} = a_{kk} \equiv d_k$. Since $A \sim \text{diag.}$,

$$A = \sum_k d_k |k\rangle\langle k|$$

Using the same expansion, and commutator relations -

$$d_\epsilon |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'| - d_{\epsilon'} |\epsilon\rangle\langle\epsilon| L_\nu |\epsilon'\rangle\langle\epsilon'| = 0$$

Meaning that diagonal elements are the same \implies

$$A = \lambda \mathbb{I}$$

The newly introduced $\rho_{mn} := A^n \rho_\infty (A^\dagger)^m$ thus do not give any additional information. The condition $\mathcal{L} \rho_{mn} = -i(m-n)\lambda \rho_{mn}$ therefore is redundant.

Assumption: continuity

- The discrete spectrum:

$$S_{\alpha}(\omega) := \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| S_{\alpha} |\epsilon'\rangle\langle\epsilon'|$$

Assumption: continuity

- ▶ The discrete spectrum:


$$S_{\alpha}(\omega) := \sum_{\omega = \epsilon' - \epsilon} |\epsilon\rangle\langle\epsilon| S_{\alpha} |\epsilon'\rangle\langle\epsilon'|$$

- ▶ Continuous spectrum:

$$S_{\alpha}(\omega) = \iint_{\epsilon, \epsilon' \in \sigma(H_S)} |\epsilon\rangle\langle\epsilon| S_{\alpha} |\epsilon'\rangle\langle\epsilon'| \delta(\omega - (\epsilon - \epsilon')) d\epsilon d\epsilon' \quad (10)$$


Degeneracies?

- In addition to the assumption (a), one can add degeneracies to the ground level $|d_1\rangle, |d_2\rangle$.




Degeneracies?

- ▶ In addition to the assumption (a), one can add degeneracies to the ground level $|d_1\rangle, |d_2\rangle$.
- ▶ What does this mean for the commutators?
- ▶ Additional terms at $\omega = \epsilon' - \epsilon$ for $0 = \omega = d_1 - d_2$



Degeneracies?


- ▶ In addition to the assumption (a), one can add degeneracies to the ground level $|d_1\rangle, |d_2\rangle$.
- ▶ What does this mean for the commutators?
- ▶ Additional terms at $\omega = \epsilon' - \epsilon$ for $0 = \omega = d_1 - d_2$



$$\left[|d_1\rangle\langle d_1| L_\nu |d_2\rangle\langle d_2| + |d_2\rangle\langle d_2| L_\nu |d_1\rangle\langle d_1|, \underbrace{A}_{\equiv a_{ij}} \right] = 0$$

Degeneracies?

- ▶ In addition to the assumption (a), one can add degeneracies to the ground level $|d_1\rangle, |d_2\rangle$.
- ▶ What does this mean for the commutators?
- ▶ Additional terms at $\omega = \epsilon' - \epsilon$ for $0 = \omega = d_1 - d_2$



$$\left[|d_1\rangle\langle d_1| L_\nu |d_2\rangle\langle d_2| + |d_2\rangle\langle d_2| L_\nu |d_1\rangle\langle d_1|, \underbrace{A}_{\equiv a_{ij}} \right] = 0$$

Using same procedures as before, one can obtain some relations as for example

$$a_{d_1, d_2} \langle d_2 | L_\nu | d_1 \rangle - a_{d_2, d_1} \langle d_1 | L_\nu | d_2 \rangle = 0$$

$$a_{d_1, d_1} \langle d_2 | L_\nu | d_1 \rangle - a_{d_2, d_2} \langle d_2 | L_\nu | d_1 \rangle = 0$$

Degeneracies

This suggest the form of A

$$\begin{pmatrix} \mathbf{A}_0^* & \begin{smallmatrix} \cdot & \cdot & \cdot \\ * & & \\ \cdot & \cdot & \cdot \end{smallmatrix} \\ \begin{smallmatrix} \cdot & \cdot & \cdot \\ * & & \\ \cdot & \cdot & \cdot \end{smallmatrix} & \lambda \mathbf{I} \end{pmatrix}$$

More on symmetries

- ▶ In fact, the conditions $[J, H] = 0$, $\dot{J} = 0$ and $U^\dagger H U = H$ are **all** equivalent.

More on symmetries

- ▶ In fact, the conditions $[J, H] = 0$, $\dot{J} = 0$ and $U^\dagger H U = H$ are **all** equivalent.
- ▶ For systems under \mathcal{L} , one introduces super-operators \mathcal{U} and \mathcal{J} , associated to U and J resp.

More on symmetries

- ▶ In fact, the conditions $[J, H] = 0$, $\dot{J} = 0$ and $U^\dagger H U = H$ are **all** equivalent.
- ▶ For systems under \mathcal{L} , one introduces super-operators \mathcal{U} and \mathcal{J} , associated to U and J resp.
- ▶ The relations are weaker and given by

$$\dot{J} \equiv \mathcal{L}(J) \iff [J, H] = [J, L_\nu] = 0 \implies [\mathcal{L}, \mathcal{U}] = 0$$

More on symmetries

- ▶ In fact, the conditions $[J, H] = 0$, $\dot{J} = 0$ and $U^\dagger H U = H$ are **all** equivalent.
- ▶ For systems under \mathcal{L} , one introduces super-operators \mathcal{U} and \mathcal{J} , associated to U and J resp.
- ▶ The relations are weaker and given by

$$\dot{J} \equiv \mathcal{L}(J) \iff [J, H] = [J, L_\nu] = 0 \implies [\mathcal{L}, \mathcal{U}] = 0$$

In fact, if \mathcal{L} does **not** have purely imaginary eigenvalues (no oscillating coherences), then for $\rho_{ss} \in \mathcal{L}_{ss}$ of $\dim = D$

$$\rho_{ss} = \lim_{t \rightarrow \infty} e^{\mathcal{L}t} \rho_{in} = \sum_{\mu}^D \langle J_{\mu} | \rho_{in} \rangle M_{\mu} \quad (11)$$

Steady-states vs Oscillating coherences

- ▶ Conserved quantities for ρ_{ss} are not conserved anymore for ρ_{∞} .

Steady-states vs Oscillating coherences

- ▶ Conserved quantities for ρ_{ss} are not conserved anymore for ρ_{∞} .
- ▶ It is possible to re-write similar explicit relations by considering other "pseudo-conserved quantities" $J \mapsto S$ together with H_{∞} , that is implicitly included in the H in the master equation.

Steady-states vs Oscillating coherences

- ▶ Conserved quantities for ρ_{ss} are not conserved anymore for ρ_{∞} .
- ▶ It is possible to re-write similar explicit relations by considering other "pseudo-conserved quantities" $J \mapsto S$ together with H_{∞} , that is implicitly included in the H in the master equation.

Q

How are those oscillating coherence quantities related to each other and to the steady-state quantities?