

Open Quantum System

Lectures:

Last time:

- ▶ Daniel Manzano, A short introduction to the Lindblad master equation (*all*)
- ▶ Breuer and Petruccione, The Theory of Open Quantum Systems (*ch. 3 - 4.3*)
- ▶ Daniel A. Lidar, Notes on the Theory of Open Quantum Systems (*up to ch. 12*)

Today:

- ▶ Breuer and Petruccione, The Theory of Open Quantum Systems (*ch. 3 - 4.3*)
- ▶ Daniel A. Lidar, Notes on the Theory of Open Quantum Systems (*up to ch. 12*)
- ▶ B.Kraus, H.P.Buchler, S. Diehl, A. Kantian, A. Micheli, & P. Zoller, Preparation of Entangled States by Quantum Markov Processes
- ▶ Buča, B., Tindall, J. & Jaksch, D. Non-stationary coherent quantum many-body dynamics through dissipation
- ▶ Victor V. Albert & Liang Jiang, Symmetries and conserved quantities in Lindblad master equations
- ▶ Cameron Booker, Berislav Buča, Dieter Jaksch, Non-stationarity and Dissipative Time Crystals: Spectral Properties and Finite-Size Effects

November 5, 2023

Last time

- Derivation of the Lindblad equation via different approximations:

Last time

- ▶ Derivation of the Lindblad equation via different approximations:
 - ▶ Von Neumann evolution equation, $H_{\text{int}} = \sum_k S_k \otimes B_k \rightarrow$ Weak coupling, Born, Markov & Rotating wave \rightarrow Redfield equation
Note: no stationarity of the system was assumed!

Last time

- Derivation of the Lindblad equation via different approximations:
 - Von Neumann evolution equation, $H_{\text{int}} = \sum_k S_k \otimes B_k \rightarrow$ Weak coupling, Born, Markov & Rotating wave \rightarrow Redfield equation
Note: no stationarity of the system was assumed!
 - Yielding (Schrodinger pic.):

$$\begin{aligned} \frac{d}{dt}\rho(t) = [H + H_{\text{LS}}, \rho(t)] + \\ + \sum_{k,l,\omega} \gamma_{k,l}(\omega) \left[S_l(\omega) \rho(t) S_k^\dagger(\omega) - \frac{1}{2} \{ S_k^\dagger(\omega) S_l(\omega), \rho(t) \} \right] \end{aligned} \quad (1)$$

Last time

- Derivation of the Lindblad equation via different approximations:
 - Von Neumann evolution equation, $H_{\text{int}} = \sum_k S_k \otimes B_k \rightarrow$ Weak coupling, Born, Markov & Rotating wave \rightarrow Redfield equation
Note: no stationarity of the system was assumed!
 - Yielding (Schrodinger pic.):

$$\begin{aligned} \frac{d}{dt}\rho(t) = [H + H_{\text{LS}}, \rho(t)] + \\ + \sum_{k,l,\omega} \gamma_{k,l}(\omega) \left[S_l(\omega) \rho(t) S_k^\dagger(\omega) - \frac{1}{2} \{ S_k^\dagger(\omega) S_l(\omega), \rho(t) \} \right] \end{aligned} \quad (1)$$

- With $\gamma_{k,l}$, $\pi_{k,l}$ - defined via $\Gamma_{k,l}(\omega)$'s real and imaginary parts.

$$\begin{aligned} H_{\text{LS}} &= \sum_{\omega,k,l} \pi_{k,l}(\omega) S_k^\dagger(\omega) S_l(\omega) \\ \Gamma_{k,l}(\omega) &= \int_0^\infty e^{i\omega s} \text{Tr}_B [B_k^\dagger(t) B_l(t-s) \rho_B(0)] \end{aligned} \quad (2)$$

- With the operators $S_k(\omega)$ defined via

$$S_k(\omega) = \sum_{\epsilon' - \epsilon = \omega} \Pi_{\epsilon} S_k \Pi_{\epsilon'} \equiv \sum_{\epsilon' - \epsilon = \omega} |\epsilon\rangle\langle\epsilon| S_k |\epsilon'\rangle\langle\epsilon'| \quad (3)$$

- With the operators $S_k(\omega)$ defined via

$$S_k(\omega) = \sum_{\epsilon' - \epsilon = \omega} \Pi_{\epsilon} S_k \Pi_{\epsilon'} \equiv \sum_{\epsilon' - \epsilon = \omega} |\epsilon\rangle\langle\epsilon| S_k |\epsilon'\rangle\langle\epsilon'| \quad (3)$$

- Yielding a time evolution (Dirac's pic) in the form

$$H_{\text{int}}(t) = \sum_{\alpha, \omega} e^{-i\omega t} S_{\alpha}(\omega) \otimes B_{\alpha}(t), \quad B_{\alpha}(t) = e^{iH_B t} B_{\alpha} e^{-iH_B t} \quad (4)$$

- With the operators $S_k(\omega)$ defined via

$$S_k(\omega) = \sum_{\epsilon' - \epsilon = \omega} \Pi_{\epsilon} S_k \Pi_{\epsilon'} \equiv \sum_{\epsilon' - \epsilon = \omega} |\epsilon\rangle\langle\epsilon| S_k |\epsilon'\rangle\langle\epsilon'| \quad (3)$$

- Yielding a time evolution (Dirac's pic) in the form

$$H_{\text{int}}(t) = \sum_{\alpha, \omega} e^{-i\omega t} S_{\alpha}(\omega) \otimes B_{\alpha}(t), \quad B_{\alpha}(t) = e^{iH_B t} B_{\alpha} e^{-iH_B t} \quad (4)$$



$$\frac{d}{dt} \rho(t) = [H + H_{\text{LS}}, \rho(t)] + \sum_{k, l, \omega} \gamma_{k, l}(\omega) \left[S_l(\omega) \rho(t) S_k^{\dagger}(\omega) - \frac{1}{2} \{ S_k^{\dagger}(\omega) S_l(\omega), \rho(t) \} \right]$$

which we can diagonalize over (k, l) , by defining $\gamma(\omega) = U \Sigma(\omega) U^{\dagger}$, with $\Sigma(\omega) = \text{Diag}(\varsigma_k(\omega))$ and the transformation of the jump operators - $L_k(\omega) = \sum_l U_{lk} S_l(\omega)$. Yielding

- With the operators $S_k(\omega)$ defined via

$$S_k(\omega) = \sum_{\epsilon' - \epsilon = \omega} \Pi_{\epsilon} S_k \Pi_{\epsilon'} \equiv \sum_{\epsilon' - \epsilon = \omega} |\epsilon\rangle\langle\epsilon| S_k |\epsilon'\rangle\langle\epsilon'| \quad (3)$$

- Yielding a time evolution (Dirac's pic) in the form

$$H_{\text{int}}(t) = \sum_{\alpha, \omega} e^{-i\omega t} S_{\alpha}(\omega) \otimes B_{\alpha}(t), \quad B_{\alpha}(t) = e^{iH_B t} B_{\alpha} e^{-iH_B t} \quad (4)$$



$$\frac{d}{dt}\rho(t) = [H + H_{\text{LS}}, \rho(t)] + \sum_{k, l, \omega} \gamma_{k, l}(\omega) \left[S_l(\omega) \rho(t) S_k^{\dagger}(\omega) - \frac{1}{2} \{ S_k^{\dagger}(\omega) S_l(\omega), \rho(t) \} \right]$$

which we can diagonalize over (k, l) , by defining $\gamma(\omega) = U \Sigma(\omega) U^{\dagger}$, with $\Sigma(\omega) = \text{Diag}(\varsigma_k(\omega))$ and the transformation of the jump operators - $L_k(\omega) = \sum_l U_{lk} S_l(\omega)$. Yielding

Lindblad

$$\frac{d}{dt}\rho(t) = [H + H_{\text{LS}}, \rho(t)] + \sum_{\omega} \sum_k \varsigma_k(\omega) \left[L_k(\omega) \rho(t) L_k^{\dagger}(\omega) - \frac{1}{2} \{ L_k^{\dagger}(\omega) L_k(\omega), \rho(t) \} \right]$$

Dissipation & Decoherence

Decoherence & Dissipation

Processes, through which a quantum system will lose its quantum properties over time, via a coupling to the environment.

Dissipation & Decoherence

Decoherence & Dissipation

Processes, through which a quantum system will lose its quantum properties over time, via a coupling to the environment.

- ▶ Loss of energy - **dissipation**

Dissipation & Decoherence

Decoherence & Dissipation

Processes, through which a quantum system will lose its quantum properties over time, via a coupling to the environment.

- ▶ Loss of energy - **dissipation**
- ▶ Loss of coherence - **decoherence**
 - ▶ Loss of information
 - ▶ Loss of entanglement
 - ▶ Loss of correlations, superpositions

Dissipation & Decoherence

Decoherence & Dissipation

Processes, through which a quantum system will lose its quantum properties over time, via a coupling to the environment.

- ▶ Loss of energy - **dissipation**
- ▶ Loss of coherence - **decoherence**
 - ▶ Loss of information
 - ▶ Loss of entanglement
 - ▶ Loss of correlations, superpositions
- ▶ Examples of such phenomena
 - ▶ Energy shift and relaxation to the ground state of the system.

Dissipation & Decoherence

Decoherence & Dissipation

Processes, through which a quantum system will lose its quantum properties over time, via a coupling to the environment.

- ▶ Loss of energy - **dissipation**
- ▶ Loss of coherence - **decoherence**
 - ▶ Loss of information
 - ▶ Loss of entanglement
 - ▶ Loss of correlations, superpositions
- ▶ Examples of such phenomena
 - ▶ Energy shift and relaxation to the ground state of the system.
 - ▶ Decay of the off-diagonal terms of the density operator, leading to loss of superpositions. In the spin-boson model, a possible solution for $\rho(t)$

$$\rho_{01}(t) \propto \exp\left(-Ct\tau \int_0^{\omega_c} d\omega \operatorname{sinc}^2(\omega\tau/2)\right)$$

- ▶ Or, in contrary, pure states can lose their purity due to the effect of decoherence.

Dissipation & Decoherence

Decoherence & Dissipation

Processes, through which a quantum system will lose its quantum properties over time, via a coupling to the environment.

- ▶ Loss of energy - **dissipation**
- ▶ Loss of coherence - **decoherence**
 - ▶ Loss of information
 - ▶ Loss of entanglement
 - ▶ Loss of correlations, superpositions
- ▶ Examples of such phenomena
 - ▶ Energy shift and relaxation to the ground state of the system.
 - ▶ Decay of the off-diagonal terms of the density operator, leading to loss of superpositions. In the spin-boson model, a possible solution for $\rho(t)$

$$\rho_{01}(t) \propto \exp\left(-Ct\tau \int_0^{\omega_c} d\omega \operatorname{sinc}^2(\omega\tau/2)\right)$$

- ▶ Or, in contrary, pure states can lose their purity due to the effect of decoherence.
- ▶ Note: relaxation mechanisms are always present in practice.

Steady state and stationarity

Why steady state?

Eigenstate thermalization hypothesis + ergodicity → Steady state system

Steady state and stationarity

Why steady state?

Eigenstate thermalization hypothesis + ergodicity \rightarrow Steady state system

- Question: *are there some systems so that this steady state is not achieved? If yes, what are those states and conditions? Do they belong to some isolated subspace of the Hilbert space $\tilde{\mathcal{H}} \in \mathcal{H}$?*

Steady state and stationarity

Why steady state?

Eigenstate thermalization hypothesis + ergodicity \rightarrow Steady state system

- ▶ Question: *are there some systems so that this steady state is not achieved? If yes, what are those states and conditions? Do they belong to some isolated subspace of the Hilbert space $\tilde{\mathcal{H}} \in \mathcal{H}$?*
- ▶ For the most widely used cases, one supposes **ergodicity**, meaning the state explores the accessible state space, as illustrated in figure (a), before reaching the supposed stationarity.




Figure 1: Illustration of ergodicity. ¹

¹Buča, B., Tindall, J. & Jaksch, D. Non-stationary coherent quantum many-body dynamics through dissipation

Steady state and stationarity

- In fact, it is **possible** to consider the microscopic properties \mathcal{L} , such that

$$\exists \quad \left\{ \{ \varphi_m, |\varphi_m\rangle \} \mid \mathcal{L} |\varphi_m\rangle = \varphi_m |\varphi_m\rangle \text{ and } \{ \varphi_m \} \in \sigma(\mathcal{L}) \right\}$$

meaning, that there can exist a decoherence-free subspace.

Steady state and stationarity

- In fact, it is **possible** to consider the microscopic properties \mathcal{L} , such that

$$\exists \left\{ \{ \varphi_m, |\varphi_m\rangle \} \mid \mathcal{L} |\varphi_m\rangle = \varphi_m |\varphi_m\rangle \text{ and } \{ \varphi_m \} \in \sigma(\mathcal{L}) \right\}$$

meaning, that there can exist a decoherence-free subspace.

Thus, in some cases, even when coupled to the environment, the system can undergo non-stationarity, which will not be the case using the *classical* ETH or ergodic assumptions.

Steady state and stationarity

- ▶ In fact, it is **possible** to consider the microscopic properties \mathcal{L} , such that

$$\exists \left\{ \{ \varphi_m, |\varphi_m\rangle \} \mid \mathcal{L} |\varphi_m\rangle = \varphi_m |\varphi_m\rangle \text{ and } \{ \varphi_m \} \in \sigma(\mathcal{L}) \right\}$$

meaning, that there can exist a decoherence-free subspace.

Thus, in some cases, even when coupled to the environment, the system can undergo non-stationarity, which will not be the case using the *classical* ETH or ergodic assumptions.

- ▶ Symmetry-preserving dissipation eliminates *non-coherent* eigenkets.
- ▶ Dark Hamiltonian \mathfrak{H} - driver of the *non-decaying* subspaces.

Steady state and stationarity

- In fact, it is **possible** to consider the microscopic properties \mathcal{L} , such that

$$\exists \quad \left\{ \{ \varphi_m, |\varphi_m\rangle \} \mid \mathcal{L} |\varphi_m\rangle = \varphi_m |\varphi_m\rangle \text{ and } \{ \varphi_m \} \in \sigma(\mathcal{L}) \right\}$$

meaning, that there can exist a decoherence-free subspace.

Thus, in some cases, even when coupled to the environment, the system can undergo non-stationarity, which will not be the case using the *classical* ETH or ergodic assumptions.

- Symmetry-preserving dissipation eliminates *non-coherent* eigenkets.
- Dark Hamiltonian \mathfrak{H} - driver of the *non-decaying* subspaces.




Figure 2: Decoherence-free subspaces, driven by \mathfrak{H} .²

² Buča, B., Tindall, J. & Jaksch, D. Non-stationary coherent quantum many-body dynamics through dissipation

Steady state and stationarity

How to identify those DF - **decoherence-free subspaces**?

Steady state and stationarity

How to identify those DF - **decoherence-free subspaces**?

$$\begin{aligned}\frac{d}{dt}\rho(t) &= \mathcal{L}\rho(t) = -[H, \rho(t)] + \sum_{\mu} (2c_{\mu}\rho c_{\mu}^{\dagger} - \{c_{\mu}c_{\mu}^{\dagger}, \rho\}) \\ \dot{\rho} &= \mathcal{L}\rho = \mathcal{E}(\rho) - Q\rho - \rho Q, \quad \mathcal{E} \equiv 2 \sum_l g_l c_l \rho c_l^{\dagger}, \quad Q \equiv P - iH, \quad P \equiv \sum_l g_l c_l^{\dagger} c_l\end{aligned}$$

Steady state and stationarity

$$\begin{aligned}\frac{d}{dt}\rho(t) &= \mathcal{L}\rho(t) = -[H, \rho(t)] + \sum_{\mu} (2c_{\mu}\rho c_{\mu}^{\dagger} - \{c_{\mu}c_{\mu}^{\dagger}, \rho\}) \\ \dot{\rho} &= \mathcal{L}\rho = \mathcal{E}(\rho) - Q\rho - \rho Q, \quad \mathcal{E} \equiv 2\sum_l g_l c_l \rho c_l^{\dagger}, \quad Q \equiv P - iH, \quad P \equiv \sum_l g_l c_l^{\dagger} c_l\end{aligned}$$

- For a steady-state ρ_{∞} , one clearly has $\mathcal{L}\rho_{\infty} = 0$

Steady state and stationarity

$$\frac{d}{dt}\rho(t) = \mathcal{L}\rho(t) = -[H, \rho(t)] + \sum_{\mu} (2c_{\mu}\rho c_{\mu}^{\dagger} - \{c_{\mu}c_{\mu}^{\dagger}, \rho\})$$

$$\dot{\rho} = \mathcal{L}\rho = \mathcal{E}(\rho) - Q\rho - \rho Q, \quad \mathcal{E} \equiv 2 \sum_l g_l c_l \rho c_l^{\dagger}, \quad Q \equiv P - iH, \quad P \equiv \sum_l g_l c_l^{\dagger} c_l$$

- For a steady-state ρ_{∞} , one clearly has $\mathcal{L}\rho_{\infty} = 0$
- Density ρ_n non stationary, if the corresponding eigenvalues ϱ_n are purely imaginary.

$$\begin{aligned} \mathcal{L}\rho_n &= \varrho_n \rho_n \\ &= -i\mathfrak{H}\rho_n = -i\lambda_n \rho_n, \quad \lambda \in \mathbb{R} \end{aligned} \tag{5}$$




Figure 3: Meaning of eigenvalues of \mathcal{L} .

Non-dissipative subspaces

Let $|\Psi\rangle$ - a pure state, that is **not** affected by dissipation. Then,

$$\blacktriangleright |\Psi\rangle \in \text{Ker}(P) \equiv \{|\Psi\rangle \mid \hat{c}_l |\Psi\rangle = 0\} = \text{Ker}(P) \cap \text{EigSp}(H) \equiv \mathcal{H}_{ND}$$

Non-dissipative subspaces

Let $|\Psi\rangle$ - a pure state, that is **not** affected by dissipation. Then,

- ▶ $|\Psi\rangle \in \text{Ker}(P) \equiv \{|\Psi\rangle \mid \hat{c}_l |\Psi\rangle = 0\} = \text{Ker}(P) \cap \text{EigSp}(H) \equiv \mathcal{H}_{ND}$
- ▶ Meaning that if for a density operator ρ , such that

$$\text{Range}(\rho) = \text{span}(\{|\varphi_i\rangle\}) \text{ ,with } \{\varphi_i\} \in \mathcal{H}_{ND} \quad (6)$$

will represent a stationary state. We call $\{\varphi_i\}$ - dark states.

Non-dissipative subspaces

Let $|\Psi\rangle$ - a pure state, that is **not** affected by dissipation. Then,

- ▶ $|\Psi\rangle \in \text{Ker}(P) \equiv \{|\Psi\rangle \mid \hat{c}_l |\Psi\rangle = 0\} = \text{Ker}(P) \cap \text{EigSp}(H) \equiv \mathcal{H}_{ND}$
- ▶ Meaning that if for a density operator ρ , such that

$$\text{Range}(\rho) = \text{span}(\{|\varphi_i\rangle\}) \text{ ,with } \{\varphi_i\} \in \mathcal{H}_{ND} \quad (6)$$

will represent a stationary state. We call $\{\varphi_i\}$ - dark states.

- ▶ We want to define the dark Hamiltonian \mathfrak{H} , that

Non-dissipative subspaces

Let $|\Psi\rangle$ - a pure state, that is **not** affected by dissipation. Then,

- ▶ $|\Psi\rangle \in \text{Ker}(P) \equiv \{|\Psi\rangle \mid \hat{c}_l |\Psi\rangle = 0\} = \text{Ker}(P) \cap \text{EigSp}(H) \equiv \mathcal{H}_{ND}$
- ▶ Meaning that if for a density operator ρ , such that

$$\text{Range}(\rho) = \text{span}(\{|\varphi_i\rangle\}) \text{ ,with } \{\varphi_i\} \in \mathcal{H}_{ND} \quad (6)$$

will represent a stationary state. We call $\{\varphi_i\}$ - dark states.

- ▶ We want to define the dark Hamiltonian \mathfrak{H} , that
 - ▶ $\text{EigSp}(\mathfrak{H}) = \{|\varphi_i\rangle\}$ such that $|\varphi_m\rangle \perp |\varphi_n\rangle$
 - ▶ $\mathfrak{H} |\varphi_m\rangle\langle\varphi_n| = (\omega_m - \omega_n) |\varphi_m\rangle\langle\varphi_n|$
 - ▶ For $\rho_{nm} = |\varphi_m\rangle\langle\varphi_n|$, $\rho_{nm}^\dagger \rho_{n'm'} = \delta_{nm} \delta_{n'm'}$
 - ▶ Hermicity - $\mathfrak{H} = \mathfrak{H}^\dagger$

Non-dissipative subspaces

There are multiple theorems giving precisions on the *Dark states*.

Non-dissipative subspaces

There are multiple theorems giving precisions on the *Dark states*.

Theorem: conditions stationarity

The set $\{|\varphi_m\rangle\} \in \text{EigSp}(\mathfrak{H})$ with $|\varphi_m\rangle \perp |\varphi_n\rangle$ IFF

- ▶ $Q^\dagger |\varphi_n\rangle = \lambda |\varphi_n\rangle$, $\forall n$
- ▶ $c_k |\varphi_n\rangle = \lambda_{kn} |\varphi_n\rangle$, with $\sum_l \gamma_l |\lambda_{ln}|^2 = \text{Re}(\lambda_n)$
- ▶ Relations between λ_{kn} and γ_k :

$$\text{Re} \left[\sum_k \gamma_k \left(2\lambda_{kn} \lambda_{km}^* - |\lambda_{kn}|^2 - |\lambda_{km}|^2 \right) \right] = 0 \quad \forall n, m$$

Non-dissipative subspaces

There are multiple theorems giving precisions on the *Dark states*.

Theorem: conditions stationarity

The set $\{|\varphi_m\rangle\} \in \text{EigSp}(\mathfrak{H})$ with $|\varphi_m\rangle \perp |\varphi_n\rangle$ IFF

- ▶ $Q^\dagger |\varphi_n\rangle = \lambda |\varphi_n\rangle$, $\forall n$
- ▶ $c_k |\varphi_n\rangle = \lambda_{kn} |\varphi_n\rangle$, with $\sum_l \gamma_l |\lambda_{ln}|^2 = \text{Re}(\lambda_n)$
- ▶ Relations between λ_{kn} and γ_k :

$$\text{Re} \left[\sum_k \gamma_k \left(2\lambda_{kn} \lambda_{km}^* - |\lambda_{kn}|^2 - |\lambda_{km}|^2 \right) \right] = 0 \quad \forall n, m$$

- ▶ Corollary for eigenvalues ω_i for the dark Hamiltonian \mathfrak{H} :

$$\omega_n - \omega_m = \text{Im} \left[\sum_k 2\gamma_k \lambda_{kn} \lambda_{km}^* - \langle \varphi_m | H | \varphi_n \rangle \right] \quad (7)$$

Non-dissipative subspaces

There are multiple theorems giving precisions on the *Dark states*.

Theorem: conditions stationarity

If there exists no subspace $\mathcal{S} \subseteq \mathcal{H}$ with $\mathcal{S} \perp \mathcal{H}_{ND}$ such that $\hat{c}_l(\mathcal{S}) \subseteq \mathcal{S}$, $\forall \hat{c}_l$, then the only oscillating coherences are the dark states, in the form $|\varphi_n\rangle\langle\varphi_m|$, such that $\{|\varphi_n\rangle\} \in \mathcal{H}_{DS}$ defined above.

Non-dissipative subspaces

There are multiple theorems giving precisions on the *Dark states*.

Theorem: conditions stationarity

If there exists no subspace $\mathcal{S} \subseteq \mathcal{H}$ with $\mathcal{S} \perp \mathcal{H}_{ND}$ such that $\hat{c}_l(\mathcal{S}) \subseteq \mathcal{S}$, $\forall \hat{c}_l$, then the only oscillating coherences are the dark states, in the form $|\varphi_n\rangle\langle\varphi_m|$, such that $\{|\varphi_n\rangle\} \in \mathcal{H}_{DS}$ defined above.

- The notations used up to now do **not** correspond to the one used previously, when defining the \mathcal{L} equation via the spectrum $\omega \in \sigma(H_S)$

Non-dissipative subspaces

There are multiple theorems giving precisions on the *Dark states*.

Theorem: conditions stationarity

If there exists no subspace $\mathcal{S} \subseteq \mathcal{H}$ with $\mathcal{S} \perp \mathcal{H}_{ND}$ such that $\hat{c}_l(\mathcal{S}) \subseteq \mathcal{S}$, $\forall \hat{c}_l$, then the only oscillating coherences are the dark states, in the form $|\varphi_n\rangle\langle\varphi_m|$, such that $\{|\varphi_n\rangle\} \in \mathcal{H}_{DS}$ defined above.

- ▶ The notations used up to now do **not** correspond to the one used previously, when defining the \mathcal{L} equation via the spectrum $\omega \in \sigma(H_S)$
- ▶ Example of change of notation: $\sum_{\mu} \mapsto \sum_k \sum_{\omega}$.

A more general case - non-stationarity

- ▶ The dark Hamiltonian \mathfrak{H} does not need to be Hermitian in general

A more general case - non-stationarity

- ▶ The dark Hamiltonian \mathfrak{H} does not need to be Hermitian in general
- ▶ How to find ρ , such that, $\mathcal{L}\rho \neq 0$?

A more general case - non-stationarity

- ▶ The dark Hamiltonian \mathfrak{H} does not need to be Hermitian in general
- ▶ How to find ρ , such that, $\mathcal{L}\rho \neq 0$?

Condition:

This is realized, if $\exists A : [H, A] = -\lambda A$ and $[c_\mu, A] = [c_\mu^\dagger, A] = 0$ with $\lambda \in \mathbb{R}$

A more general case - non-stationarity


- ▶ The dark Hamiltonian \mathfrak{H} does not need to be Hermitian in general
- ▶ How to find ρ , such that, $\mathcal{L}\rho \neq 0$?

Condition:

This is realized, if $\exists A : [H, A] = -\lambda A$ and $[c_\mu, A] = [c_\mu^\dagger, A] = 0$ with $\lambda \in \mathbb{R}$

Corollary:

As a consequence, $\exists \rho_{mn} \equiv A^n \rho_\infty (A^m)^\dagger$ $m, n \in \mathbb{Z}$ and $\mathcal{L}\rho_{mn} = i(m - n)\lambda\rho_{mn}$



Recap & terminology

- ▶ Decoherence, dissipation - loss of quantum properties

Recap & terminology

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state \Leftrightarrow ergodicity & ETH

Recap & terminology

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state \Leftrightarrow ergodicity & ETH
- ▶ **Dark states** $c_k |\varphi_n\rangle = 0 \ \forall n$ - invisible for decoherence.

Recap & terminology

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state \rightleftharpoons ergodicity & ETH
- ▶ **Dark states** $c_k |\varphi_n\rangle = 0 \ \forall n$ - invisible for decoherence.
- ▶ $\rho_{mn} = |\varphi_m\rangle\langle\varphi_n| \in \text{EigSp}(\mathcal{L})$ with eigenval. $\lambda_{mn} = -i(E_n - E_m)$

Recap & terminology

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state \rightleftharpoons ergodicity & ETH
- ▶ **Dark states** $c_k |\varphi_n\rangle = 0 \ \forall n$ - invisible for decoherence.
- ▶ $\rho_{mn} = |\varphi_m\rangle\langle\varphi_n| \in \text{EigSp}(\mathcal{L})$ with eigenval. $\lambda_{mn} = -i(E_n - E_m)$
- ▶ Mixed coherences $\rho_{nm} = (A^\dagger)^n \rho_\infty A^m$ also in $\text{EigSp}(\mathcal{L})$ with eigenvalues $\lambda_{nm}^j = i(n - m)\omega_j$, with $\omega_j A_j = [H, A_j]$

Recap & terminology

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state \rightleftharpoons ergodicity & ETH
- ▶ **Dark states** $c_k |\varphi_n\rangle = 0 \ \forall n$ - invisible for decoherence.
- ▶ $\rho_{mn} = |\varphi_m\rangle\langle\varphi_n| \in \text{EigSp}(\mathcal{L})$ with eigenval. $\lambda_{mn} = -i(E_n - E_m)$
- ▶ Mixed coherences $\rho_{nm} = (A^\dagger)^n \rho_\infty A^m$ also in $\text{EigSp}(\mathcal{L})$ with eigenvalues $\lambda_{nm}^j = i(n - m)\omega_j$, with $\omega_j A_j = [H, A_j]$

Remark

It is important to distinguish the mixed coherences and decoherence free subspace of dark space.

Namely, $\hat{c}_l |\varphi_n\rangle = 0$ for dark states and $\hat{c}_l \rho_{nm} \hat{c}_l^\dagger \neq 0$

Recap & terminology

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state \rightleftharpoons ergodicity & ETH
- ▶ **Dark states** $c_k |\varphi_n\rangle = 0 \ \forall n$ - invisible for decoherence.
- ▶ $\rho_{mn} = |\varphi_m\rangle\langle\varphi_n| \in \text{EigSp}(\mathcal{L})$ with eigenval. $\lambda_{mn} = -i(E_n - E_m)$
- ▶ Mixed coherences $\rho_{nm} = (A^\dagger)^n \rho_\infty A^m$ also in $\text{EigSp}(\mathcal{L})$ with eigenvalues $\lambda_{nm}^j = i(n - m)\omega_j$, with $\omega_j A_j = [H, A_j]$

Remark

It is important to distinguish the mixed coherences and decoherence free subspace of dark space.

Namely, $\hat{c}_l |\varphi_n\rangle = 0$ for dark states and $\hat{c}_l \rho_{nm} \hat{c}_l^\dagger \neq 0$

Corollary

The mixed coherence states' evolution depends on dissipative processes.