

# Open Quantum System

## Lectures:

Last time:

- ▶ Daniel Manzano, A short introduction to the Lindblad master equation (*all*)
- ▶ Breuer and Petruccione, The Theory of Open Quantum Systems (*ch. 3 - 4.3*)
- ▶ Daniel A. Lidar, Notes on the Theory of Open Quantum Systems (*up to ch. 12*)

Today:

- ▶ Breuer and Petruccione, The Theory of Open Quantum Systems (*ch. 3 - 4.3*)
- ▶ Daniel A. Lidar, Notes on the Theory of Open Quantum Systems (*up to ch. 12*)
- ▶ B.Kraus, H.P.Buchler, S. Diehl, A. Kantian, A. Micheli, & P. Zoller, Preparation of Entangled States by Quantum Markov Processes
- ▶ Buča, B., Tindall, J. & Jaksch, D. Non-stationary coherent quantum many-body dynamics through dissipation
- ▶ Victor V. Albert & Liang Jiang, Symmetries and conserved quantities in Lindblad master equations
- ▶ Cameron Booker, Berislav Buča, Dieter Jaksch, Non-stationarity and Dissipative Time Crystals: Spectral Properties and Finite-Size Effects

November 12, 2023

# Recalling

- Derivation of the Lindblad equation via different approximations:
  - Von Neumann evolution equation,  $H_{\text{int}} = \sum_k S_k \otimes B_k \rightarrow$  Weak coupling, Born, Markov & Rotating wave  $\rightarrow$  Redfield equation  
*Note: no stationarity of the system was assumed!*
  - Yielding (Schrodinger pic.):

$$\begin{aligned} \frac{d}{dt}\rho(t) = [H + H_{\text{LS}}, \rho(t)] + \\ + \sum_{k,l,\omega} \gamma_{k,l}(\omega) \left[ S_l(\omega)\rho(t)S_k^\dagger(\omega) - \frac{1}{2}\{S_k^\dagger(\omega)S_l(\omega), \rho(t)\} \right] \end{aligned} \quad (1)$$

- With  $\gamma_{k,l}$ ,  $\pi_{k,l}$  - defined via  $\Gamma_{k,l}(\omega)$ 's real and imaginary parts.

$$\begin{aligned} H_{\text{LS}} &= \sum_{\omega,k,l} \pi_{k,l}(\omega) S_k^\dagger(\omega) S_l(\omega) \\ \Gamma_{k,l}(\omega) &= \int_0^\infty e^{i\omega s} \text{Tr}_B [B_k^\dagger(t) B_l(t-s) \rho_B(0)] \end{aligned} \quad (2)$$

- With the operators  $S_k(\omega)$  defined via

$$S_k(\omega) = \sum_{\epsilon' - \epsilon = \omega} \Pi_{\epsilon} S_k \Pi_{\epsilon'} \equiv \sum_{\epsilon' - \epsilon = \omega} |\epsilon\rangle\langle\epsilon| S_k |\epsilon'\rangle\langle\epsilon'| \quad (3)$$

- Yielding a time evolution (Dirac's pic) in the form

$$H_{\text{int}}(t) = \sum_{\alpha, \omega} e^{-i\omega t} S_{\alpha}(\omega) \otimes B_{\alpha}(t), \quad B_{\alpha}(t) = e^{iH_B t} B_{\alpha} e^{-iH_B t} \quad (4)$$



$$\frac{d}{dt}\rho(t) = [H + H_{\text{LS}}, \rho(t)] + \sum_{k,l,\omega} \gamma_{k,l}(\omega) \left[ S_l(\omega) \rho(t) S_k^{\dagger}(\omega) - \frac{1}{2} \{ S_k^{\dagger}(\omega) S_l(\omega), \rho(t) \} \right]$$

which we can diagonalize over  $(k, l)$ , by defining  $\gamma(\omega) = U \Sigma(\omega) U^{\dagger}$ , with  $\Sigma(\omega) = \text{Diag}(\varsigma_k(\omega))$  and the transformation of the jump operators -  $L_k(\omega) = \sum_l U_{lk} S_l(\omega)$ . Yielding

## Lindblad

$$\frac{d}{dt}\rho(t) = [H + H_{\text{LS}}, \rho(t)] + \sum_{\omega} \sum_k \varsigma_k(\omega) \left[ L_k(\omega) \rho(t) L_k^{\dagger}(\omega) - \frac{1}{2} \{ L_k^{\dagger}(\omega) L_k(\omega), \rho(t) \} \right]$$

# Last time

- ▶ Decoherence, dissipation - loss of quantum properties

# Last time

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state  $\Leftrightarrow$  ergodicity & ETH

# Last time

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state  $\Leftrightarrow$  ergodicity & ETH
- ▶ Dark states - **invisible for decoherence**

# Last time

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state  $\Leftrightarrow$  ergodicity & ETH
- ▶ Dark states - **invisible for decoherence**
- ▶ Oscillating coherences (OC)  $\neq$  dark states (DS)

$$\varphi \in \text{DS} : L_k |\varphi\rangle = 0 \quad \& \quad L_k \rho L_k^\dagger \neq 0$$

# Last time

- ▶ Decoherence, dissipation - loss of quantum properties
- ▶ Steady-state  $\Leftrightarrow$  ergodicity & ETH
- ▶ Dark states - **invisible for decoherence**
- ▶ Oscillating coherences (OC)  $\neq$  dark states (DS)


$$\varphi \in \text{DS} : L_k |\varphi\rangle = 0 \quad \& \quad L_k \rho L_k^\dagger \neq 0$$

- ▶ Dark Hamiltonian - driver of the non stationary effects




# (non)-Stationarity

- ▶ Question - how to determine the steady states and non-steady states?




(a) Ergodicity




(b) Oscillating coherences  
via  $J$

# (non)-Stationarity

- ▶ Question - how to determine the steady states and non-steady states?
- ▶ Before that, one defines:
  - ▶ Symmetry-preserving dissipation
  - ▶ Dark Hamiltonian  $\mathfrak{H}$  - not necessarily hermitian




(a) Ergodicity




(b) Oscillating coherences  
via  $\mathfrak{H}$

# (non)-Stationarity


- ▶ Question - how to determine the steady states and non-steady states?
- ▶ Before that, one defines:
  - ▶ Symmetry-preserving dissipation
  - ▶ Dark Hamiltonian  $\mathfrak{H}$  - not necessarily hermitian
- ▶ Evolution of the states  $|\rho\rangle\rangle$  under Liouvillian -  $\mathcal{L}|\rho\rangle\rangle = \rho|\rho\rangle\rangle$  with  $\text{Re}(\lambda) \leq 0$



(a) Ergodicity




(b) Oscillating coherences via  $\mathfrak{H}$




(c) Eigenvalues of  $\mathcal{L}$

# (non)-Stationarity


- ▶ Question - how to determine the steady states and non-steady states?
- ▶ Before that, one defines:
  - ▶ Symmetry-preserving dissipation
  - ▶ Dark Hamiltonian  $\mathfrak{H}$  - not necessarily hermitian
- ▶ Evolution of the states  $|\rho\rangle\rangle$  under Liouvillian -  $\mathcal{L}|\rho\rangle\rangle = \rho|\rho\rangle\rangle$  with  $\text{Re}(\lambda) \leq 0$ 
  - ▶ Decays
  - ▶ Oscillating coherences
  - ▶ Spirals



(a) Ergodicity



(b) Oscillating coherences via  $\mathfrak{H}$



(c) Eigenvalues of  $\mathcal{L}$

# (non)-Stationarity

## Theorem

Let  $|\rho_\infty\rangle\rangle$  - steady state, s.t.  $\mathcal{L}|\rho_\infty\rangle\rangle = 0$ . If  $\exists A$  s.t.

$$[H, A]\rho_\infty = \lambda A\rho_\infty \quad [L_k, A]\rho_\infty = [L_k^\dagger, A]\rho_\infty = 0$$

Then,

$$\mathcal{L}(A|\rho_\infty\rangle\rangle) = i\lambda A|\rho_\infty\rangle\rangle \quad \text{i.e. } A|\rho_\infty\rangle\rangle \in \text{EigSp}(\mathcal{L})$$

with  $\lambda$  - purely imaginary (See oscillating coherences).

# (non)-Stationarity

## Theorem

Let  $|\rho_\infty\rangle\rangle$  - steady state, s.t.  $\mathcal{L}|\rho_\infty\rangle\rangle = 0$ . If  $\exists A$  s.t.

$$[H, A]\rho_\infty = \lambda A\rho_\infty \quad [L_k, A]\rho_\infty = [L_k^\dagger, A]\rho_\infty = 0$$

Then,

$$\mathcal{L}(A|\rho_\infty\rangle\rangle) = i\lambda A|\rho_\infty\rangle\rangle \quad \text{i.e. } A|\rho_\infty\rangle\rangle \in \text{EigSp}(\mathcal{L})$$

with  $\lambda$  - purely imaginary (See oscillating coherences).

Intuitively, given  $|\rho_\infty\rangle\rangle$ , the "symmetry"  $A$  will "induce" the non-stationary states.

# (non)-Stationarity

One defines

- ▶ Super-operator  $\hat{\hat{A}}$ :  $\hat{\hat{A}}|\rho\rangle\rangle \equiv \hat{A}|\rho\rangle\rangle$ , with operator  $\hat{A}$  with defined properties.

# (non)-Stationarity

One defines

- ▶ Super-operator  $\hat{\hat{A}}$ :  $\hat{\hat{A}}|\rho\rangle\rangle \equiv \hat{A}|\rho\rangle\rangle$ , with operator  $\hat{A}$  with defined properties.
- ▶ The state  $|\rho_\infty\rangle\rangle$  as before,  $\mathcal{L}|\rho_\infty\rangle\rangle = 0$



# (non)-Stationarity

One defines

- ▶ Super-operator  $\hat{\hat{A}}$ :  $\hat{\hat{A}}|\rho\rangle\rangle \equiv \hat{A}|\rho\rangle\rangle$ , with operator  $\hat{A}$  with defined properties.
- ▶ The state  $|\rho_\infty\rangle\rangle$  as before,  $\mathcal{L}|\rho_\infty\rangle\rangle = 0$
- ▶ The eigen-operator relation for super-operators

$$[\mathcal{L}, \hat{\hat{A}}] = i\lambda\hat{\hat{A}}$$

## Corollary

If there exists an operator  $\hat{A}$ , such that

$$[H, \hat{A}] = \lambda \hat{A} \quad \text{and} \quad [L_k, \hat{A}] = [L_k^\dagger, \hat{A}] = 0 \quad \forall k$$

## Corollary

If there exists an operator  $\hat{A}$ , such that

$$[H, \hat{A}] = \lambda \hat{A} \quad \text{and} \quad [L_k, \hat{A}] = [L_k^\dagger, \hat{A}] = 0 \quad \forall k$$

Then, one can define

$$|\rho_{nm}\rangle\rangle = (\hat{A})^n |\rho_\infty\rangle\rangle (\hat{A}^\dagger)^m$$

## Corollary

If there exists an operator  $\hat{A}$ , such that

$$[H, \hat{A}] = \lambda \hat{A} \quad \text{and} \quad [L_k, \hat{A}] = [L_k^\dagger, \hat{A}] = 0 \quad \forall k$$

Then, one can define

$$|\rho_{nm}\rangle\rangle = (\hat{A})^n |\rho_\infty\rangle\rangle (\hat{A}^\dagger)^m$$

Then, one has

$$\boxed{\mathcal{L}|\rho_{nm}\rangle\rangle = i\lambda(n - m)|\rho_{nm}\rangle\rangle}$$

## Corollary

If there exists an operator  $\hat{A}$ , such that

$$[H, \hat{A}] = \lambda \hat{A} \quad \text{and} \quad [L_k, \hat{A}] = [L_k^\dagger, \hat{A}] = 0 \quad \forall k$$

Then, one can define

$$|\rho_{nm}\rangle\rangle = (\hat{A})^n |\rho_\infty\rangle\rangle (\hat{A}^\dagger)^m$$

Then, one has

$$\mathcal{L}|\rho_{nm}\rangle\rangle = i\lambda(n - m)|\rho_{nm}\rangle\rangle$$

- Properties of eigenstates  $|\rho\rangle\rangle \leftrightarrow \sigma(\mathcal{L})$  derived independently.




Figure 2

## Corollary

If there exists an operator  $\hat{A}$ , such that

$$[H, \hat{A}] = \lambda \hat{A} \quad \text{and} \quad [L_k, \hat{A}] = [L_k^\dagger, \hat{A}] = 0 \quad \forall k$$

Then, one can define

$$|\rho_{nm}\rangle\rangle = (\hat{A})^n |\rho_\infty\rangle\rangle (\hat{A}^\dagger)^m$$

Then, one has

$$\mathcal{L}|\rho_{nm}\rangle\rangle = i\lambda(n - m)|\rho_{nm}\rangle\rangle$$

- Properties of eigenstates  $|\rho\rangle\rangle \leftrightarrow \sigma(\mathcal{L})$  derived independently.
- Main ingredients -  $\{H, L_k, \hat{A}\}$  and  $\rho_\infty$




Figure 2

## Corollary

If there exists an operator  $\hat{A}$ , such that

$$[H, \hat{A}] = \lambda \hat{A} \quad \text{and} \quad [L_k, \hat{A}] = [L_k^\dagger, \hat{A}] = 0 \quad \forall k$$

Then, one can define

$$|\rho_{nm}\rangle\rangle = (\hat{A})^n |\rho_\infty\rangle\rangle (\hat{A}^\dagger)^m$$

Then, one has

$$\mathcal{L}|\rho_{nm}\rangle\rangle = i\lambda(n - m)|\rho_{nm}\rangle\rangle$$

- ▶ Properties of eigenstates  $|\rho\rangle\rangle \leftrightarrow \sigma(\mathcal{L})$  derived independently.
- ▶ Main ingredients -  $\{H, L_k, \hat{A}\}$  and  $\rho_\infty$
- ▶ Symmetry-preserving dissipations & dark Hamiltonian




Figure 2

# Implications?

Last times:

$$\mathcal{L}\rho = \dot{\rho} = -i[H + H_{\text{LS}}, \rho] + \sum_{\omega,k} \varsigma_k(\omega) \left( L_k(\omega) \rho L_k^\dagger(\omega) - \frac{1}{2} \{ L_k^\dagger(\omega) L_k(\omega), \rho \} \right)$$



# Implications?

Last times:

$$\mathcal{L}\rho = \dot{\rho} = -i[H + H_{LS}, \rho] + \sum_{\omega,k} \varsigma_k(\omega) \left( L_k(\omega) \rho L_k^\dagger(\omega) - \frac{1}{2} \{ L_k^\dagger(\omega) L_k(\omega), \rho \} \right)$$

With the diagonalized elements  $L_k(\omega) = \sum_l U_{lk} S_l(\omega)$  and

$$S_l(\omega) = \sum_{\omega=\epsilon'-\epsilon} \Pi_\epsilon S_l \Pi_{\epsilon'} \equiv \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| S_l |\epsilon'\rangle\langle\epsilon'|$$

# Implications?

Last times:

$$\mathcal{L}\rho = \dot{\rho} = -i[H + H_{LS}, \rho] + \sum_{\omega,k} \varsigma_k(\omega) \left( L_k(\omega) \rho L_k^\dagger(\omega) - \frac{1}{2} \{ L_k^\dagger(\omega) L_k(\omega), \rho \} \right)$$

With the diagonalized elements  $L_k(\omega) = \sum_l U_{lk} S_l(\omega)$  and

$$S_l(\omega) = \sum_{\omega=\epsilon'-\epsilon} \Pi_\epsilon S_l \Pi_{\epsilon'} \equiv \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| S_l |\epsilon'\rangle\langle\epsilon'|$$

In the literature in question -

$$\mathcal{L}\rho = -i[H, \rho] + \sum_\mu \left( 2\tilde{L}_\mu \rho \tilde{L}_\mu^\dagger - \{ \tilde{L}_\mu^\dagger \tilde{L}_\mu, \rho \} \right)$$

# Implications?

Last times:

$$\mathcal{L}\rho = \dot{\rho} = -i[H + H_{LS}, \rho] + \sum_{\omega,k} \varsigma_k(\omega) \left( L_k(\omega) \rho L_k^\dagger(\omega) - \frac{1}{2} \{ L_k^\dagger(\omega) L_k(\omega), \rho \} \right)$$

With the diagonalized elements  $L_k(\omega) = \sum_l U_{lk} S_l(\omega)$  and

$$S_l(\omega) = \sum_{\omega=\epsilon'-\epsilon} \Pi_\epsilon S_l \Pi_{\epsilon'} \equiv \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| S_l |\epsilon'\rangle\langle\epsilon'|$$

In the literature in question -

$$\mathcal{L}\rho = -i[H, \rho] + \sum_\mu \left( 2\tilde{L}_\mu \rho \tilde{L}_\mu^\dagger - \{ \tilde{L}_\mu^\dagger \tilde{L}_\mu, \rho \} \right)$$

One may try to suggest to go from the 2nd to the 1st:

$$H \mapsto H + H_{LS} \quad \mu \mapsto (\omega, k) \quad \Longrightarrow \quad \tilde{L}_{\mu=(\omega,k)} \mapsto \sqrt{\frac{\varsigma_k(\omega)}{2}} L_k(\omega)$$

# Implications?

Using the definitions:

$$L_\mu \mapsto \sqrt{\frac{\varsigma_k(\omega)}{2}} L_k(\omega) = \sqrt{\frac{\varsigma_k(\omega)}{2}} \sum_{\omega = \epsilon' - \epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|$$

# Implications?

Using the definitions:

$$L_\mu \mapsto \sqrt{\frac{\varsigma_k(\omega)}{2}} L_k(\omega) = \sqrt{\frac{\varsigma_k(\omega)}{2}} \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|$$

For an operator  $A$  such that  $[H, A] = \lambda A$  and  $[L_\mu, A] = [L_\mu^\dagger, A] = 0$ ,

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

which must be hold  $\forall \omega, k$

# Implications?

For an operator  $A$  such that  $[H, A] = \lambda A$  and  $[L_\mu, A] = [L_\mu^\dagger, A] = 0$ ,

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

which must hold  $\forall \omega, k$

# Implications?

For an operator  $A$  such that  $[H, A] = \lambda A$  and  $[L_\mu, A] = [L_\mu^\dagger, A] = 0$ ,

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

which must hold  $\forall \omega, k$

How about the condition on  $H$ ? That is,  $[H, A] = \lambda A$ . One suggested:

$$[H, A] \mapsto [H + H_{\text{LS}}] \quad \implies \quad [H + H_{\text{LS}}, A] = \lambda A$$

# Implications?

For an operator  $A$  such that  $[H, A] = \lambda A$  and  $[L_\mu, A] = [L_\mu^\dagger, A] = 0$ ,

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

which must be hold  $\forall \omega, k$

How about the condition on  $H$ ? That is,  $[H, A] = \lambda A$ . One suggested:

$$[H, A] \mapsto [H + H_{\text{LS}}] \implies [H + H_{\text{LS}}] = \lambda A$$

How to interpret  $A$ ?



# Implications?

For an operator  $A$  such that  $[H, A] = \lambda A$  and  $[L_\mu, A] = [L_\mu^\dagger, A] = 0$ ,

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

which must hold  $\forall \omega, k$

How about the condition on  $H$ ? That is,  $[H, A] = \lambda A$ . One suggested:

$$[H, A] \mapsto [H + H_{\text{LS}}] \implies [H + H_{\text{LS}}] = \lambda A$$

How to interpret  $A$ ?

- Symmetry generator? (angular momentum, energy)
- Ladders?

# Implications?

For an operator  $A$  such that  $[H, A] = \lambda A$  and  $[L_\mu, A] = [L_\mu^\dagger, A] = 0$ ,

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

which must hold  $\forall \omega, k$

How about the condition on  $H$ ? That is,  $[H, A] = \lambda A$ . One suggested:

$$[H, A] \mapsto [H + H_{\text{LS}}] \implies [H + H_{\text{LS}}] = \lambda A$$

How to interpret  $A$ ?

- ▶ Symmetry generator? (angular momentum, energy)
- ▶ Ladders?
- ▶ Example:  $[L_z, L_\pm] = \pm \hbar L_\pm$ ,  $[H, a^\dagger] = \hbar\omega a^\dagger$

# Implications?

Let us recall again:

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

# Implications?

Let us recall again:

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

► Can we expand more?

# Implications?

Let us recall again:

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

- ▶ Can we expand more?
- ▶ Is  $L_k$  Hermitian? How are  $A$  and  $L_k$  related?

# Implications?

Let us recall again:

$$\left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon\rangle\langle\epsilon| L_k |\epsilon'\rangle\langle\epsilon'|, A \right] = \left[ \sum_{\omega=\epsilon'-\epsilon} |\epsilon'\rangle\langle\epsilon'| L_k^\dagger |\epsilon\rangle\langle\epsilon|, A \right] = 0$$

- ▶ Can we expand more?
- ▶ Is  $L_k$  Hermitian? How are  $A$  and  $L_k$  related?
- ▶ Remember the definition of  $L_k$  - unitary transform of  $S_k$ , defined via  $H_{\text{int}} = \sum_k S_k \otimes B_k$ . Thus  $H_{\text{int}} = H_{\text{int}}^\dagger$ ,  $S_k$  - not necessarily hermitian.