Notas de aula- MAT01355

Leonardo Suarez

15 de abril de 2026

Notas de aula referentes à disciplina MAT01355 - Álgebra Linear I-A, ministrada pelo prof. Leandro Farina no semestre letivo 24/2. A bibliografia principal é a sexta edição do livro *Linear Algebra and Its Applications*, de David Lay, Steven R. Lay e Judi J. McDonald.

Conteúdo

I Linear Equations in Linear Algebra **2**
1 Row reduction and echelon forms 2
2 The matrix equation $Ax = b$ 2
3 Solution sets of linear systems 3
4 Linear independence 3
5 The matrix of a linear transformation 3

II Matrix Algebra **3**
1 Matrix operations 3
2 The inverse of a matrix 4
3 Characterizations of invertible matrices 5

III Determinants **5**
1 Introduction to determinants 5
2 Properties of determinants 6

IV Vector Spaces **6**
1 Vector spaces and subspaces 6
2 Null space, column space, row space, and linear transformations 7
3 Linearly independent sets; bases 7
4 Coordinate systems 8
5 The dimension of a vector space 8
6 Change of basis 9

V Eigenvalues and Eigenvectors **9**
1 Eigenvectors and eigenvalues 9
2 The characteristic equation 9
3 Diagonalization 10

VI Orthogonality and Least Squares **10**
1 Inner product, length, and orthogonality 10
2 Orthogonal sets 11
3 Orthogonal projections 12

4 The Gram-Schmidt process 12
5 Least-squares problems 13

VII Symmetric Matrices and Quadratic Forms 13
1 Quadratic forms 14
2 Constrained optimization 14

I Linear Equations in Linear Algebra

1 Row reduction and echelon forms

Teorema 1 (uniqueness of the reduced echelon form). *Each matrix is row equivalent to one and only one reduced echelon matrix.*

Teorema 2 (existence and uniqueness theorem). *A linear system is consistent iff the rightmost column of the augmented matrix is not a pivot column—that is, iff an echelon form of the augmented matrix has no row of the form*

$$[0 \quad \cdots \quad 0 \quad b], \qquad b \neq 0$$

If a linear system is consistent, then the solution set contains either (i) a unique solution, when there are no free variables, or (ii) infinitely many solutions, when there is at least one free variable.

2 The matrix equation $Ax = b$

Teorema 3. *If A is an $m \times n$ matrix, with columns $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_n$, and if $\boldsymbol{b}$ is in $\mathbb{R}^m$, the matrix equation*

$$A\boldsymbol{x} = \boldsymbol{b}$$

has the same solution set as the vector equation

$$x_1\boldsymbol{a}_1 + x_2\boldsymbol{a}_2 + \cdots + x_n\boldsymbol{a}_n$$

which, in turn, has the same solution set as the system of linear equations whose augmented matrix is

$$\begin{bmatrix} \boldsymbol{a}_1 & \boldsymbol{a}_2 & \cdots & \boldsymbol{a}_n & \boldsymbol{b} \end{bmatrix}$$

Teorema 4. *Let A be an $m \times n$ matrix. Then the following statements are logically equivalent.*

a. *For each $\boldsymbol{b}$ in $\mathbb{R}^m$, the equation $A\boldsymbol{x} = \boldsymbol{b}$ has a solution;*

b. *Each $\boldsymbol{b}$ is a linear combination of the columns of A;*

c. *The columns of A span $\mathbb{R}^m$;*

d. *A has a pivot in every row.*

Teorema 5 (properties of matrix-vector product). *If A is an $m \times n$ matrix, $\boldsymbol{u}$ and $\boldsymbol{v}$ are vectors in $\mathbb{R}^m$, and c is a scalar, then:*

a. $A(\boldsymbol{u} + \boldsymbol{v}) = A\boldsymbol{u} + A\boldsymbol{v}$;

b. $A(c\boldsymbol{u}) = c(A\boldsymbol{u})$.

3 Solution sets of linear systems

Teorema 6. *Suppose the equation $A\boldsymbol{x} = \boldsymbol{b}$ is consistent for some given $\boldsymbol{b}$, and let $\boldsymbol{p}$ be a solution. Then the solution set of $A\boldsymbol{x} = \boldsymbol{b}$ is the set of all vectors of the form $\boldsymbol{w} = \boldsymbol{p} + \boldsymbol{v}_h$, where $\boldsymbol{v}_h$ is any solution to the homogeneous equation $A\boldsymbol{x} = \boldsymbol{0}$.*

4 Linear independence

Teorema 7 (characterization of linearly dependent sets). *An indexed set $S = \{\boldsymbol{v}_1, \ldots, \boldsymbol{v}_p\}$ of two or more vectors is linearly dependent iff at least one of the vectors in S is a linear combination of the others. In fact, if S is linearly dependent and $\boldsymbol{v}_1 \neq= \boldsymbol{0}$, then some $\boldsymbol{v}_j$ (with $j > 1$) is a linear combination of the preceding vectors, $\boldsymbol{v}_1, \ldots, \boldsymbol{v}_{j-1}$.*

Teorema 8. *Any set $\{\boldsymbol{v}_1, \ldots, \boldsymbol{v}_p\}$ in $\mathbb{R}^n$ is linearly dependent if $p > n$.*

Teorema 9. *If a set $S = \{\boldsymbol{v}_1, \ldots, \boldsymbol{v}_p\}$ in $\mathbb{R}^n$ contains the zero vector, then the set is linearly dependent.*

5 The matrix of a linear transformation

Teorema 10. *Let $T : \mathbb{R}^n \to \mathbb{R}^m$ be a linear transformation. Then there exists a unique matrix A such that*

$$T(\boldsymbol{x}) = A\boldsymbol{x}, \qquad \forall \boldsymbol{x} \in \mathbb{R}^n$$

In fact,

$$A = [T(\boldsymbol{e}_1) \quad \cdots \quad T(\boldsymbol{e}_n)]$$

Teorema 11. *Let $T : \mathbb{R}^n \to \mathbb{R}^m$ be a linear transformation. Then T is one-to-one iff the equation $T(\boldsymbol{x}) = \boldsymbol{0}$ has only the trivial solution.*

Teorema 12. *Let $T : \mathbb{R}^n \to \mathbb{R}^m$ be a linear transformation, and let A be the standard matrix for T. Then:*

a. T maps $\mathbb{R}^n$ onto $\mathbb{R}^m$ iff the columns of A span $\mathbb{R}^m$;

b. T is one-to-one iff the columns of A are linearly independent.

II Matrix Algebra

1 Matrix operations

Teorema 1. *Let A, B, and C be matrices of the same size, and let r and s be scalars.*

a. $A + B = B + A$

b. $(A + B) + C = A + (B + C)$

c. $A + 0 = A$

d. $r(A + B) = rA + rB$

e. $(r + s)A = rA + sA$

f. $r(sA) = (rs)A$

Teorema 2 (properties of matrix multiplication). *Let A be an $m \times n$ matrix, and let B and C have sizes for which the indicated sums and products are defined.*

a. $A(BC) = (AB)C$ *(associative law of multiplication)*

b. $A(B + C) = AB + AC$ *(left distributive law)*

c. $(B + C)A = BA + CA$ *(right distributive law)*

d. $r(AB) = (rA)B = A(rB)$ *for any scalar* r

e. $I_m A = A = AI_n$

Teorema 3 (properties of matrix transposes). *Let* A *and* B *denote matrices whose sizes are appropriate for the following sums and products.*

a. $(A^T)^T = A$

b. $(A + B)^T = A^T + B^T$

c. For any scalar r, $(rA)^T = rA^T$

d. $(AB)^T = B^T A^T$

2 The inverse of a matrix

Definição 1. An $n \times n$ matrix A is said to be *invertible* if there exists an $n \times n$ matrix C such that $CA = I_n$ and $AC = I_n$. In this case, there is only one such matrix C, and we call it the *inverse* of A and denote it by A^{-1}. A matrix that is not invertible is sometimes called a *singular* matrix, and an invertible matrix is called a *nonsingular* matrix.

Teorema 4. *Let* $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. *If* $ad - bc \neq 0$, *then* A *is invertible and*

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

If $ad - bc = 0$, *then* A *is not invertible.*

Teorema 5. *If* A *is an invertible* $n \times n$ *matrix, then for each* $\boldsymbol{b}$ *in* $\mathbb{R}^n$, *the equation* $A\boldsymbol{x} = \boldsymbol{b}$ *has the unique solution* $\boldsymbol{x} = A^{-1}\boldsymbol{b}$.

Teorema 6. *a. If* A *is an invertible matrix, then* A^{-1} *is invertible and*

$$(A^{-1})^{-1} = A$$

b. If A *and* B *are* $n \times n$ *invertible matrices, then so is* AB, *and the inverse of* AB *is the product of the inverses in reverse order. That is,*

$$(AB)^{-1} = B^{-1}A^{-1}$$

c. If A *is an invertible matrix, then so is* A^T, *and the inverse of* A^T *is the transpose of* A^{-1}. *That is,*

$$(A^T)^{-1} = (A^{-1})^T$$

Teorema 7. *An* $n \times n$ *matrix* A *is invertible iff* A *is row equivalent to* I_n, *and in this case, any sequence of elementary row ops that reduce* A *to* I_n *also transforms* I_n *into* A^{-1}.

3 Characterizations of invertible matrices

Teorema 8 (the invertible matrix theorem). *Let A be a square $n \times n$ matrix. Then the following statements are equivalent.*

a. *A is an invertible matrix.*

b. *A is row equivalent to the $n \times n$ identity matrix.*

c. *A has n pivot positions.*

d. *The equation $Ax = 0$ has only the trivial solution.*

e. *The columns of A form a linearly independent set.*

f. *The transformation $x \mapsto Ax$ is one-to-one.*

g. *The equation $Ax = b$ has a unique solution for each b in $\mathbb{R}^n$*

h. *The columns of A span $\mathbb{R}^n$.*

i. *The linear transformation $x \mapsto Ax$ maps $\mathbb{R}^n$ onto $\mathbb{R}^n$.*

j. *There is an $n \times n$ matrix C such that $CA = I$.*

k. *There is an $n \times n$ matrix D such that $AD = I$.*

l. *A^T is an invertible matrix.*

m. *The columns of A form a basis for $\mathbb{R}^n$.*

n. $\operatorname{Col} A = \mathbb{R}^n$.

o. $\operatorname{rank} A = n$.

p. $\dim \ker A = 0$.

q. $\ker A = 0$.

r. *The number 0 is not an eigenvalue of A.*

Teorema 9. *Let $T : \mathbb{R}^n \to \mathbb{R}^n$ be a linear transformation and let A be the standard matrix for T. Then T is invertible iff A is an invertible matrix. In that case, the linear transformation S given by $S(x) = A^{-1}x$ is the unique function satisfying equations*

$$S(T(x)) = x \quad \text{for all } x \text{ in } \mathbb{R}^n \tag{1}$$

$$T(S(x)) = x \quad \text{for all } x \text{ in } \mathbb{R}^n \tag{2}$$

III Determinants

1 Introduction to determinants

Definição 1. For $n \geq 2$, the *determinant* of an $n \times n$ matrix A is defined recursively by the formula

$$\det A = \sum_{j=1}^{n} (-1)^{1+j} a_{1j} \det A_{1j}$$

where A_{1j} is the $(n-1) \times (n-1)$ matrix that results from deleting the first row and the j-th column of A. The determinant of a 1×1 matrix $[a]$ is defined to be a.

Teorema 1 (determinant by cofactor expansion). *The determinant of an $n \times n$ matrix A can be computed by a cofactor expansion across any row or down any column. The expansion across the i-th row is*

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in}.$$

The cofactor expansion down the j-th column is

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj}.$$

In either case, $C_{ij} = (-1)^{i+j} \det A_{ij}$, where A_{ij} is the $(n-1) \times (n-1)$ matrix that results from deleting the i-th row and j-th column of A.

Teorema 2. *If A is a triangular matrix, then* $\det A$ *is the product of the entries on the main diagonal.*

2 Properties of determinants

Teorema 3 (row operations). *Let A be a square matrix.*

a. *If a multiple of one row of A is added to another row to produce matrix B, then* $\det B = \det A$.

b. *If two rows of A are interchanged to produce B, then* $\det B = -\det A$.

c. *If one row of A is multiplied by k to produce B, then* $\det B = k \ \det A$.

Teorema 4. *A square matrix A is invertible iff* $\det A \neq 0$.

Teorema 5. *If A is an $n \times n$ matrix, then* $\det A^T = \det A$.

Teorema 6 (multiplicative property of determinants). *If A and B are $n \times n$ matrices, then* $\det AB = (\det A)\,(\det B)$.

IV Vector Spaces

1 Vector spaces and subspaces

Definição 1. A *vector space* is a nonempty set V of objects, called *vectors*, on which are defined two operations, called *vector addition* and *scalar multiplication*, that satisfy the following ten axioms below for all u, v, w in V and all scalars c and d.

1. $u + v$ is in V
2. $u + v = v + u$
3. $(u + v) + w = u + (v + w)$
4. There is a zero vector 0 in V such that $u + 0 = u$
5. For each u in V, there is a vector $-u$ in V such that $u + (-u) = 0$
6. cu is in V
7. $c(u + v) = cu + cv$
8. $(c + d)u = cu + du$
9. $c(du) = (cd)u$

10. $1u = u$

Definição 2. A *subspace* of a vector space V is a subset H of V that has three properties:

a. The zero vector of V is in H.

b. H is closed under vector addition. That is, for each u and v in H, the sum $u + v$ is in H.

c. H is closed under scalar multiplication. That is, for each u in H and each scalar c, the product cu is in H.

Teorema 1. *If $v_1, \ldots, v_p$ are in a vector space V, then Span$\{v_1, \ldots, v_p\}$ is a subspace of V.*

2 Null space, column space, row space, and linear transformations

Definição 3. The *null space* of a matrix A is the set of all vectors $\mathbf{x}$ such that $A\mathbf{x} = \mathbf{0}$.

Teorema 2. *The null space of an $m \times n$ matrix A is a subspace of $\mathbb{R}^n$.*

Definição 4. The *column space* of an $m \times n$ matrix A is the set of all linear combinations of the columns of A.

Teorema 3. *The column space of an $m \times n$ matrix A is a subspace of $\mathbb{R}^m$.*

Definição 5. A *linear transformation* T from a vector space V to a vector space W is a function from V to W that assigns to each vector $x \in V$ a unique vector $T(x) \in W$ such that for all vectors $u, v \in V$ and all scalars c, the following two properties hold:

1. $T(u + v) = T(u) + T(v)$
2. $T(cu) = cT(u)$

3 Linearly independent sets; bases

Definição 6. An indexed set of vectors $\{v_1, \ldots, v_p\}$ in a vector space V is said to be *linearly independent* if the vector equation

$$x_1v_1 + x_2v_2 + \cdots + x_pv_p = 0$$

has only the trivial solution $x_1 = x_2 = \cdots = x_p = 0$. The set is said to be *linearly dependent* if there exist weights $c_1, \ldots, c_p$, not all zero, such that

$$c_1v_1 + c_2v_2 + \cdots + c_pv_p = 0$$

Teorema 4. *An indexed set $v_1, \ldots, v_p$ of two or more vectors, with $v_1 \neq 0$, is linearly dependent iff some v_j (with $j > 1$) is a linear combination of the preceding vectors $v_1, \ldots, v_{j-1}$.*

Definição 7. Let H be a subspace of a vector space V. A *basis* for H is a linearly independent set in H that spans H.

Teorema 5 (spanning set theorem). *Let $S = \{v_1, \ldots, v_p\}$ be a set in a vector space V, and let $H =$ span$\{v_1, \ldots, v_p\}$.*

a. If v_k is a linear combination of the remaining vectors in S, then the set formed by $S - \{v_k\}$ still spans H.

b. If $H \neq 0$, some subset of S is a basis for H.

Teorema 6. *The pivot columns of a matrix A form a basis for Col A.*

Teorema 7. *If two matrices A and B are row equivalent, then their row spaces are the same. If B is in echelon form, the nonzero rows of B form a basis for the row space of A as well as for that of B.*

4 Coordinate systems

Teorema 8 (the unique representation theorem). *Let $\mathcal{B} = \{\boldsymbol{b}_1, \ldots, \boldsymbol{b}_n\}$ be a basis for a vector space V. Then for each $\boldsymbol{x}$ in V, there exists a unique set of scalars $c_1, \ldots, c_n$ such that*

$$\boldsymbol{x} = c_1\boldsymbol{b}_1 + \cdots + c_n\boldsymbol{b}_n$$

Teorema 9. *Let $\mathcal{B} = \{b_1, \ldots, b_n\}$ be a basis for a vector space V. Then the coordinate mapping $x \mapsto [x]_{\mathcal{B}}$ is a one-to-one linear transformation from V onto $\mathbb{R}^n$.*

Definição 8. Suppose $\mathcal{B} = \{b_1, \ldots, b_n\}$ is a basis for a vector space V. Then the *coordinates* of a vector x in V relative to the basis $\mathcal{B}$ are the weights $c_1, \ldots, c_n$ such that $x = c_1b_1 + \cdots + c_nb_n$.

Definição 9. A one-to-one linear transformation from a vector space V onto a vector space W is called an *isomorphism* from V to W. If there exists an isomorphism from V to W, we say that V and W are *isomorphic*.

5 The dimension of a vector space

Teorema 10. *If a vector space V has a basis $\mathcal{B} = \{b_1, \ldots, b_n\}$, then any set in V containing more than n vectors must be linearly dependent.*

Teorema 11. *If a vector space V has a basis of n vectors, then every basis of V must consist of exactly n vectors.*

Definição 10. If a vector space V is spanned by a finite set, then V is said to be *finite-dimensional*, and the dimension of V is the number of vectors in a basis for V. We denote the dimension of V by $\dim V$. The dimension of the zero vector space is defined to be zero. If V is not spanned by a finite set, then V is said to be *infinite-dimensional*.[1]

Teorema 12 (linearly independent sets can be expanded to bases). *Let H be a subspace of a finite-dimensional vector space V. Any linearly independent set in H can be expanded, if necessary, to a basis for H. Also, H if finite-dimensional and*

$$\dim H \leq \dim V$$

Teorema 13 (basis theorem). *Let V be a p-dimensional vector space, $p \geq 1$. Any linearly independent set of exactly p elements in V is automatically a basis for V. Any set of exactly p elements that spans V is automatically a basis for V.*

Definição 11. The *rank* of an $m \times n$ matrix A is the dimension of the column space of A. The *nullity* of A is the dimension of the null space of A.

Teorema 14 (rank theorem). *The dimensions of the column space and the null space of an $m \times n$ matrix A satisfy the equation*

$$\text{rank } A + \text{nullity } A = \text{number of columns in } A$$

Teorema 15. *Let $\mathcal{B} = \{b_1, \ldots, b_n\}$ and $\mathcal{C} = \{c_1, \ldots, c_n\}$ be bases of a vector space V. Then there is a unique $n \times n$ matrix $\underset{C \leftarrow B}{P}$ such that*

$$[\boldsymbol{x}_{\mathcal{C}}] = \underset{C \leftarrow B}{P}[\boldsymbol{x}_{\mathcal{B}}]$$

The columns of $\underset{C \leftarrow B}{P}$ are the $\mathcal{C}$-coordinate vectors of the vectors in the basis $\mathcal{B}$. That is,

$$\underset{C \leftarrow B}{P} = [[\boldsymbol{b}_1]_{\mathcal{C}} \ [\boldsymbol{b}_2]_{\mathcal{C}} \ \cdots [\boldsymbol{b}_n]_{\mathcal{C}}]$$

[1] Infinite-dimensional vector spaces are studied in Functional Analysis courses (phD level).

6 Change of basis

Teorema 16. *Let $\mathcal{B} = \{b_1, \ldots, b_n\}$ and $\mathcal{C} = \{c_1, \ldots, c_n\}$ be bases of a vector space V. Then there is a unique $n \times n$ matrix $\underset{C \leftarrow B}{P}$ such that*

$$[x]_{\mathcal{C}} = \underset{C \leftarrow B}{P} [x]_{\mathcal{B}}$$

The columns of $\underset{C \leftarrow B}{P}$ are the $\mathcal{C}$-coordinate vectors of the vectors in the basis $\mathcal{B}$. That is,

$$\underset{C \leftarrow B}{P} = [[b]_{\mathcal{C}} \, [b_2]_{\mathcal{C}} \, \cdots [b_n]_{\mathcal{C}}]$$

The columns of $\underset{C \leftarrow B}{P}$ are linearly independent because they are the coordinate vectors of a basis. Hence, $\underset{C \leftarrow B}{P}$ is an invertible matrix, and

$$\underset{B \leftarrow C}{P} = \underset{C \leftarrow B}{P}^{-1}$$

V Eigenvalues and Eigenvectors

1 Eigenvectors and eigenvalues

Definição 1. An *eigenvector* of an $n \times n$ matrix A is a nonzero vector x such that $Ax = \lambda x$ for some scalar λ. A scalar λ is called an *eigenvalue* of A if there is a nontrivial solution x of the equation $Ax = \lambda x$. In this case, we say that x is an eigenvector of A that corresponds to the eigenvalue λ.

Definição 2. The *eigenspace* of an $n \times n$ matrix A corresponding to an eigenvalue λ, denoted by $E(A, \lambda)$, is the set of all eigenvectors corresponding to λ, together with the zero vector. That is,

$$E(A, \lambda) = \{x \in \mathbb{R}^n : (A - \lambda I)x = 0\} = \ker(A - \lambda I)$$

Teorema 1. *The eigenvalues of a triangular matrix are the entries on its main diagonal.*

Teorema 2 (eigenvectors corresponding to distinct eigenvalues are linearly independent). *If $v_1, \ldots, v_r$ are eigenvectors that correspond to distinct eigenvalues $\lambda_1, \ldots, \lambda_r$ of an $n \times n$ matrix A, then the set $\{v_1, \ldots, v_r\}$ is linearly independent.*

Demonstração. □

2 The characteristic equation

Teorema 3 (properties of determinants). *Let A and B be $n \times n$ matrices.*

a. A is invertible iff $\det A \neq 0$

b. $\det AB = (\det A)(\det B)$

c. $\det A^T = \det A$

d. If A is triangular, then $\det A$ *is the product of the entries on the main diagonal of A.*

e. A row replacement operation on A does not change the determinant. A row interchange changes the sign of the determinant. A row scaling also scales the determinant by the same scalar factor.

Definição 3. The *characteristic polynomial* of an $n \times n$ matrix A is the polynomial defined by the formula

$$p(\lambda) \stackrel{\text{def}}{=} \det(A - \lambda I).$$

The *characteristic equation* of A is the equation $p(\lambda) = 0$.

Observação. A scalar λ is an eigenvalue of A iff λ is a root of the characteristic polynomial $p(\lambda)$. The (algebraic) multiplicity of an eigenvalue λ is the multiplicity of λ as a root of $p(\lambda)$.

Definição 4. If A and B are $n \times n$ matrices, then A and B are *similar* if there is an invertible matrix P such that $A = PBP^{-1}$.

Teorema 4. *If $n \times n$ matrices A and B are similar, then they have the same characteristic polynomial and hence the same eigenvalues (with the same multiplicities).*

3 Diagonalization

Teorema 5 (diagonalization theorem). *An $n \times n$ matrix A is diagonalizable iff A has n linearly independent eigenvectors. In fact, $A = PDP^{-1}$, with D a diagonal matrix, iff the columns of P are the n linearly independent eigenvectors of A. In this case, the diagonal entries of D are the eigenvalues of A that correspond, respectively, to the eigenvectors in P.*

Teorema 6. *An $n \times n$ matrix with n distinct eigenvalues is diagonalizable.*

Teorema 7. *Let A be an $n \times n$ matrix whose distinct eigenvalues are $\lambda_1, \ldots, \lambda_p$.*

a. *For $1 \leq k \leq p$, the dimension of the eigenspace for λ_k is less than or equal to the multiplicity of the eigenvalue λ_k.*

b. *The matrix A is diagonalizable iff the sum of the dimensions of the eigenspaces equals n, and this happens iff (i) the characteristic polynomial factors completely into linear factors and (ii) the dimension of the eigenspace for each λ_k equals the multiplicity of λ_k.*

c. *If A is diagonalizable and $\mathcal{B}_k$ is a basis for the eigenspace corresponding to λ_k for each k, then the total collection of vectors in $\mathcal{B}_1, \ldots, \mathcal{B}_p$ forms an eigenvector basis for $\mathbb{R}^n$.*

Observação. The *trace* of an $n \times n$ matrix A, denoted by tr A, is the sum of the diagonal entries of A. If $\lambda_1, \ldots, \lambda_n$ are the eigenvalues of A, then tr $A = \lambda_1 + \lambda_2 + \cdots + \lambda_n$.

Observação. The determinant of an $n \times n$ matrix A is the product of the eigenvalues of A:

$$\det A = \lambda_1 \lambda_2 \cdots \lambda_n$$

VI Orthogonality and Least Squares

1 Inner product, length, and orthogonality

Teorema 1. *Let u, v, and w be vectors in $\mathbb{R}^n$, and let c be a scalar. Then*

a. $u \cdot v = v \cdot u$

b. $(u + v) \cdot w = u \cdot w + v \cdot w$

c. $(cu) \cdot v = c(u \cdot v) = u \cdot (cv)$

d. $u \cdot u \geq 0$, *and* $u \cdot u = 0 \iff u = 0$

Definição 1. The *length* (or *norm*) of v is the nonnegative scalar defined by

$$\|v\| \stackrel{\text{def}}{=} \sqrt{v \cdot v}$$

Definição 2. For u and v in $\mathbb{R}^n$, the *distance* between u and v is the nonnegative scalar defined by

$$d(u, v) \stackrel{\text{def}}{=} \|u - v\|$$

Definição 3. Two vectors u and v in $\mathbb{R}^n$ are *orthogonal* if $u \cdot v = 0$.

Teorema 2 (pythagorean theorem). *Two vectors u and v are orthogonal iff $\|u + v\|^2 = \|u\|^2 + \|v\|^2$.*

Definição 4. If a vector z is orthogonal to every vector in a subspace W of $\mathbb{R}^n$, then z is said to be *orthogonal* to W. The set of all vectors that are orthogonal to W is called the *orthogonal complement* of W and is denoted by $W^{\perp}$.

Teorema 3. *Let A be an $m \times n$ matrix. The orthogonal complement of the row space of A is the null space of A, and the orthogonal complement of the column space of A is the null space of A^T:*

$$(\textit{Row } A)^{\perp} = \textit{Nul } A \quad \textit{and} \quad (\textit{Col } A)^{\perp} = \textit{Nul } A^T$$

2 Orthogonal sets

Definição 5. A set of vectors $u_1, \ldots, u_p$ in $\mathbb{R}^n$ is an *orthogonal set* if each pair of distinct vectors in the set is orthogonal. That is, $u_i \cdot u_j = 0$ whenever $i \neq j$.

Teorema 4. *If $S = \{u_1, \ldots, u_p\}$ is an orthogonal set of nonzero vectors in $\mathbb{R}^n$, then S is linearly independent and hence is a basis for the subspace spanned by S.*

Definição 6. An *orthogonal basis* for a subspace W of $\mathbb{R}^n$ is a basis for W that is also an orthogonal set.

Teorema 5. *Let $\{u_1, \ldots, u_p\}$ be an orthogonal basis for a subspace W of $\mathbb{R}^n$. For each y in W, the weights in the linear combination*

$$y = c_1 u_1 + \cdots + c_p u_p$$

are given by

$$c_j = \frac{y \cdot u_1}{u_1 \cdot u_1} \qquad (j = 1, \ldots, p)$$

Definição 7 (orthogonal projection). For a vector y in $\mathbb{R}^n$ and a subspace W of $\mathbb{R}^n$, the *orthogonal projection* of y onto W, denoted by $\hat{y} = \text{proj}_W\, y$, is the vector in W given by

$$\text{proj}_W\, y = \frac{y \cdot u_1}{u_1 \cdot u_1} u_1 + \cdots + \frac{y \cdot u_p}{u_p \cdot u_p} u_p$$

where $\{u_1, \ldots, u_p\}$ is any orthogonal basis of W.

Definição 8. A set $\{u_1, \ldots, u_p\}$ of vectors in $\mathbb{R}^n$ is an *orthonormal set* if it is an orthogonal set of unit vectors. An *orthonormal basis* for a subspace W of $\mathbb{R}^n$ is a basis for W that is also an orthonormal set.

Teorema 6. *An $m \times n$ matrix U has orthonormal columns iff $U^T U = I$.*

Teorema 7. *Let U be an $m \times n$ matrix with orthonormal columns, and let x and y be in $\mathbb{R}^n$. Then*

a. $\|Ux\| = \|x\|$

b. $(Ux) \cdot (Uy) = x \cdot y$

c. $(Ux) \cdot (Uy) = 0 \iff x \cdot y = 0$

Definição 9. An *orthogonal matrix* is a square invertible matrix U such that $U^{-1} = U^T$.

Corolário 7.1. *By* (6)*, an $n \times n$ matrix U is orthogonal iff it has orthonormal columns.*

3 Orthogonal projections

Teorema 8 (orthogonal decomposition theorem). *Let W be a subspace of $\mathbb{R}^n$. Then each y in $\mathbb{R}^n$ can be written uniquely in the form*

$$y = \hat{y} + z$$

where $\hat{y}$ is in W and z is in $W^\perp$. In fact, if $\{u_1, \ldots, u_p\}$ is any orthogonal basis of W, then

$$\hat{y} = \frac{y \cdot u_1}{u_1 \cdot u_1} u_1 + \cdots + \frac{y \cdot u_p}{u_p \cdot u_p} u_p$$

and

$$z = y - \hat{y}$$

.

Teorema 9 (best approximation theorem). *Let W be a subspace of $\mathbb{R}^n$, let y be any vector in $\mathbb{R}^n$, and let $\hat{y}$ be the orthogonal projection of y onto W. Then $\hat{y}$ is the closest point in W to y, in the sense that*

$$\|y - \hat{y}\| < \|y - v\|$$

for all v in W distinct from $\hat{y}$.

Teorema 10 (orthogonal projection theorem). *If $\{\boldsymbol{u}_1, \ldots, \boldsymbol{u}_p\}$ is an orthonormal basis for a subspace W of $\mathbb{R}^n$, then*

$$\mathit{proj}_W \boldsymbol{y} = (\boldsymbol{y} \cdot \boldsymbol{u}_1)\boldsymbol{u}_1 + (\boldsymbol{y} \cdot \boldsymbol{u}_2)\boldsymbol{u}_2 + \cdots + (\boldsymbol{y} \cdot \boldsymbol{u}_p)\boldsymbol{u}_p$$

If $U = [\boldsymbol{u}_1 \quad \boldsymbol{u}_2 \quad \cdots \quad \boldsymbol{u}_p]$*, then*

$$\mathit{proj}_W \boldsymbol{y} = UU^T \boldsymbol{y}$$

for all y in $\mathbb{R}^n$.

4 The Gram-Schmidt process

Teorema 11 (Gram-Schmidt process). *Given a basis $\{x_1, \ldots, x_p\}$ for a nonzero subspace W of $\mathbb{R}^n$, define*

$$\begin{aligned}
v_1 &= x_1 \\
v_2 &= x_2 - \frac{x_2 \cdot v_1}{v_1 \cdot v_1} v_1 \\
v_3 &= x_3 - \frac{x_3 \cdot v_1}{v_1 \cdot v_1} v_1 - \frac{x_3 \cdot v_2}{v_2 \cdot v_2} v_2 \\
&\vdots \\
v_p &= x_p - \frac{x_p \cdot v_1}{v_1 \cdot v_1} v_1 - \frac{x_p \cdot v_2}{v_2 \cdot v_2} v_2 - \cdots - \frac{x_p \cdot v_{p-1}}{v_{p-1} \cdot v_{p-1}} v_{p-1}
\end{aligned}$$

Then $\{v_1, \ldots, v_p\}$ is an orthogonal basis for W. In addition

$$\text{span}\{v_1, \ldots, v_k\} = \text{span}\{x_1, \ldots, x_k\}$$

for $1 \leq k \leq p$.

Teorema 12 (QR factorization). *If A is an $m \times n$ matrix with linearly independent columns, then A can be factored as $A = QR$, where Q is an $m \times n$ matrix whose columns form an orthonormal basis for* $\text{Col}\, A$ *and R is an $n \times n$ upper triangular invertible matrix with positive entries on its diagonal.*

5 Least-squares problems

Definição 10. If A is an $m \times n$ matrix and b is a vector in $\mathbb{R}^m$, then a *least-squares solution* of the equation $Ax = b$ is a vector $\hat{x}$ in $\mathbb{R}^n$ such that

$$\|A\hat{x} - b\| \leq \|Ax - b\|$$

for all x in $\mathbb{R}^n$.

Teorema 13. *The set of least-squares solutions of $Ax = b$ coincides with the nonempty set of solutions of the normal equations $A^T Ax = A^T b$.*

Teorema 14. *Let A be an $m \times n$ matrix. The following statements are logically equivalent:*

a. *The equation $Ax = b$ has a unique least-squares solution for each b in $\mathbb{R}^m$.*

b. *The columns of A are linearly independent.*

c. *The matrix $A^T A$ is invertible.*

When these statements are true, the least-squares solution $\hat{x}$ is given by

$$\hat{x} = (A^T A)^{-1} A^T b$$

Teorema 15. *Given an $m \times n$ matrix A with linearly independent columns, let $A = QR$ be a QR factorization of A. Then, for each b in $\mathbb{R}^m$, the equation $Ax = b$ has a unique least-squares solution, given by*

$$\hat{x} = R^{-1} Q^T b$$

VII Symmetric Matrices and Quadratic Forms

Definição 1. A *symmetric* matrix is a square matrix A such that $A^T = A$.

Teorema 1. *If A is symmetric, then any two eigenvectors from different eigenspaces are orthogonal.*

Definição 2. An $n \times n$ matrix A is *orthogonally diagonalizable* if there are an orthogonal matrix P and a diagonal matrix D such that $A = PDP^T = PDP^{-1}$.

Teorema 2. *An $n \times n$ matrix A is orthogonally diagonalizable iff A is a symmetric matrix.*

Definição 3. The *spectrum* of an $n \times n$ matrix A is the set of eigenvalues of A.

Teorema 3 (spectral theorem for symmetric matrices). *An $n \times n$ symmetric matrix A has the following properties:*

a. A has n real eigenvalues, counting multiplicities.

b. For each eigenvalue, the dimension of the eigenspace equals the multiplicity of the eigenvalue.

c. The eigenspaces are mutually orthogonal, in the sense that eigenvectors corresponding to different eigenvalues are orthogonal.

d. A is orthogonally diagonalizable; that is, there are an orthogonal matrix P and a diagonal matrix D such that $A = PDP^T = PDP^T = PDP^{-1}$.

1 Quadratic forms

Definição 4. A *quadratic form* on $\mathbb{R}^n$ is a function $Q : \mathbb{R}^n \to \mathbb{R}$ such that there is an $n \times n$ symmetric matrix A such that $Q(x) = x^T Ax$ for all x in $\mathbb{R}^n$.

Definição 5. If x is a variable vector in $\mathbb{R}^n$, then a *change of variable* is an equation of the form $x = Py$, where P is an invertible $n \times n$ matrix and y is a new variable vector in $\mathbb{R}^n$.

Teorema 4 (principal axes theorem). *Let A be an $n \times n$ symmetric matrix. Then there is an orthogonal change of variable, $\boldsymbol{x} = P\boldsymbol{y}$, that transforms the quadratic form $\boldsymbol{x}^T A\boldsymbol{x}$ into a quadratic form $\boldsymbol{y}^T D\boldsymbol{y}$ with no cross-product term.*

Definição 6. A quadratic form Q is

a. *positive definite* if $Q(x) > 0$ for all $x \neq 0$.

b. *negative definite* if $Q(x) < 0$ for all $x \neq 0$.

c. *indefinite* if $Q(x)$ assumes both positive and negative values.

Observação. If $Q(x) \geq 0$ for all x in $\mathbb{R}^n$, then Q is said to be *positive semidefinite*. If $Q(x) \leq 0$ for all x in $\mathbb{R}^n$, then Q is said to be *negative semidefinite*.

Teorema 5 (quadratic forms and eigenvalues). *Let A be an $n \times n$ symmetric matrix. Then a quadratic form $Q(\boldsymbol{x}) = \boldsymbol{x}^T A\boldsymbol{x}$ is*

a. positive definite ($Q(\boldsymbol{x}) > 0$ for all $\boldsymbol{x} \neq 0$) iff the eigenvalues of A are all positive.

b. negative definite ($Q(\boldsymbol{x}) < 0$ for all $\boldsymbol{x} \neq 0$) iff the eigenvalues of A are all negative.

c. indefinite ($Q(\boldsymbol{x})$ assumes both positive and negative values) iff A has both positive and negative eigenvalues.

2 Constrained optimization

Teorema 6. *Let A be a symmetric matrix, and define m and M as $m = \min\{x^T Ax : |x| = 1\}$ and $M = \max\{x^T Ax : |x| = 1\}$. Then M is the greatest eigenvalue λ_1 of A and m is the least eigenvalue of A. The value of $x^T Ax$ is M when x is a unit vector u_1 corresponding to M. The value of $x^T Ax$ is m when x is a unit vector u_2 corresponding to m.*

Teorema 7. *Let A, λ_1, and u_1 be as in* (6)*. Then the maximum value of $x^T Ax$ subject to the constraints*

$$x^T x = 1, \quad x^T u_1 = 0$$

is the second greatest eigenvalue, λ_2, and this maximum value is attained when x is an eigenvector u_2 corresponding to λ_2.

Teorema 8. *Let A be a symmetric $n \times n$ matrix with orthogonal diagonalization $A = PDP^{-1}$, where the entries on the diagonal of D are $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$ and where the columns of P are the corresponding orthonormal eigenvectors $u_1, u_2, \ldots, u_n$. Then, for $k = 2, \ldots, n$, the maximum value of $x^T Ax$ subject to the constraints*

$$x^T x = 1, \quad x^T u_1 = 0, \quad \ldots, \quad x^T u_{k-1} = 0$$

is the eigenvalue λ_k, and this maximum value is attained at $x = u_k$.