

DSA5103 Assignment 4

Instructions Due on April 21, 11:59pm. The final exam will take the same format as this assignment.

1. (5 marks) Given $b \in \mathbb{R}^n$, $c \in \mathbb{R}$, show that the half-space $H = \{x \in \mathbb{R}^n \mid b^T x \leq c\}$ is a convex set.
2. (5 marks) Given two convex sets C_1 and C_2 in \mathbb{R}^n ,
 - (a) Show that $C = C_1 \cap C_2$ is also convex (\cap denotes the intersection of two sets).
 - (b) Give an example where $C_1 \cup C_2$ is not convex but C_1 and C_2 are convex (\cup denotes the union of two sets).
3. (5 marks) Let $C = \{x \in \mathbb{R}^n \mid \|x\|_2 \leq r\}$.
 - (a) Give the formula for the projection onto C : $\Pi_C(x)$.
 - (b) When $r = 2$ and $x = (3, 1)^T$, compute $\Pi_C(x)$.
4. (5 marks) Consider the function

$$f(x) = 100(x_2 - x_1^2)^2 + (1 - x_1)^2, \quad x = (x_1; x_2) \in \mathbb{R}^2.$$

- (a) Compute the gradient and Hessian of the function.
 - (b) Show that $x^* = (1, 1)^T$ is a local minimizer of the function. [Hint: consider the sufficient condition]
 - (c) Show that $x^* = (1, 1)^T$ is the only local minimizer of the function. [Hint: consider the necessary condition]
5. (10 marks) Consider the function

$$f(x) = (x_1 + x_2^2)^2, \quad x = (x_1; x_2) \in \mathbb{R}^2.$$

- (a) At $x^{(0)} = (1, 0)^T$, compute the steepest descent direction.
 - (b) At $x^{(0)} = (1, 0)^T$, construct another descent direction (different from the steepest descent direction).

(c) Start from $x^{(0)} = (1, 0)^T$, apply steepest descent method with exact line search and determine the next iterate $x^{(1)}$.

(d) Start from $x^{(0)} = (1, 0)^T$, apply steepest descent method with backtracking line search and determine the next iterate $x^{(1)}$. In particular, set $\bar{\alpha} = 1, \rho = 0.9, c_1 = 10^{-4}$ and do backtracking line search until the Armijo condition holds

$$f(x^{(k)} + \alpha p^{(k)}) \leq f(x^{(k)}) + c_1 \alpha \nabla f(x^{(k)})^T p^{(k)}$$

(e) At $x^{(0)} = (1, 0)^T$, compute the Newton direction.

(f) Start from $x^{(0)} = (1, 0)^T$, apply pure Newton's method (step size = 1) and determine the next iterate $x^{(1)}$.

6. (10 marks) (a) Let $f(x) = \frac{1}{2}x^2 + 4x, x \in \mathbb{R}$. Write down the effective domain of f . Compute its conjugate function f^* .

(b) Let $f(x) = -\sum_{i=1}^n \log x_i, x \in \mathbb{R}^n$. Write down the effective domain of f . Compute its conjugate function f^* . ($\log x = \ln x, (\log x)' = \frac{1}{x}$.)

7. (10 marks) Let $f(x) = \lambda \|x\|_2, x \in \mathbb{R}^n$.

(a) Compute its conjugate function f^* .

(b) Give the formula for $P_{f^*}(\cdot)$ (the proximal mapping of f^*).

(c) Give the formula for $P_f(\cdot)$ (the proximal mapping of f). [Hint: Moreau decomposition]

(d) Set $\lambda = 1, b = (1, 1)^T$, compute $P_{f^*}(b)$ and $P_f(b)$.

8. (10 marks) Consider the problem

$$\min_{x \in \mathbb{R}^n} \quad \frac{1}{2} \|Ax - b\|_2^2 \quad \text{s.t.} \quad \|x\|_\infty \leq r$$

where $r > 0, A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$ are given, $\|x\|_\infty = \max_i |x_i|$.

(a) Transform the above problem into the structure

$$\min_x \quad f(x) + g(x)$$

where f is differentiable and g is non-differentiable.

(b) Give the PG (proximal gradient) iterations.

(c) Set $r = 1, \alpha = 1$ (step length), $m = n = 2, A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, b = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$. Initialize $x^{(0)} = (0, 0)^T$. Compute the first iteration $x^{(1)}$ of PG.

9. (10 marks) Consider the problem

$$\min_{x \in \mathbb{R}^n} \|x\|_1 \quad \text{s.t.} \quad \|x - b\|_2 \leq r$$

where $r > 0$, and $b \in \mathbb{R}^n$ are given, $\|\cdot\|_1, \|\cdot\|_2$ denote the ℓ_1, ℓ_2 norms.

- (a) Transform the above problem into the 2-block separable structure

$$\min_{y, z} f(y) + g(z) \quad \text{s.t.} \quad Ay + Bz = c$$

that ADMM can handle.

- (b) Give the ADMM iterations.

- (c) Set $r = 1, \sigma = 1, \tau = 1, m = n = 2, b = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$. Initialize every variable by zero. Compute the first iteration of ADMM.

10. (10 marks) Apply coordinate descent method for

$$\min_{x=(x_1; x_2) \in \mathbb{R}^2} f(x_1, x_2) = x_1^4 - 4x_1x_2 + 5x_2^2 - 10x_2$$

with initial point $x^{(0)} = (0; 0)$. Find $x^{(1)}$ and $x^{(2)}$.

11. (10 marks) Two supplies $a_1 = 0.4$ and $a_2 = 0.6$ of goods must be transport to meet two demands $b_1 = 0.2$ and $b_2 = 0.8$ of customers. The cost of transporting one unit of i th good to j th customer is c_{ij} ($c_{11} = 2, c_{21} = 1, c_{12} = 3, c_{22} = 4$). Determine the optimal transportation plan (x_{ij} unit of i th good transported to j th customer) to minimize the cost.

- (a) Form the above problem into an optimization problem.

- (b) Give the KKT conditions.

- (c) Construct two feasible points (i.e., two feasible transportation plans).

- (d) For the two feasible transportation plans in (c), which one is better (i.e., with less cost)?

12. (10 marks) Consider the problem

$$\begin{aligned} \max_{x_1, x_2} \quad & 6x_1 + 4x_2 - 13 - x_1^2 - x_2^2 \\ \text{s.t.} \quad & x_1 + x_2 \leq 3 \end{aligned}$$

- (a) Convert the problem into an equivalent minimization problem. Solve it graphically and give the solution x_1^*, x_2^* .

- (b) Give the KKT conditions.

- (c) Derive the Lagrange dual problem.

- (d) Solve the Lagrange dual problem and give the dual solution y^* .

- (e) Verify that x_1^*, x_2^*, y^* satisfy the KKT conditions.