

Appendix S1 – Supplementary Information

Capitani L. *et al.*

Beaver-engineering amplifies dissolved organic carbon concentration

This document contains the material supporting the main findings of the manuscript (tested hypotheses of the structural causal model, priors, residuals, posterior predictive checks, posterior parameter distributions, a stream-level classification of ΔDOC , and an alternative SCM including lateral stream–wetland connectivity). The full description of data collection, variable derivation and model specification is given in the Methods of the main manuscript.

SCM: tested hypothesis

Table S1: Tested hypotheses for the structural causal model (SCM) represented in Figure 1b of the main document.

Causal path	Predictor	Response	Hypothesis
1	DOC upstream	ΔDOC	If DOC upstream increases, then ΔDOC decreases because beaver ponds can act as sink, retaining or transforming DOC through processes such as microbial degradation, sedimentation, and photodegradation, thereby reducing downstream DOC concentrations relative to upstream inputs (Larson et al. 2007; Wohl 2013; Johnston 2014).
2	Macrophytes	ΔDOC	If macrophyte abundance increases, we expect ΔDOC to increase because the decomposition of macrophyte biomass releases labile dissolved organic carbon, enhancing downstream concentrations (Reitsema et al. 2018).
3	Phytoplankton	ΔDOC	If phytoplankton abundance increases, then ΔDOC increases is likely to increase, because phytoplankton contribute to DOC content through exudation during growth and release upon cell lysis, thereby enhancing downstream DOC levels relative to upstream inputs (Søndergaard et al. 1985; Sell and Overbeck 1992).
4	Soil litter cover	ΔDOC	If soil litter cover increases, then ΔDOC increases as decomposition of litter releases dissolved organic compounds into the water (Aitkenhead et al. 1999; Cincotta et al. 2019).

continued on next page

Table S1 (continued)

Causal path	Predictor	Response	Hypothesis
5	Water residence time	ΔDOC	If water residence time increases, then ΔDOC increases, because longer water residence times allow for more extensive microbial decomposition and processing of dissolved organic carbon, enhancing downstream DOC concentrations (Jepsen et al. 2019; Abbasi et al. 2024).
6	Solar radiation	Phytoplankton	If solar radiation increases then phytoplankton abundance is expected to increase because greater light availability enhances photosynthetic rates (Reynolds 2006).
7	Solar radiation	Macrophytes	If solar radiation increases, then macrophyte abundance increases because higher light availability enhances photosynthetic activity and supports plant growth (Feuchtmayr et al. 2009).
8	Solar radiation	Litter cover	If solar radiation increases, as in summer, then soil litter cover decreases because lower rates of tree leaf fall coincide with enhanced decomposition driven by higher temperatures and solar radiation (Gessner et al. 1999).
9	Water residence time	Macrophytes	If water residence time increases, macrophyte abundance may increase because longer residence time allows for higher nutrient retention, promoting plant growth and increasing the likelihood of macrophyte establishment and persistence (Manolaki et al. 2021).
10	Water residence time	Phytoplankton	If water residence time increases, phytoplankton abundance may increase because longer residence time allows for greater nutrient retention and stabilization, which can support enhanced phytoplankton growth, particularly in systems with nutrient limitation (Finger et al. 2007).
11	Channel gradient (terrain slope)	Water residence time	If valley slope increases, then water residence time decrease because steeper terrain causes water to move more quickly through the landscape, reducing the time available for subsurface and surface interactions (McGuire et al. 2005).
12	Max dam height	Water residence time	If dam height increases, water residence time increases because taller dams impound more water, expanding ponded areas and slowing streamflow, which prolongs the time water remains within the stream system (Majerova et al. 2015; Dittbrenner et al. 2022).
13	Number of dams	Water residence time	If the number of beaver dams increases, water residence time is expected to increase because dams slow down streamflow and create ponded areas that retain water for longer periods (Westbrook et al. 2006; Nyssen et al. 2011; Majerova et al. 2015).

Structural Causal Model: the priors

To ensure the model was not over-constrained by prior knowledge, we used weakly informative priors for all parameters. This approach allows the data to inform the model without assuming strong beliefs about the parameter values beforehand (Figure S1). We note, however, that for macrophytes, phytoplankton, and litter cover—variables measured in 16 of the 176 streams—we used more informative priors that incorporate established ecological knowledge about their influence on DOC concentrations (Figure S1 panel b).




Figure S1: Half-eye plots display the prior distributions used in the model. The shaded areas (slabs) represent the probability density of each prior distribution, illustrating their shape and spread. The central dot indicates the median of the distribution, while the thick and thin horizontal lines represent the 66% and 95% credible intervals, respectively. Text labels to the left of each distribution indicate the corresponding prior specification.

Structural Causal Model: the residuals

We can evaluate the model fit by looking at the difference between observed data and their predictions. That are the residuals (r_i).

$$r_i = y_i - y_i^{pred} \quad (S1)$$

If the model is a reasonable representation of the observational process that originates the data then the residuals should look roughly randomly scattered in comparison to a horizontal line.




Figure S2: Residual plot showing the relationship between model-predicted ΔDOC values (mg L^{-1}) and corresponding residuals. Points represent the difference between observed and predicted values; a random scatter around zero (grey line) indicates good model fit and no systematic bias in predictions.

Structural Causal Model: posterior predictive checks

Posterior predictive checks assess how well the fitted SCM model reproduces observed data by comparing the observed values to data simulated from the model's posterior predictive distribution.

Structural Causal Model: posterior parameters distribution

Structural Causal Model: stream-level classification of ΔDOC

To avoid conflating the posterior predictive probability of a response (Figure 2a, main document) with the proportion of streams showing that response, we classified each stream individually. For each of the 179 streams we computed the posterior distribution of the expected ΔDOC (population-level effects plus the stream-level intercept, averaged over the stream's observations; 2,000 posterior draws) and assigned the stream to the region (sink $< -0.5 \text{ mg L}^{-1}$, ROPE $\pm 0.5 \text{ mg L}^{-1}$, source $> 0.5 \text{ mg L}^{-1}$) that held the majority of its posterior mass. The expected ΔDOC of all 179 streams fell within the ROPE (posterior medians from -0.40 to 0.38 mg L^{-1} ; median width of the 90% credible interval 0.41 mg L^{-1} ; Figure S5), reflecting the small average effect (intercept 0.00 mg L^{-1} ; 95% BCI -0.12 ; 0.12) and the small between-stream standard deviation (0.05 mg L^{-1} ; 95% BCI 0.00 ; 0.14). The heavy-tailed posterior predictive distribution of Figure 2a (Student- t residual with $\nu \approx 2$) thus captures the variability of individual samplings around a null average rather than consistent stream-level differences.




Figure S3: Posterior predictive checks of the mean and standard deviation statistics for the response variable delta DOC, with 1000 posterior draws used to generate the predictive distribution. This plot provides a visual assessment of the model's fit by comparing the observed data (dark blue) against the model's predictions (light blue) from the posterior distribution.




Figure S4: Posterior parameters distributions of the fitted SCM. Median shown with point; 66% and 95% credible intervals shown with black bars. Parameters that have more than 90% probability mass on either side of zero are considered to have a positive or negative influence on the response variable. a) The direct effects on ΔDOC b) The indirect effects on ΔDOC .




Figure S5: Stream-level ΔDOC . Blue points and bars: posterior median and 90% credible interval of the model-based expected ΔDOC of each stream (ranked by posterior median). Orange crosses: observed stream-averaged ΔDOC , shown for reference (one value of 29.9 mg L^{-1} is clipped at 5 mg L^{-1}). The grey band is the region of practical equivalence ($\pm 0.5 \text{ mg L}^{-1}$).

Alternative structural causal model: lateral stream–wetland connectivity

Because a DAG encodes a specific set of causal assumptions, we also tested an alternative, extended SCM that incorporates the lateral connectivity between the stream and adjacent wetlands as an additional driver of water residence time (WRT). Hydrologically connected off-channel storage zones (wetlands, floodplains) exchange water with the channel and are known to extend the residence time of water in the system (transient-storage theory; Bencala and Walters 1983). We therefore hypothesized that beaver-pond WRT varies with the degree of lateral stream–wetland connectivity and, consequently, that connectivity indirectly influences ΔDOC via WRT and primary producers.

Connectivity variable. For each site, we first determined whether a wetland was present within a 50 m buffer around the river reach between the upstream and downstream sampling locations, using the wetland layer (*Feuchtgebiet*: marshes, bogs, swamp forests, wet meadows, floodplains and fens) of the swissTLM3D topographic landscape model (SwissTopo 2024). For sites with a wetland, we then assessed visually on the swisstopo maps and orthophotos whether the wetland was hydrologically connected to the river water, or separated from it by roads, trails, tree lines or other barriers. Each site was thereby assigned to one of three categories: *No wetland* (no wetland within the buffer; $n = 136$ sites), *Wetland, not connected* (wetland present but not directly connected to the river; $n = 11$), and *Wetland, connected* (wetland present and directly connected to the river; $n = 13$). Twenty-two sites could not be classified and were excluded from this analysis (310 observations retained). Note that most of these wetlands are themselves strongly influenced by beaver damming and would likely not exist without beavers.

DAG and identification. In the extended DAG (Figure S6) lateral connectivity has no parents, hence there are no open back-door paths between connectivity and ΔDOC and the

total effect of connectivity on ΔDOC is identified without adjustment (empty adjustment set, *dagitty*). The causal influence flows through two front-door paths: connectivity $\rightarrow$ WRT $\rightarrow$ macrophytes $\rightarrow \Delta\text{DOC}$ and connectivity $\rightarrow$ WRT $\rightarrow$ phytoplankton $\rightarrow \Delta\text{DOC}$. Because the total effect of WRT on ΔDOC is likewise unconfounded, we fitted a parsimonious two-equation Bayesian model in *brms*: (i) ΔDOC (with the 0.283 mg L^{-1} measurement error, Student-*t* likelihood, site-level random intercept) regressed on WRT; (ii) WRT regressed on number of dams, maximum dam height, channel gradient, solar radiation and lateral connectivity (reference level: *No wetland*). Missing values in WRT and dam height were imputed within the model (`mi()`). Four chains were run ($\hat{R} \leq 1.005$). The total effect of each connectivity level on ΔDOC was computed from the posterior draws as the product of the connectivity $\rightarrow$ WRT contrast and the WRT $\rightarrow \Delta\text{DOC}$ coefficient.

Results. Relative to sites without wetland, the presence of an unconnected wetland increased WRT by 0.20 standardized units (90% BCI: -0.20 ; 0.59 ; $P(\beta > 0) = 0.80$) and a connected wetland by 0.01 (90% BCI: -0.35 ; 0.39). The WRT $\rightarrow \Delta\text{DOC}$ path was positive but weak (0.09 ; 90% BCI: 0.01 ; 0.16). The resulting total effects of connectivity on ΔDOC were negligible: 0.02 (90% BCI: -0.02 ; 0.07) for unconnected and 0.00 (90% BCI: -0.04 ; 0.04) for connected wetlands (Figure S7). We therefore found no evidence that lateral stream–wetland connectivity modifies the beaver-mediated DOC change through the water-residence-time pathway. Given that only 7% and 6% of sites fell into the connected and unconnected wetland categories, we regard this result as power-limited rather than as strong evidence of absence of an effect. Importantly, the estimates of the remaining paths were consistent with those of the main SCM (Figure 2b, main manuscript), indicating that our main conclusions are robust to this alternative causal structure.




Figure S6: Extended directed acyclic graph (DAG) including lateral stream–wetland connectivity (exposure, green) as a driver of water residence time (WRT). Green edges mark the open front-door paths from connectivity to ΔDOC (outcome, blue).




Figure S7: Posterior effects of lateral stream–wetland connectivity (relative to sites without wetland) on water residence time (left) and the total mediated effect on ΔDOC (right), controlling for beaver-dam variables, channel gradient and solar radiation. Points show posterior means; thick and thin bars show the 66% and 90% credible intervals. The dashed line marks zero.

References

- Bencala KE and Walters RA. 1983. Simulation of solute transport in a mountain pool-and-riffle stream: A transient storage model. *Water Resources Research* **19**: 718–24.
- Abbasi M, Peacock M, Drakare S, et al. 2024. Water residence time is an important predictor of dissolved organic matter composition and drinking water treatability. *Water Research* **260**: 121910.
- Aitkenhead JA, Hope D, and Billett MF. 1999. The relationship between dissolved organic carbon in stream water and soil organic carbon pools at different spatial scales. *Hydrological Processes* **13**: 1289–302.
- Arif S and MacNeil MA. 2023. Applying the structural causal model framework for observational causal inference in ecology. *Ecological Monographs* **93**: e1554.
- Cincotta MM, Perdrial JN, Shavitz A, et al. 2019. Soil aggregates as a source of dissolved organic carbon to streams: An experimental study on the effect of solution chemistry on water extractable carbon. *Frontiers in Environmental Science* **7**.
- Clubb, F., Mudd, S., Milodowski, D., Valters, D. A., Slater, L. J., Hurst, M. D., & Limaye, A. B. 2017. Geomorphometric delineation of floodplains and terraces from objectively defined topographic thresholds. *Earth Surface Dynamics*, 5(3), 369-385.
- Dittbrenner BJ, Schilling JW, Torgersen CE, and Lawler JJ. 2022. Relocated beaver can increase water storage and decrease stream temperature in headwater streams. *Ecosphere* **13**: e4168.
- Feuchtmayr H, Moran R, Hatton K, et al. 2009. Global warming and eutrophication: effects on water chemistry and autotrophic communities in experimental hypertrophic shallow lake mesocosms. *Journal of Applied Ecology* **46**: 713–23.
- Finger D, Schmid M, and Wüest A. 2007. Comparing effects of oligotrophication and upstream hydropower dams on plankton and productivity in perialpine lakes. *Water Resources Research* **43**.
- FOEN, 2021. Vegetation height model NFI.
- Gelman A, Carlin JB, Stern HS, et al. 2013. *Bayesian Data Analysis*. CRC Press.
- Gelman A and Rubin DB. 1992. Inference from iterative simulation using multiple sequences. *Statistical Science* **7**: 457–72.
- Gessner MO, Chauvet E, and Dobson M. 1999. A perspective on leaf litter breakdown in streams. *Oikos* **85**: 377–84.
- Hijmans, R. (2025). terra: Spatial Data Analysis. R package version 1.8-87.
- Jepsen SM, Harmon TC, Sadro S, et al. 2019. Water residence time (age) and flow path exert synchronous effects on annual characteristics of dissolved organic carbon in terrestrial runoff. *Science of The Total Environment* **656**: 1223–37.
- Johnston CA. 2014. Beaver pond effects on carbon storage in soils. *Geoderma* **213**: 371–8.
- Larson JH, Frost PC, Zheng Z, et al. 2007. Effects of upstream lakes on dissolved organic matter in streams. *Limnology and Oceanography* **52**: 60–9.
- Lindsay, J. B. 2016. Whitebox GAT: A case study in geomorphometric analysis. *Computers & Geosciences*, 95, 75-84.

- Majerova M, Neilson BT, Schmadel NM, et al. 2015. Impacts of beaver dams on hydrologic and temperature regimes in a mountain stream. *Hydrology and Earth System Sciences* **19**: 3541–56.
- Manolaki P, Olesen A, Hvidt BG, et al. 2021. Investigating emergent macrophytes establishment rate and propagation towards constructed wetlands efficacy optimization. *Knowledge & Management of Aquatic Ecosystems* 23.
- McGuire KJ, McDonnell JJ, Weiler M, et al. 2005. The role of topography on catchment-scale water residence time. *Water Resources Research* **41**.
- Nyssen J, Pontzele J, and Billi P. 2011. Effect of beaver dams on the hydrology of small mountain streams: Example from the chevral in the ourthe orientale basin, ardennes, belgium. *Journal of Hydrology* **402**: 92–102.
- Reitsema RE, Meire P, and Schoelynck J. 2018. The future of freshwater macrophytes in a changing world: Dissolved organic carbon quantity and quality and its interactions with macrophytes. *Frontiers in Plant Science* **9**.
- Reynolds CS. 2006. The ecology of phytoplankton. Cambridge: Cambridge University Press.
- Sell AF and Overbeck J. 1992. Exudates: Phytoplankton-bacterioplankton interactions in plusee. *Journal of Plankton Research* **14**: 1199–215.
- Søndergaard M, Riemann B, and Jørgensen NOG. 1985. Extracellular organic carbon (EOC) released by phytoplankton and bacterial production. *Oikos* **45**: 323–32.
- SwissTopo. 2022. swissALTI3D.
- Vats D and Knudson C. 2021. Revisiting the gelmanrubin diagnostic. *Statistical Science* **36**: 518–29.
- Viviroli D, Zappa M, Gurtz J, and Weingartner R. 2009. An introduction to the hydrological modelling system PREVAH and its pre- and post-processing-tools. *Environmental Modelling & Software* **24**: 1209–22.
- Westbrook CJ, Cooper DJ, and Baker BW. 2006. Beaver dams and overbank floods influence groundwatersurface water interactions of a Rocky Mountain riparian area. *Water Resources Research* **42**.
- Wohl E. 2013. Landscape-scale carbon storage associated with beaver dams. *Geophysical Research Letters* **40**: 3631–6.
- Wu, Q., Brown, A. 2022. 'whitebox': 'WhiteboxTools' R Frontend. R package, version 2.2.0.