SeeWandel Climate

Interreg
Alpenrhein-Bodensee-Hochrhein

Kofinanziert von der Europäischen Union

Regime Shifts in Plankton Dynamics: What Drives Them and Can We Predict Them?

Dr. Leonardo Capitani
Department of Aquatic Ecology, Eawag · SeeWandel-Climate

www.seewandel.org

Ecomeets at Eawag, Dübendorf
28 October 2026 · 11:00–12:00

Slides and links

Scan to download the slides

SORTEE
Society for Open, Reliable, and Transparent Ecology and Evolutionary Biology

Slides, code and data shared openly
leomarameo7.github.io/science_communication.html

The Pomati lab



Pomati lab
Plankton Ecology, Eawag

SeeWandel-Climate
Lake Constance

Take home message


Plankton regime shifts can be detected and partly explained by biotic drivers. Whether they propagate and can be predicted is still open.



Outline

1. What we know: background
2. Research questions
3. Methods: our pipeline
4. Results: detection of regime shifts
5. Explain regime shifts
6. Propagation of regime shifts
7. Conclusions

What we know: background




Lake Constance
473 km² · 252 m deep

Phosphate ↑ until 1980, then ↓

Temperature ↑ about 1 °C

Figure: Dietmar Straile, University of Konstanz

What we know: the plankton community



2005

Spotlight: what we expect to drive the omnivores (Om)

Network: Merz et al. 2023, Nat. Clim. Change · hypotheses: notebook 18 · Ci, Ro: not in the data

Research questions



1 · Detect
Abrupt shifts?



2 · Explain
What drives them?



3 · Propagate
Does a shift spread?



4 · Predict
Can we forecast them? (in progress)

Methods: our pipeline

1

Data

Monthly data 1975–2022

2

Detect

BEAST + two criteria

3

Explain

GP-EDM + hypotheses

4

Propagate

Shift-removal test

5


Predict

Forward validation

WORK IN PROGRESS

Gilarranz et al. 2022 · Munch et al. 2017 · Merz et al. 2023

Results: detection of regime shifts






Shift: Aug 2014

abruptness 1.19

Omnivores, monthly biovolume. Dashed: shift; red: period means.

The omnivores dropped abruptly in 2014

Explain regime shifts






LOO R^2 = **0.81**

Omnivores ↓ with cyanobacteria, small diatoms, large gold algae

Prey hypothesis: **not supported**

Heat maps: axes in log10(mm³ m⁻² + 1); dots: observed months

Explain regime shifts: partial dependence





x: predictor 1
lines: predictor 2

Cyanobacteria ↓ omnivores

Lines never cross

Other predictors at their median; axes in log10(mm³ m⁻² + 1); lines: 5th to 95th percentile of the second predictor

Propagation of regime shifts

1 · Real predictor



2 candidate pairs

0 testable

Predictor shifted **after**

2 · Shift-free surrogate (× 100)



Gap between real and surrogate prediction = evidence for propagation

No evidence yet: too few shifts

Conclusions



Shifts are detectable



Biotic drivers matter



Propagation, prediction: open

Take home message

Plankton regime shifts can be detected and partly explained by biotic drivers. Whether they propagate and can be predicted is still open.



Acknowledgments

Interreg
Alpenrhein-Bodensee-Hochrhein

Kofinanziert von der Europäischen Union

igkb
Internationale Gewässerschutzkommission für den Bodensee

IBKF


SeeWandel-Climate · Pomati lab and colleagues at Eawag
Luis J. Gilarranz · Stefanie Eyring · Stephan B. Munch · Tanya L. Rogers · Francesco Pomati
Data: IGKB, IBKF, University of Konstanz (D. Straile)

Supplementary material


Methods in detail



What is GP-EDM?



Gaussian process: no fixed equation




Next month = f(past states, drivers)

f learned from the data: Gaussian process


Takens 1981 · Munch et al. 2017

Automatic relevance determination (ARD)



One length scale per predictor: **phi near 0** = predictor dropped

Detecting regime shifts: BEAST and two criteria



Kept if **both** hold


2 interval excludes 0


3 abruptness > 1

abruptness = |difference| / mean SD of the two periods

Gilarranz et al. 2022

In-sample versus out-of-sample R^2





In-sample: $R^2 = 1$, a GP interpolates

LOO: one month left out

LFO: forecast the last 10 %