The Beaver as an Ecosystem Engineer

Effects on Water Quality and Freshwater Ecosystems

Dr. Leonardo Capitani

Senior Scientist, Department of Aquatic Ecology, Eawag

leomarameo7.github.io

23 October 2026

Slides and links

Scan to download the slides



open science

Acknowledgments

Christof Angst

Leiter Biberfachstelle

Dr. Francesco Pomati

Dr. Anita C. Risch

Dr. Annegret Larsen

Dr. Joshua R. Larsen

… and many other researchers and stakeholders!

Main message of this talk

EXCAVATION, CANAL DIGGING



+ Wetland area
+ Lateral connectivity
+ Aquatic–terrestrial interface

DAM BUILDING

+ Water storage
+ Groundwater connection
– Flow velocity
+ Sedimentation rate

FORAGING, TREE CUTTING

+ Plant diversity
– Canopy cover
+ Landscape heterogeneity

> Beaver engineering can create locally climate-resistant wetlands

Index of this presentation

1. **Water quality:** carbon and nutrients

2. **Biodiversity:** habitats and species

3. **Resilience:** floods and droughts

4. **Management:** living with beavers

> International synthesis + Swiss 🇨🇭 cases

Beavers are back across Europe

2002



2020



Switzerland, 2022: 4,900 beavers

info fauna

Beavers are recolonising Europe

How many beavers?



info fauna

- 150 beavers in 2008
- 400+ beavers in 2022
- Growth slows after 2020

Rebound now levelling off

Living with beavers: the cost of damage



Damage compensation increased over 30 years (1988–2021)

Hydrological impacts



Incised channel, dams blow out

Wider channel, stable dams



Sediment builds, water table rises



Floodplain reconnected

Beavers reduce longitudinal connectivity, while increasing lateral connectivity

Water quality · Why nutrient cycling?

1 Inorganic nutrients

Nitrate NO_3^-

Ammonium NH_4^+

Phosphate PO_4^{3-}

(N and P, essential nutrients)



2

DOM

dissolved organic matter

DOC

dissolved organic carbon

= carbon part of DOM

DIC

dissolved inorganic carbon

Nutrients and organic matter drive ecosystems

Water quality · International synthesis: C and nutrients

❶ Upstream of dam

- More open water
- Longer residence time
- Less oxygen

❷ Storage

- C carbon
- N nitrogen
- P phosphorus
- sediment



❸ To the atmosphere

- N_2 nitrogen gas
- CO_2 carbon dioxide
- CH_4 methane

❹ Export downstream

- DOC dissolved organic C ↑
- NH_4^+ ammonium ↑
- NO_3^- nitrate ↓
- PO_4^{3-} phosphate ↕

❺ Downstream of dam

- Less open water
- Oxygen returns

Carbon and nutrient budgets: no unique direction

Water quality: the Swiss case



Beaver ponds are local control points of DOC

Water quality · drivers of beaver's influence



Beavers slow the water; algae and plants add DOC

Biodiversity: the Swiss case





Beavers influence stream biodiversity

Biodiversity · Bats at beaver pools



Increased canopy heterogeneity

Increased standing deadwood density

Increased arthropod abundance

Echolocation guild	*Species richness*	*Activity*	*Feeding*
Short-range	n.s.	n.s.	n.s.
Mid-range	+	+	+
Long-range	+	+	*no data*

\+ significant increase at beaver pools n.s. not significant

- Mid-range gain most
- Driver: deadwood
- Endangered bats:

Western barbastelle
Barbastella barbastellus

Geoffroy's bat
Myotis emarginatus

Beaver contribution: habitat complexity and more food

Resilience: the Belgian case



Study site: Chevral, Ardennes, Belgium

Flatter peaks, higher low flows

Resilience · Storing water, buffering flow

243 m³

surface water stored per 100 m of stream (first year)

Dittbrenner et al. 2022

+60%

more water held in dry summers (incl. water table)

Hood 2012

–2.3 °C

cooler summer base flow downstream of dams (mean; varies by site)

Dittbrenner et al. 2022

–30%

peak flow in storms below beaver ponds, vs. water entering

Puttock et al. 2017

Dams buffer droughts and floods

Beaver management: conflicts and solutions

Conflict	Solutions
Foraging crop and ornamental tree loss	**fence or wrap trees in wire; plant species beavers avoid**
Burrowing bank erosion, undermined infrastructure	**leave a vegetated strip along the bank; strengthen the bank with stone or mesh**
Damming flooded crops, trees, roads	**pipe through the dam to keep the water low; cut a gap in the dam; wider pipes under roads**

Each conflict has a practical fix

Beaver management: zoning in the Czech Republic

Zone A

Full protection

links key conservation sites (Natura 2000)

smallest area

Zone B

Managed, with limits

dam management; removal if non-lethal methods fail

85.5%

Zone C

Not tolerated

farmed-fish ponds and infrastructure at high risk of beaver damage

13.3%

> Zoning separates protection from conflict areas

Beaver management: the Swiss case



info fauna Biber-Auenmodell

Beaver floodplain: 450 km²

- Driver: channel gradient
- **298 km²** (66%) opportunity
- **132 km²** (29%) conflict

Maximise restoration, minimise conflicts

Beaver management: how should we proceed?



Beaver management should consider social context

Main message: take-home

EXCAVATION, CANAL DIGGING



+ Wetland area
+ Lateral connectivity
+ Aquatic–terrestrial interface

DAM BUILDING

+ Water storage
+ Groundwater connection
– Flow velocity
+ Sedimentation rate

FORAGING, TREE CUTTING

+ Plant diversity
– Canopy cover
+ Landscape heterogeneity

> Beaver engineering can create locally climate-resistant wetlands

Thank you

Photo: Dominic Tinner

Dr. Leonardo Capitani
leomarameo7.github.io

Supplementary material

Additional slides

Beaver management: a ladder of options



> Start with information; remove beavers last

Beaver life history and diet

Eurasian beaver eating berry leaves

- **Life history**
- **Diet**

17.8 kg average adult body mass	**105–107 d** gestation (about 3.5 months)	**2–3** baby beavers (kits) per birth, once a year
~20 mo age at sexual maturity	**3.8** animals per family (parents + offspring)	**9 yr** average age at death (up to 20–24 yr)
~2 kg woody plants eaten per day (captive)	**86 → 17%** woody share of diet, winter → August	**~1,200 kcal** energy needed per day (15-kg beaver)

Slow breeders: few kits, long family life

Sampling design and sites





Connectivity and expansion



Isolated northern populations expand slowly

Fig. 3 in Falaschi et al. 2024, Animal Conservation 27, 324–337: connectivity map of Eurasian beaver in Italy and surrounding countries. Source points

Beavers dam streams in human-used landscapes

Hersiwil
Canton Solothurn (SO)

Beaver dam

Beaver lodge

Dams and lodge sit next to a field

Beavers dam streams right next to farmland

Beaver dams beside paths and railways

Herzogenbuchsee
Canton Bern (BE)

Beaver dam
Beaver lodge

Territory between a footpath and a railway

Beaver territories can sit next to infrastructure

References (1/2)

- Angst, C., Auberson, C. & Nienhuis, C. (2023). Biberbestandeserhebung 2022 in der Schweiz und Liechtenstein.
- Brazier, R.E., Puttock, A., Graham, H.A., Auster, R.E., Davies, K.H. & Brown, C.M.L. (2021). Beaver: Nature's ecosystem engineers. *Wires. Water*, 8, e1494.
- Browning, E. (2025). Direct and indirect impacts of beaver ecosystem engineering have mixed effects on bats across feeding guilds. *Journal of Animal Ecology*.
- Dennis, M., Angst, C., Larsen, J.R., Rey, E. & Larsen, A. (2024). A national scale floodplain model revealing channel gradient as a key determinant of beaver dam occurrence and inundation potential can anticipate land-use based opportunities and conflicts for river restoration. *Global Ecology and Conservation*, 56, e03304.
- Dittbrenner, B.J., Schilling, J.W., Torgersen, C.E. & Lawler, J.J. (2022). Relocated beaver can increase water storage and decrease stream temperature in headwater streams. *Ecosphere*, 13, e4168.
- Fairfax, E. & Westbrook, C. (2024). The Ecology and Evolution of Beavers: Ecosystem Engineers that Ameliorate Climate Change. *Annual Review of Ecology, Evolution, and Systematics*, 55, 323–345.
- Falaschi, M., Ficetola, G.F., Viviano, A., Mazza, G. & Mori, E. (2024). Environmental suitability and potential range expansion of the Eurasian beaver in Italy. *Animal Conservation*, 27, 324–337.
- Halley, D.J., Saveljev, A.P. & Rosell, F. (2020). Population and distribution of beavers Castor fiber and Castor canadensis in Eurasia. *Mammal Review*, 51, 1–24.

References (2/2)

- Jansen, J., MacIntyre, S., Barrett, D.C., Chin, Y.-P., Cortés, A., Forrest, A.L., *et al.* (2021). Winter limnology: How do hydrodynamics and biogeochemistry shape ecosystems under ice? *Journal of Geophysical Research: Biogeosciences*, 126, e2020JG006237.
- Moser, V., Capitani, L., Zehnder, L., Hürbin, A., Obrist, M.K., Ecker, K., *et al.* (2025). Habitat heterogeneity and food availability in beaver-engineered streams foster bat richness, activity and feeding. *Journal of Animal Ecology*.
- Moser, V., Minnig, S., Capitani, L., Boch, S., Cramer, N., Edman, O., *et al.* (2026). Rewilding beyond the wilderness: Beavers can restore stream biodiversity from urban to agricultural to natural landscapes. *Journal of Applied Ecology*.
- Nationale Biberfachstelle / info fauna. (2023). Beaver counts in the canton of Zürich, 2008–2022.
- Nyssen, J., Pontzeele, J. & Billi, P. (2011). Effect of beaver dams on the hydrology of small mountain streams: Example from the Chevral in the Ourthe Orientale basin, Ardennes, Belgium. *Journal of Hydrology*, 402, 92–102.
- Puttock, A., Graham, H.A., Cunliffe, A.M., Elliott, M. & Brazier, R.E. (2017). Eurasian beaver activity increases water storage, attenuates flow and mitigates diffuse pollution from intensively-managed grasslands. *Science of The Total Environment*, 576, 430–443.
- Rosell, F. & Campbell-Palmer, R. (2022). *Beavers: Ecology, Behaviour, Conservation, and Management*. Oxford University Press.
- Sheppy, J., Sudduth, E.B., Clinton, S., Riveros-Iregui, D. & Ledford, S.H. (2024). Urban beaver ponds show limited impact on stream carbon quantity in contrast to stormwater ponds. *Urban Ecosystems*, 27, 1477–1491.