

Session 2: Linear transformations and matrices

Optimization and Computational Linear Algebra for Data Science

Contents

1. Recap of the videos
2. Operation on matrices
3. Kernel and Image
4. Why do we care about all these things ?

Solving linear systems

Questions?


Linear maps & matrices

Two sides of the same coin

Linear map

function transformation

$$L : \mathbb{R}^m \rightarrow \mathbb{R}^n$$



$$\begin{cases} L(u+v) = L(u) + L(v) \\ L(\alpha u) = \alpha L(u) \end{cases}$$

$$L(\alpha u) = \alpha L(u)$$

for all $\alpha \in \mathbb{R}$, $u, v \in \mathbb{R}^m$

Matrix

$$\tilde{L} \in \mathbb{R}^{n \times m}$$

$$\tilde{L} = \begin{pmatrix} l_{1,1} & \dots & l_{1,m} \\ \vdots & & \vdots \\ l_{n,1} & \dots & l_{n,m} \end{pmatrix}$$

The matrix \tilde{L} is shown with dimensions m (width) and n (height) indicated by red arrows. The entries are labeled $l_{i,j}$ in blue.

Two sides of the same coin

Linear map

$$L: \mathbb{R}^m \rightarrow \mathbb{R}^n$$

$$L(x) = \tilde{L}x$$

for all $x \in \mathbb{R}^m$

matrix vector product

Matrix

$$\tilde{L} \in \mathbb{R}^{n \times m}$$

$$\tilde{L} = \begin{pmatrix} | & & | \\ L(e_1) & \dots & L(e_m) \\ | & & | \end{pmatrix}$$

n

$$L: \mathbb{R}^m \rightarrow \mathbb{R}^n$$


$$x \mapsto \tilde{L}x$$

L is linear

$$\tilde{L} = \begin{pmatrix} l_{1,1} & \dots & l_{1,m} \\ \vdots & & \vdots \\ l_{n,1} & \dots & l_{n,m} \end{pmatrix}$$

$$\rightarrow \begin{pmatrix} | & & | \\ L(e_1) & \dots & L(e_m) \\ | & & | \end{pmatrix}$$



$$L(e_1) = \tilde{L} e_1 =$$



Rotations in \mathbb{R}^2

Let $\theta \in \mathbb{R}$. The rotation $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ of angle θ about the origin is linear.

Exercise: what is the canonical matrix of R_θ ?



$$R_\theta = \begin{pmatrix} | & | \\ R_\theta(e_1) & R_\theta(e_2) \\ | & | \end{pmatrix}$$

$$R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

$$R_\theta \left(\begin{pmatrix} x \\ y \end{pmatrix} \right) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix}$$

$$v = \begin{pmatrix} x \\ y \end{pmatrix} \quad \underline{\underline{R_\theta(v)}}$$

Operations on matrices

Addition and scalar multiplication

- Sum of two matrices of the **same** dimensions:

$$\begin{pmatrix} a_{1,1} & \cdots & a_{1,m} \\ \vdots & \ddots & \vdots \\ a_{n,1} & \cdots & a_{n,m} \end{pmatrix} + \begin{pmatrix} b_{1,1} & \cdots & b_{1,m} \\ \vdots & \ddots & \vdots \\ b_{n,1} & \cdots & b_{n,m} \end{pmatrix} = \begin{pmatrix} a_{1,1} + b_{1,1} & \cdots & a_{1,m} + b_{1,m} \\ \vdots & \ddots & \vdots \\ a_{n,1} + b_{n,1} & \cdots & a_{n,m} + b_{n,m} \end{pmatrix}$$

- Multiplication by a scalar λ :

$$\lambda \begin{pmatrix} a_{1,1} & \cdots & a_{1,m} \\ \vdots & \ddots & \vdots \\ a_{n,1} & \cdots & a_{n,m} \end{pmatrix} = \begin{pmatrix} \lambda a_{1,1} & \cdots & \lambda a_{1,m} \\ \vdots & \ddots & \vdots \\ \lambda a_{n,1} & \cdots & \lambda a_{n,m} \end{pmatrix}$$

A new vector space!

Proposition

❑ $\mathbb{R}^{n \times m}$ is a vector space.

❑ $\dim(\mathbb{R}^{n \times m}) = n \cdot m$

Proof. *sketch*

→ We can verify that the matrices


$$\underline{E_{i,j}} = \begin{pmatrix} 0 & \cdots & \overset{\downarrow j}{1} & \cdots & 0 \\ \vdots & & \uparrow i & & \vdots \\ 0 & \cdots & 0 & \cdots & 0 \end{pmatrix} \leftarrow i, \text{ for } \begin{matrix} 1 \leq i \leq n \\ 1 \leq j \leq m \end{matrix}$$

form a basis of $\mathbb{R}^{n \times m}$, with $n \cdot m$ elements.

□

Product of two matrices

Warning:

$$\begin{pmatrix} a_{1,1} & \cdots & a_{1,m} \\ \vdots & \ddots & \vdots \\ a_{n,1} & \cdots & a_{n,m} \end{pmatrix} \times \begin{pmatrix} b_{1,1} & \cdots & b_{1,m} \\ \vdots & \ddots & \vdots \\ b_{n,1} & \cdots & b_{n,m} \end{pmatrix} \neq \begin{pmatrix} a_{1,1} \times b_{1,1} & \cdots & a_{1,m} \times b_{1,m} \\ \vdots & \ddots & \vdots \\ a_{n,1} \times b_{n,1} & \cdots & a_{n,m} \times b_{n,m} \end{pmatrix}$$



We will never consider such notion of product
because it doesn't have any "geometrical
meaning"

Matrix product

Let $\underline{L} \in \mathbb{R}^{n \times m}$ and $\underline{M} \in \mathbb{R}^{m \times k}$.

Definition (Matrix product)

The matrix product \underline{LM} is the $n \times k$ matrix of the linear map $L \circ M$.



The composition $L \circ M$ is linear.

Matrix product


Theorem

Let $L \in \mathbb{R}^{n \times m}$ and $M \in \mathbb{R}^{m \times k}$.

The entries matrix product LM are given by

$$(LM)_{i,j} = \sum_{\ell=1}^m L_{i,\ell} M_{\ell,j},$$

for $1 \leq i \leq n$ and $1 \leq j \leq k$.



Proof

$$LM \stackrel{\text{def}}{=} \begin{pmatrix} | & & | \\ \text{LoM}(e_1) & \dots & \text{LoM}(e_n) \\ | & & | \end{pmatrix}$$

$$(LM)_{i,j} = (\text{LoM}(e_j))_i \leftarrow i^{\text{th}} \text{ coefficient of } \text{LoM}(e_j)$$

$$= \left(L(\underline{M}(e_j)) \right)_i$$

$$M(e_j) = \begin{pmatrix} M_{1,j} \\ \vdots \\ M_{m,j} \end{pmatrix}$$

$$= (L(v))_i$$

$$= (Lv)_i$$

(matrix-vector product)

$$= \sum_{e=1}^m L_{i,e} v_e =$$

$$\sum_{e=1}^m L_{i,e} M_{e,j}$$

Rotations in \mathbb{R}^2

The R_a and R_b denote respectively the matrices of the rotations of angles a and b about the origin, in \mathbb{R}^2 .

Exercise: Compute the product $R_a R_b$.

$$R_a = \begin{pmatrix} \cos a & -\sin a \\ \sin a & \cos a \end{pmatrix}$$

1st method

$$R_a R_b = \begin{pmatrix} \cos a & -\sin a \\ \sin a & \cos a \end{pmatrix} \cdot \begin{pmatrix} \cos b & -\sin b \\ \sin b & \cos b \end{pmatrix} = \begin{pmatrix} \cos a \cos b - \sin a \sin b & * \\ \sin b \sin a + \cos a \sin b & * \end{pmatrix}$$

2nd method : $R_a R_b$ is the composition of R_b and R_a which is a rotation of angle $a+b$:

$$R_a R_b = R_{a+b} = \begin{pmatrix} \cos(a+b) & -\sin(a+b) \\ \sin(a+b) & \cos(a+b) \end{pmatrix}$$

(As a by product: $\cos(a+b) = \cos a \cos b - \sin a \sin b$)

Matrix product properties


• Let $A, B \in \mathbb{R}^{n \times m}$ and $C, D \in \mathbb{R}^{m \times k}$

$$(A+B) \cdot C = A \cdot C + B \cdot C$$

$$A(C+D) = AC + AD$$

Recall $\text{Id}_n = \begin{pmatrix} 1 & & (0) \\ & \ddots & \\ (0) & & 1 \end{pmatrix} : \begin{cases} \text{Id}_n A = A \\ A \cdot \text{Id}_m = A \end{cases}$

Issue : $\underline{AC} \neq \textcircled{CA}$ if $\underline{k \neq n}$



Can we divide two matrices ?

For instance if ~~A~~ $B = \cancel{A} C$
do we have $B = C$?

Take : $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ $B = \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}$ $C = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$

• $AC = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$

• $AB = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$

$AB = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = AC$ but $C \neq B$.

Invertible matrices

Definition (Matrix inverse)

A **square** matrix $M \in \mathbb{R}^{n \times n}$ is called *invertible* if there exists a matrix $M^{-1} \in \mathbb{R}^{n \times n}$ such that

$$MM^{-1} = M^{-1}M = \text{Id}_n.$$

Such matrix M^{-1} is unique and is called the *inverse* of M .

Exercise: Let $A, B \in \mathbb{R}^{n \times n}$. Show that if $AB = \text{Id}_n$ then $BA = \text{Id}_n$.

If $AB = AC$ and A is invertible
then $\underbrace{A^{-1}A}_\text{Id}_n B = \underbrace{A^{-1}A}_\text{Id}_n C$ hence $\underline{B = C}$

Kernel and image

Definitions

Let $L : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear transformation.

$$0^n = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

notation

Definition (Kernel)

The kernel $\text{Ker}(L)$ (or nullspace) of L is defined as the set of all vectors $\underline{v \in \mathbb{R}^m}$ such that $L(v) = 0$, i.e.


$$\underline{\text{Ker}(L)} \stackrel{\text{def}}{=} \{v \in \mathbb{R}^m \mid L(v) = 0\}.$$

$\in \mathbb{R}^n$ ← zero vector of \mathbb{R}^n

Definition (Image)

The image $\text{Im}(L)$ (or column space) of L is defined as the set of all vectors $u \in \mathbb{R}^n$ such that there exists $v \in \mathbb{R}^m$ such that $L(v) = u$.

Picture



Remarks

Let $L : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear transformation.

Proposition

❖ $\text{Ker}(L)$ is a subspace of \mathbb{R}^m .

❖ $\text{Im}(L)$ is a subspace of \mathbb{R}^n .

proved in HW2.


Remark: $\text{Im}(L)$ is also the Span of the columns of the matrix representation of L .

$$L = \begin{pmatrix} | & & | \\ c_1 & \dots & c_m \\ | & & | \end{pmatrix} \begin{matrix} \begin{pmatrix} x_1 \\ \vdots \\ x_m \end{pmatrix} \end{matrix} = x_1 c_1 + \dots + x_m c_m$$

$$\text{Im}(L) = \text{Span}(c_1, \dots, c_m)$$

Example: orthogonal projection

Consider $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ to be the orthogonal projection onto the x -axis.



Why do we care about this ?

Linear systems

Assume that we given a dataset:

$$a_i = (a_{i,1}, \dots, a_{i,m}) \in \mathbb{R}^m, \quad y_i \in \mathbb{R} \quad \text{for } i = 1, \dots, n.$$

We would like to find $x \in \mathbb{R}^m$ such that


$$x_1 a_{i,1} + \dots + x_m a_{i,m} = y_i \quad \text{for all } i \in \{1, \dots, n\}.$$

We need to solve :

$$\begin{cases} x_1 a_{1,1} + \dots + x_m a_{1,m} = y_1 \\ \vdots \\ x_1 a_{n,1} + \dots + x_m a_{n,m} = y_n \end{cases}$$

Matrix notation

Let's write

$$A = \begin{pmatrix} a_{1,1} & \cdots & a_{1,m} \\ \vdots & \ddots & \vdots \\ a_{n,1} & \cdots & a_{n,m} \end{pmatrix} \in \mathbb{R}^{n \times m} \quad \text{and} \quad y = \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} \in \mathbb{R}^n.$$


Then the system of equations becomes $\boxed{Ax = y}$

- Case 1: $y \notin \text{Im}(A)$: my system does not have any solution.
- Case 2: $y \in \text{Im}(A)$, then there exists $x_0 \in \mathbb{R}^m$ such that $Ax_0 = y$.

Let's find all solutions !

Is x_0 the only solution to $Ax = y$?

x is solution

$$\Leftrightarrow Ax = y$$

$$Ax_0 = y$$

$$\Leftrightarrow Ax - Ax_0 = y - y$$

$$\Leftrightarrow A(x - x_0) = 0.$$

$$\Leftrightarrow \underline{x - x_0} \in \underline{\text{Ker}(A)}$$

$$\Leftrightarrow \text{there exists } v \in \text{Ker}(A)$$

The set of all solutions is such that : $x = x_0 + v$.

$$S = \{ x_0 + v \mid v \in \text{Ker}(A) \}$$

Conclusion: 3 possible cases

1. $y \notin \text{Im}(A)$: there is no solution to $Ax = y$.
2. $y \in \text{Im}(A)$, then there exists $x_0 \in \mathbb{R}^m$ such that $Ax_0 = y$. The set of solutions is then

$$S = \{x_0 + v \mid v \in \text{Ker}(A)\}.$$

- ❖ If $\text{Ker}(A) = \{0\}$, then $S = \{x_0\}$: x_0 is the unique solution.
- ❖ If $\text{Ker}(A) \neq \{0\}$, then $\text{Ker}(A)$ contains infinitely many vectors: there are infinitely many solutions.

Gaussian elimination

$$A = \begin{pmatrix} 1 & -1 & 0 & 1 \\ 2 & 0 & 1 & -1 \\ -1 & 5 & 2 & 0 \end{pmatrix} \in \mathbb{R}^{n \times m}$$

$$\text{and } y = \begin{pmatrix} 1 \\ 1 \\ 4 \end{pmatrix} \in \mathbb{R}^n.$$

$$\begin{cases} x_1 - x_2 + x_4 = 1 & (R_1) \\ 2x_1 + x_3 - x_4 = 1 & (R_2) \\ -x_1 + 5x_2 + 2x_3 = 4 & (R_3) \end{cases}$$

$$\begin{cases} x_1 - x_2 + x_4 = 1 & (R_1) \\ 0 + 2x_2 + x_3 - 3x_4 = -1 & (R_2) - 2(R_1) \\ 0 + 4x_2 + 2x_3 + x_4 = 5 & (R_3) + (R_1) \end{cases}$$

Handwritten notes: "leading coefficient" with arrows pointing to the circled x_1 in the first equation and the circled $2x_2$ in the second equation.

Gaussian elimination

$$\begin{cases} x_1 - x_2 + x_4 = 1 & (R_1) \\ 2x_2 + x_3 - 3x_4 = -1 & (R_2) \\ 0 + 0 + 7x_4 = 7 & (R_3) - 2(R_2) \end{cases}$$

$$\begin{cases} x_1 - x_2 + x_4 = 1 \\ 2x_2 - 3x_4 = -1 - x_3 \\ x_4 = 1 \end{cases}$$

x_3 will play a role of "parameter" of the set of solutions

$$\begin{cases} x_1 = 1 - \frac{x_3}{2} \\ x_2 = 1 - \frac{x_3}{2} \\ x_3 = x_3 \\ x_4 = 1 \end{cases}$$

Gaussian elimination


The set of solution

$$S = \left\{ \begin{pmatrix} 1 - x_3/2 \\ 1 - x_3/2 \\ x_3 \\ 1 \end{pmatrix} \mid \underline{x_3 \in \mathbb{R}} \right\}$$

$$= \left\{ \underbrace{\begin{pmatrix} 1 \\ 1 \\ 0 \\ 1 \end{pmatrix}}_{x_0} + v \mid v \in \text{Span} \left(\underbrace{\begin{pmatrix} -1/2 \\ -1/2 \\ 1 \\ 0 \end{pmatrix}}_{\text{Ker}(A)} \right) \right\}$$

Gaussian elimination

Questions?



Questions?

Questions?