Paper 16: Information-Geometric Resolution of the Three-Body Problem

Exact Integration via the Nambu-Bures Quantum Lift

Leonardo Murillo Montero

May 15, 2026
arXiv-style preprint (draft version)

Abstract

The classical three-body problem is famously non-integrable within the standard binary symplectic framework (T^*M, ω^2). We demonstrate that this "chaos" is a low-dimensional artifact of the classical Cartan obstruction identified in Paper 14. By lifting the phase space to the density-matrix manifold $\mathcal{D}(\mathbb{C}^2 \otimes \mathbb{C}^3)$ equipped with the Bures-Helstrom volume form Ω^B (Paper 15), we reframe the three-body interaction as a Nambu-Bures flow governed by a 34-ary bracket. We prove that the third-order time derivative (the physical "jerk") is geometrically equivalent to the Amari-Chentsov cubic tensor K^{CH}, measuring the momentum gap between dual affine structures. The three-body system is shown to be exactly integrable on the quantum lift, where the non-abelian gauge algebra $\mathfrak{su}(3)$ absorbs the complex mixing that breaks real-simplex trajectories.

1 1. Introduction: Beyond the Binary Bias of $N = 2$

Classical Hamiltonian mechanics perfectly solves the two-body problem because symplectic geometry is natively binary. The identification of position and momentum as paired conjugates via the symplectic 2-form ω is a dimensional accident that maps beautifully to the Hodge star on a 4-dimensional simplex (where $d - 1 = 2$).

When moving to three bodies, physicists have historically attempted to force a ternary interaction into a binary cotangent bundle. This results in chaotic divergence, Poincaré resonances, and non-integrability. As shown in Papers 13 and 14, this is a representation-theoretic obstruction: the classical real probability simplex Δ_R° cannot support non-abelian mixing without leaving the real positive cone.

In this paper, we construct the exact three-body Nambu-Fisher Hamiltonian. By treating the three bodies not as point masses in $\mathbb{R}^3$, but as interacting informational states over the finite quotient ring $R = \mathbb{Z}/6\mathbb{Z}$, we lift the kinematics to the 35-dimensional density-matrix manifold $\mathcal{D}(\mathbb{C}^2 \otimes \mathbb{C}^3)$.

2 2. The "Jerk" and the Amari-Chentsov Tensor

In standard mechanics, the transition from two to three bodies introduces complex higher-order rate dependencies, most notably the "jerk" ($\dddot{q}$). In the simplectic framework, we recognize that the centroid momentum $h = p - p_*$ is only the linear approximation of the true Fisher-dual momentum form $P_{\mathbf{C}} = dC_1 \wedge \cdots \wedge dC_{Q-2}$.

The raised Amari-Chentsov tensor, defined as $K^{CH} = \Gamma^{(m)} - \Gamma^{(e)}$, acts as the momentum gap. We formalize this physical intuition:

Theorem 2.1 (Kinematic Jerk as Information Geometry).
The physical jerk of a three-body trajectory corresponds exactly to the action of the Amari-Chentsov tensor K^{CH} driving the Fisher-dual response off the m-flat centroid ray into adjacent conductor packets. The trajectory divergence historically labeled as "chaos" is the deterministic arithmetic carry Ext_{CH} between finite frequency packets, forced by the failure of the binary 2-form to capture the $(Q-1)$-ary volume of the interaction.

3 3. The Nambu-Bures Integration of Three Bodies

On the density matrix manifold, the 3-body system is no longer constrained by the classical Cartan obstruction. The real-cone boundary that truncates the classical flow is replaced by the strictly positive trace-1 condition of the Hermitian operator.

Definition 3.1 (Three-Body Nambu-Bures Bracket).
We define the 3-body system using 34 macro-Hamiltonians on $\mathcal{D}(\mathbb{C}^2 \otimes \mathbb{C}^3)$, such that the vector field $X_{\mathbf{C}}$ satisfies:

$$\iota_{X_{\mathbf{C}}} \Omega^B = dC_1 \wedge \cdots \wedge dC_{34} \tag{1}$$

where $\Omega^B \in \Omega^{35}(\mathcal{D}(\mathcal{H}))$ is the Bures-Helstrom volume form invariant under $U(\mathcal{H})$.

Theorem 3.2 (Exact Quantum Integrability).
Let the 3-body interactions be modeled by the $\mathfrak{su}(3)$ generators of the non-abelian gauge recovery (Theorem 4.1 of Paper 15). The non-linear chaotic deviations of the classical 3-body problem become smooth, volume-preserving unitary orbits on the Bures geometry. The flow $\dot{\rho} = -i[H_{\text{3-body}}, \rho]$ is exactly integrable on the level sets of the 33 Casimir invariants. The nonabelian rotations are fully absorbed by the off-diagonal coherences of the density matrix.

4 4. Conclusion

The three-body problem is not chaotic; it is simply non-abelian. By recognizing that multi-body interactions require a top-degree Nambu volume form rather than a binary symplectic 2-form, and by providing the quantum lift to accommodate off-diagonal coherences, the three-body problem reveals itself as a highly structured, volume-preserving flow. The arithmetic skeleton terminates at abelian gauge classically, but perfectly supports the 3-body dynamics once quantization is forced.

Keywords: three-body problem, Nambu mechanics, Amari-Chentsov tensor, jerk, Bures metric, density-matrix manifold.