

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines™)

Antiemesis

Version 1.2012

NCCN.org

Continue



Comprehensive NCCN GuidelinesTM Version 1.2012 Panel Members Cancer Network® Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

David S. Ettinger, MD/Chair †
The Sidney Kimmel Comprehensive Cancer
Center at Johns Hopkins

Debra K. Armstrong, RN, OCN # Vanderbilt-Ingram Cancer Center

Sally Barbour, PharmD, BCOP, CCP Σ Duke Cancer Institute

Michael J. Berger, PharmD, BCOP ∑
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital and
Solove Research Institute

Philip J. Bierman, MD † ‡
UNMC Eppley Cancer Center at The Nebraska
Medical Center

Bob Bradbury, BCPS \sum H. Lee Moffitt Cancer Center & Research Institute

Georgianna Ellis, MD †
Fred Hutchinson Cancer Research Center/
Seattle Cancer Care Alliance

Steve Kirkegaard, PharmD Σ Huntsman Cancer Institute at the University of Utah

Dwight D. Kloth, PharmD, FCCP, BCOP \sum Fox Chase Cancer Center

Mark G. Kris, MD †
Memorial Sloan-Kettering Cancer Center

Dean Lim, MD †
City of Hope Comprehensive Cancer Center

Laura Boehnke Michaud, PharmD, BCOP ∑ The University of Texas M.D. Anderson Cancer Center

Lida Nabati, MD £ Þ
Dana-Farber/Brigham and Women's Cancer
Center | Massachusetts General Hospital
Cancer Center

Kim Noonan, RN, NP, AOCN #
Dana-Farber/Brigham and Women's Cancer
Center | Massachusetts General Hospital
Cancer Center

Hope S. Rugo, MD † ‡
UCSF Helen Diller Family Comprehensive
Cancer Center

Darby Siler, PharmD ∑
University of Alabama at
Birmingham Comprehensive Cancer
Center

Steven M. Sorscher, MD †
Siteman Cancer Center at BarnesJewish Hospital and Washington
University School of Medicine

Sundae Stelts, PharmD ∑ St. Jude Children's Hospital / University of Tennessee Cancer Institute

Lisa Stucky-Marshall, RN, MS, AOCN # Robert H. Lurie Cancer Center of Northwestern University

Barbara Todaro, PharmD ∑ Roswell Park Cancer Institute

Susan Urba, MD † £ University of Michigan Comprehensive Cancer Center

NCCN
Maoko Naganuma, MSc
Dorothy A. Shead, MS

Continue

Specialties Index ‡Hematology/hematology oncology ^bInternal medicine

†Medical oncology

*Nurse

 Σ Pharmacology

£Supportive care including palliative, pain management, pastoral care and oncology social work

Writing committee member

Comprehensive NCCN GuidelinesTM Version 1.2012 Table of Contents Cancer Network® Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

NCCN Antiemesis Panel Members

Guidelines Updates

Principles of Emesis Control for the Cancer Patient (AE-1)

CHEMOTHERAPY-INDUCED EMESIS:

High Emetic Risk Intravenous Chemotherapy -

Acute and Delayed Emesis Prevention (AE-2)

Moderate Emetic Risk Intravenous Chemotherapy - Emesis Prevention (AE-3)

Low and Minimal Emetic Risk Intravenous Chemotherapy - Emesis Prevention (AE-4)

Oral Chemotherapy - Emesis Prevention (AE-5)

Breakthrough Treatment for Chemotherapy-Induced Nausea/Vomiting (AE-6)

Emetogenic Potential of Intravenous Antineoplastic Agents (AE-7)

Emetogenic Potential of Oral Antineoplastic Agents (AE-9)

Principles of Managing Multiday Emetogenic Chemotherapy Regimens (AE-A)

Principles for Managing Breakthrough Emesis (AE-B)

RADIATION-INDUCED EMESIS:

Radiation-Induced Nausea/Vomiting (AE-10)

ANTICIPATORY EMESIS:

Anticipatory Emesis Prevention/Treatment (AE-11)

Clinical Trials: The NCCN believes that the best management for any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN member institutions, <u>click here:</u> <u>nccn.org/clinical_trials/physician.html.</u>

NCCN Categories of Evidence and Consensus: All recommendations are Category 2A unless otherwise specified.

See NCCN Categories of Evidence and Consensus

The NCCN Guidelines™ are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2011.

Comprehensive NCCN GuidelinesTM Version 1.2012 Updates Cancer Network[®] Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

Updates in Version 1.2012 of the NCCN Guidelines from Version 3.2011 include:

AE-2

• High emetic risk intravenous chemotherapy - acute and delayed emesis prevention, added preferred to palonosetron 0.25 mg IV day 1.

AE-3

- Moderate emetic risk intravenous chemotherapy emesis prevention, added preferred to palonosetron 0.25 mg IV day 1.
- Added footnote h: "Data with palonosetron is based on randomized studies with 2 drug combinations."
- Following granisetron 2 mg PO or 1 mg PO BID or 0.01 mg/kg (max 1 mg) IV day 1, added (category 1).

AE-5

• Oral chemotherapy - emesis prevention, removed dolasetron 100 mg PO daily from high to moderate emetic risk.

AE-6

- Breakthrough treatment for chemotherapy-induced nausea/vomiting, changed miscellaneous to "Other."
- Added olanzapine 2.5 5 mg PO BID (category 2B) and scopolamine 1 patch every 72 h to "Other" category.

AE-7, AE-8, AE-9

 Added footnote t: "Potential drug interactions between antineoplastic agents / antiemetic therapies and various other drugs should always be considered."

AE-8

- Low emetic risk, added eribulin.
- Minimal emetic risk, added ipilimumab and ofatumumab.

AE-9

• Minimal to low risk, added vandetanib.

AE-A

• Added "Alternatively, for highly emetogenic regimens, fosaprepitant 150 mg IV with recommended dexamethasone dosing may be given on day 1 with no need for oral aprepitant on days 2 and 3."

Note: All recommendations are category 2A unless otherwise indicated.

Comprehensive NCCN GuidelinesTM Version 1.2012 Cancer Network[®] Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

PRINCIPLES OF EMESIS CONTROL FOR THE CANCER PATIENT

- Prevention of nausea/vomiting is the goal.
- ➤ The risk of nausea/vomiting for persons receiving chemotherapy of high and moderate emetic risk lasts for at least 3 days for high and 2 days for moderate after the last dose of chemotherapy. Patients need to be protected throughout the full period of risk.
- Oral and intravenous 5-HT3 antagonists have equivalent efficacy when used at the appropriate doses.
- Consider the toxicity of the specific antiemetic(s).
- Choice of antiemetic(s) used should be based on the emetic risk of the therapy, prior experience with antiemetics, and patient factors.
- There are other potential causes of emesis in cancer patients.

These may include:

- > Partial or complete bowel obstruction
- Vestibular dysfunction
- ➤ Brain metastases
- > Electrolyte imbalance: hypercalcemia, hyperglycemia, or hyponatremia
- ▶ Uremia
- ➤ Concomitant drug treatments including opiates
- ➤ Gastroparesis: tumor or chemotherapy (eg, vincristine) induced or other causes (eg, diabetes)
- > Psychophysiologic:
 - * Anxiety
 - * Anticipatory nausea/vomiting
- For use of antiemetics for nausea/vomiting that are not related to radiation and/or chemotherapy, See NCCN Palliative Care Guidelines.
- For multi-drug regimens, select antiemetic therapy based on the drug with the highest emetic risk. See Emetogenic Potential of Intravenous Antineoplastic Agents (AE-7).
- Consider using an H2 blocker or proton pump inhibitor to prevent dyspepsia, which can mimic nausea.
- Lifestyle measures may help to alleviate nausea/vomiting, such as eating small frequent meals, choosing healthful foods, controlling the amount of food consumed, and eating food at room temperature. A dietary consult may also be useful. See NCI's "What You Should Know About Cancer Treatment, Eating Well, and Eating Problems." (http://www.cancer.gov/cancertopics/coping/eatinghints/page2#4)

Note: All recommendations are category 2A unless otherwise indicated.

Comprehensive NCCN Guidelines™ Version 1.2012 **Antiemesis**

NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

HIGH EMETIC RISK INTRAVENOUS CHEMOTHERAPY - ACUTE AND DELAYED EMESIS PREVENTION a,b,c

Start before chemotherapy c,d

- Serotonin (5-HT3) antagonist: e
- ➤ Dolasetron 100 mg PO f

➤ Granisetron 2 mg PO or 1 mg PO BID or 0.01 mg/kg (max 1 mg) IV day 1^f or transdermal patch as 3.1 mg/24 h patch (containing 34.3 mg granisetron total dose) applied approximately 24 - 48 h prior to first dose of chemotherapy; maximum duration of patch is 7 days

or

- ➤ Ondansetron 16 24 mg PO or 8 24 mg (max 32 mg/day) IV day 1^f
- ➤ Palonosetron 0.25 mg IV day 1 (preferred)

AND

- Steroid:^g
- > Dexamethasone 12 mg PO or IV day 1, 8 mg PO daily days 2 4 (with aprepitant 125 mg or fosaprepitant 115 mg IV day 1) or
- > Dexamethasone 12 mg PO or IV day 1, 8 mg PO day 2, then 8 mg PO BID days 3 and 4 (with fosaprepitant 150 mg IV day 1)

AND

- Neurokinin 1 antagonist:
- > Aprepitant 125 mg PO day 1, 80 mg PO daily days 2 3
- ➤ Fosaprepitant 150 mg IV day 1 only

- > Fosaprepitant 115 mg IV on day 1 only, then aprepitant 80 mg PO daily days 2 3
- ± Lorazepam 0.5 -2 mg PO or IV or sublingual either every 4 hours or every 6 hours days 1 4
- ± H2 blocker or proton pump inhibitor

^aData for post-cisplatin (≥50 mg/m²) emesis prevention are category 1; others are ^dSee Principles of Managing Multiday Emetogenic Chemotherapy Regimens (AE-A). category 2A.

^eOrder of listed antiemetics is alphabetical and does not reflect preference.

bSee Emetogenic Potential of Intravenous Antineoplastic Agents (AE-7).

^fSome NCCN Member institutions use a 5-HT3 antagonist on days 2 - 3.

risk as well as patient-specific risk factors.

^cAntiemetic regimens should be chosen based on the drug with the highest emetic ^gUse of steroids is contraindicated with drugs such as interleukin-2 (ie, IL-2, aldesleukin) and interferon.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

See Breakthrough **Treatment** (AE-6)

category 1 for combined regimens^c

Comprehensive NCCN Guidelines™ Version 1.2012 Cancer Network® Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

MODERATE EMETIC RISK INTRAVENOUS CHEMOTHERAPY - EMESIS PREVENTIONb,c,i

DAY 1

Start before chemotherapy c,d

- Serotonin (5-HT3) antagonist:e
- ➤ Dolasetron 100 mg PO (category 1) or
- ➤ Granisetron 2 mg PO or 1 mg PO BID or 0.01 mg/kg (max 1 mg) IV day 1 (category 1) or transdermal patch as 3.1 mg/24 h patch (containing 34.3 mg granisetron total dose) applied approximately 24 48 h prior to first dose of chemotherapy; maximum duration of patch is 7 days
- ➤ Ondansetron 16 24 mg PO or 8 12 mg (maximum 32 mg/day) IV (category 1) or
- ➤ Palonosetron 0.25 mg IV (category 1) day 1 only (preferred)^h

AND

• Steroid: 9

or

➤ Dexamethasone 12 mg PO or IV

WITH/WITHOUT

- Neurokinin 1 antagonist (for selected patients, where appropriate)
- ➤ Aprepitant 125 mg PO or
- ➤ Fosaprepitant 115 mg IV day 1 only
- ± Lorazepam 0.5 2 mg PO or IV or sublingual either every 4 or every 6 h PRN
- ± H2 blocker or proton pump inhibitor

bSee Emetogenic Potential of Intravenous Antineoplastic Agents (AE-7).

- ^cAntiemetic regimens should be chosen based on the drug with the highest emetic risk as well as patient-specific risk factors.
- dSee Principles of Managing Multiday Emetogenic Chemotherapy Regimens (AE-A).

DAYS 2 and 3

- Serotonin (5-HT3) antagonist monotherapy:e
- Dolasetron 100 mg PO daily or
- ➤ Granisetron 1 2 mg PO daily or 1 mg PO BID or 0.01 mg/kg (maximum 1 mg) IV
- ➤ Ondansetron 8 mg PO bid or 16 mg PO daily or 8 mg (maximum 32 mg/day) IV

OR

- Steroid monotherapy:9
- ➤ Dexamethasone 8 mg PO or IV daily

OR

- Neurokinin 1 antagonist ± steroid (if NK-1 antagonist used on day 1)^j
- ➤ Aprepitant 80 mg PO ± dexamethasone 8 mg PO or IV daily
- ± Lorazepam 0.5 2 mg PO or IV or sublingual either every 4 or every 6 h PRN
- ± H2 blocker or proton pump inhibitor

See Breakthrough
Treatment (AE-6)


- eOrder of listed antiemetics is alphabetical and does not reflect preference.
- ^gUse of steroids is contraindicated with drugs such as interleukin-2 (ie, IL-2, aldesleukin) and interferon.
- ^hData with palonosetron is based on randomized studies with 2 drug combinations.
- ⁱData for post-carboplatin ≥300 mg/m², cyclophosphamide ≥600 1000 mg/m², doxorubicin ≥50 mg/m² emesis prevention are category 1.
- ^jAs per high emetic risk prevention, aprepitant should be added (to dexamethasone and a 5-HT3 antagonist regimen) for select patients receiving other chemotherapies of moderate emetic risk (eg, carboplatin, cisplatin, doxorubicin, epirubicin, ifosfamide, irinotecan, or methotrexate) (See AE-2).

Note: All recommendations are category 2A unless otherwise indicated.

Comprehensive NCCN GuidelinesTM Version 1.2012 Cancer Network[®] Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

LOW AND MINIMAL EMETIC RISK INTRAVENOUS CHEMOTHERAPY - EMESIS PREVENTIONC, d, I



See Principles of Emesis Control for the Cancer Patient (AE-1)

Note: All recommendations are category 2A unless otherwise indicated.

^cAntiemetic regimens should be chosen based on the drug with the highest emetic risk as well as patient-specific risk factors.

dSee Principles of Managing Multiday Emetogenic Chemotherapy Regimens (AE-A).

⁹Use of steroids is contraindicated with drugs such as interleukin-2 (ie, IL-2, aldesleukin) and interferon.


^kMonitor for dystonic reactions; use diphenhydramine 25 - 50 mg PO or IV either every 4 or every 6 h for dystonic reactions. If allergic to diphenhydramine, use benztropine at 1 - 2 mg IV or IM x 1 dose, followed by oral dose of 1 - 2 mg daily or BID if needed to control the reaction.

See Emetogenic Potential of Intravenous Antineoplastic Agents (AE-8).

Comprehensive NCCN Guidelines™ Version 1.2012 Cancer Network® Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

ORAL CHEMOTHERAPY - EMESIS PREVENTION C,d,m,n



^cAntiemetic regimens should be chosen based on the drug with the highest emetic risk as well as patient-specific risk factors.

Note: All recommendations are category 2A unless otherwise indicated.

d See Principles of Managing Multiday Emetogenic Chemotherapy Regimens (AE-A).

^eOrder of listed antiemetics is alphabetical and does not reflect preference.

^kMonitor for dystonic reactions; use diphenhydramine 25 - 50 mg PO or IV either every 4 or every 6 h for dystonic reactions. If allergic to diphenhydramine, use benztropine at 1 - 2 mg IV or IM x 1 dose, followed by oral dose of 1 - 2 mg daily or BID if needed to control the reaction.

^mSee Emetogenic Potential of Oral Antineoplastic Agents (AE-9).

ⁿThese antiemetic recommendations apply to oral chemotherapy only. When combined with IV agents in a combination chemotherapy regimen, the antiemetic recommendations for the agent with the highest level of emetogenicity should be followed. If multiple oral agents are combined, emetic risk may increase and require prophylaxis.

Comprehensive NCCN GuidelinesTM Version 1.2012 Cancer Network[®] Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

BREAKTHROUGH TREATMENT FOR CHEMOTHERAPY-INDUCED NAUSEA/VOMITING^{d,o}

RESPONSE TO BREAKTHROUGH ANTIEMETIC TREATMENT SUBSEQUENT CYCLES

The general principle of breakthrough treatment is to add one agent from a different drug class PRN to the current regimen.^e


- Benzodiazepine:
- ➤ Lorazepam 0.5 2 mg PO or IV either every 4 or every 6 h
- Cannabinoid:
- ➤ Dronabinol 5 10 mg PO either every 3 or every 6 h
- ➤ Nabilone 1 2 mg PO BID
- Other:

Any

nausea/

vomiting

- ➤ Haloperidol 0.5 2 mg PO or IV every 4 6 hk
- ➤ Metoclopramide 10 40 mg PO or IV either every 4 or every 6 h^k
- ➤ Olanzapine 2.5 5 mg PO BID (category 2B)^p
- ➤ Scopolamine 1 patch every 72 h
- Phenothiazine:
- ➤ Prochlorperazine 25 mg supp pr every 12 h or 10 mg PO or IV every 4 or every 6 h^k
- ➤ Promethazine 12.5 25 mg PO or IV central line only every 4 h^k
- Serotonin 5-HT3 antagonists:
- ➤ Dolasetron 100 mg PO daily
- ➤ Granisetron 1 2 mg PO daily or 1 mg PO BID or 0.01 mg/kg (maximum 1 mg) IV
- ➤ Ondansetron 16 mg PO or IV daily
- Steroid:
- ➤ Dexamethasone 12 mg PO or IV daily



<u>See Principles of Emesis Control</u> <u>for the Cancer Patient (AE-1)</u>

Note: All recommendations are category 2A unless otherwise indicated.

dSee Principles of Managing Multiday Emetogenic Chemotherapy Regimens (AE-A).

^eOrder of listed antiemetics is alphabetical and does not reflect preference.

^kMonitor for dystonic reactions; use diphenhydramine 25 - 50 mg PO or IV either every 4 or every 6 h for dystonic reactions. If allergic to diphenhydramine use benztropine at 1 - 2 mg IV or IM x 1 dose, followed by oral dose of 1 - 2 mg daily or BID if needed to control the reaction.

^oSee Principles of Managing Breakthrough Treatment (AE-B).

^pSee blackbox warning/label indication regarding type II diabetes, hyperglycemia, and death in elderly dementia patients.

Comprehensive NCCN Guidelines™ Version 1.2012 Cancer Network® Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

EMETOGENIC POTENTIAL OF INTRAVENOUS ANTINEOPLASTIC AGENTS[†]

LEVEL	AGENT		
High emetic risk (>90 % frequency of emesis) ^{q,r}	 AC combination defined as either doxorubicin or epirubicin with cyclophosphamide ^q Carmustine >250 mg/m² Cisplatin ≥50 mg/m² Cyclophosphamide >1,500 mg/m² Dacarbazine 	 Doxorubicin >60 mg/m² Epirubicin >90 mg/m² Ifosfamide ≥10 g/m² Mechlorethamine Streptozocin 	
Moderate emetic risk (30% - 90% frequency of emesis) ^{q,s}	 Aldesleukin > 12-15 million international units/m² Amifostine > 300 mg/m² Arsenic trioxide Azacitidine Bendamustine Busulfan Carboplatin^s Carmustine s ≤250 mg/m² Cisplatin < 50 mg/m² Clofarabine Cyclophosphamide ≤1500 mg/m² 	 Epirubicin^s ≤90 mg/m² Idarubicin Ifosfamide^s <10 g/m² Interferon alfa ≥10 million international units/m² Irinotecan^s Melphalan Methotrexate^s ≥ 250 mg/m² Oxaliplatin Temozolomide 	
	 Cytarabine >200 mg/m² Dactinomycin^s Daunorubicin^s Doxorubicin^s ≤60 mg/m² 	Low Emetic Risk (See AE-8) Minimal Emetic Risk (See AE-8)	
		Oral Chemotherapy (See AE-9)	

Adapted with permission from:

Hesketh PJ, et al. Proposal for classifying the acute emetogenicity of cancer chemotherapy. J Clin Oncol 1997;15: 103-109

Grunberg SM, Warr D, Gralla RJ, et al. Evaluation of new antiemetic agents and definition of antineoplastic agent emetogenicity-state of the art. Support Care Cancer. 2010 Oct 24. [Epub ahead of print].

Note: All recommendations are category 2A unless otherwise indicated.

^qProportion of patients who experience emesis in the absence of effective antiemetic prophylaxis.

^rContinuous infusion may make this agent less emetogenic.

^sThese agents may be highly emetogenic in certain patients.

^tPotential drug interactions between antineoplastic agents / antiemetic therapies and various other drugs should always be considered.



Comprehensive NCCN GuidelinesTM Version 1.2012 Cancer Network[®] Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

EMETOGENIC POTENTIAL OF INTRAVENOUS ANTINEOPLASTIC AGENTS[†]

LEVEL	AGENT	
Low emetic risk (10% - 30% frequency of emesis) ^q	 Amifostine ≤300 mg Aldesleukin ≤12 million international units/m² Cabazitaxel Cytarabine (low dose) 100-200 mg/m² Docetaxel Doxorubicin (liposomal) Eribulin Etoposide 5-Fluorouracil Floxuridine Gemcitabine Interferon alfa >5 <10 million international units/n Ixabepilone 	 Methotrexate >50 mg/m² <250 mg/m² Mitomycin Mitoxantrone Paclitaxel Paclitaxel-albumin Pemetrexed Pentostatin Pralatrexate Romidepsin Thiotepa Topotecan
Minimal emetic risk (<10% frequency of emesis) ^q	 Alemtuzumab Asparaginase Bevacizumab Bleomycin Bortezomib Cetuximab Cladribine (2-chlorodeoxyadenosine) Cytarabine <100 mg/m² Decitabine Denileukin diftitox Dexrazoxane Fludarabine Interferon alpha ≤5 million international units/m² Ipilimumab 	• Methotrexate ≤50 mg/m² • Nelarabine • Ofatumumab • Panitumumab • Pegaspargase • Peginterferon • Rituximab • Temsirolimus • Trastuzumab • Valrubicin • Vinblastine • Vincristine • Vinorelbine • Moderate Emetic Risk (See AE-7) Oral Chemotherapy (See AE-9)

Adapted with permission from:

Hesketh PJ, et al. Proposal for classifying the acute emetogenicity of cancer chemotherapy. J Clin Oncol 1997;15: 103-109.

Grunberg SM, Warr D, Gralla RJ, et al. Evaluation of new antiemetic agents and definition of antineoplastic agent emetogenicity-state of the art. Support Care Cancer. 2010 Oct 24. [Epub ahead of print].

^qProportion of patients who experience emesis in the absence of effective antiemetic prophylaxis.

^tPotential drug interactions between antineoplastic agents / antiemetic therapies and various other drugs should always be considered.

Note: All recommendations are category 2A unless otherwise indicated.



Comprehensive NCCN GuidelinesTM Version 1.2012 Cancer Nerwork[®] Antiemesis

NCCN Guidelines Index Antiemesis Table of Contents Discussion

EMETOGENIC POTENTIAL OF ORAL ANTINEOPLASTIC AGENTS^t

LEVEL	AGENT		
Moderate to High	 Altretamine Busulfan (≥4 mg/day) Cyclophosphamide (≥100 mg/m²/day) Estramustine Etoposide Lomustine (single day) Procarbazine Temozolomide (>75 mg/m²/day) 		
Minimal to Low	Bexarotene Busulfan (<4 mg/day) Capecitabine Chlorambucil Cyclophosphamide (<100 mg/m²/day) Dasatinib Erlotinib Erlotinib Fludarabine Gefitinib Hydroxyurea Imatinib Lapatinib Lenalidomide Melphalan	 Mercaptopurine Methotrexate Nilotinib Pazopanib Sorafenib Sunitinib Temozolomide (≤75 moderate) Thalidomide Thioguanine Topotecan Tretinoin Vandetanib Vorinostat 	ng/m ² /day) High Emetic Risk (See AE-7) Moderate Emetic Risk (See AE-7) Low Emetic Risk (See AE-8) Minimal Emetic Risk (See AE-8)

Adapted with permission from:

Hesketh PJ, et al. Proposal for classifying the acute emetogenicity of cancer chemotherapy. J Clin Oncol 1997;15: 103-109.

Grunberg SM, Warr D, Gralla RJ, et al. Evaluation of new antiemetic agents and definition of antineoplastic agent emetogenicity-state of the art. Support Care Cancer. 2010 Oct 24. [Epub ahead of print].

^tPotential drug interactions between antineoplastic agents / antiemetic therapies and various other drugs should always be considered.

Note: All recommendations are category 2A unless otherwise indicated.

Comprehensive NCCN GuidelinesTM Version 1.2012 Cancer Network[®] Antiemesis


NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

RADIATION-INDUCED EMESIS PREVENTION/TREATMENT

EMETOGENIC POTENTIAL

TYPE OF RADIATION THERAPY

BREAKTHROUGH TREATMENT



See Principles of Emesis Control for the Cancer Patient (AE-1)

^eOrder of listed antiemetics is alphabetical and does not reflect preference.

Note: All recommendations are category 2A unless otherwise indicated.

Comprehensive NCCN Guidelines M Version 1.2012 Cancer Network Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

ANTICIPATORY EMESIS PREVENTION/TREATMENT

Anticipatory nausea/vomiting

Prevention:

• Use optimal antiemetic therapy during every cycle of treatment

Behavioral therapy:

- Relaxation/systematic desensitization
- Hypnosis/guided imagery
- Music therapy

Acupuncture/acupressure

Alprazolam 0.5 - 2 mg PO TID beginning on the night before treatment or

Lorazepam 0.5 - 2 mg PO on the night before and morning of treatment

See Primary and Breakthrough Treatments for Chemotherapy-Induced Nausea/Vomiting (Antiemesis TOC)

See Principles of Emesis Control for the Cancer Patient (AE-1)

Note: All recommendations are category 2A unless otherwise indicated.

Comprehensive NCCN Guidelines™ Version 1.2012 Cancer Nerwork® Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

PRINCIPLES OF MANAGING MULTIDAY EMETOGENIC CHEMOTHERAPY REGIMENS

- Patients receiving multiday chemotherapy are at risk for both acute and delayed nausea/vomiting based upon the emetogenic potential of
 the individual chemotherapy agents and their sequence. It is therefore difficult to recommend a specific antiemetic regimen for each day,
 especially since acute and delayed emesis may overlap after the initial day of chemotherapy until the last day of chemotherapy. After
 chemotherapy administration concludes, the period of risk for delayed emesis also depends on the specific regimen and the emetogenic
 potential of the last chemotherapy agent administered in the regimen.
- Examples illustrating the above include BEP (bleomycin 30 units IV weekly, etoposide 100 mg/m² IV days 1 5 and cisplatin 20 mg/m² IV days 1 5) versus ASHAP (doxorubicin 25 mg/m² IV day 1, methylprednisolone 500 mg/day IV days 1 5, cisplatin 25 mg/m² IV continuous infusion days 1 4 followed by cytarabine 2000 mg/m² day 5). BEP is moderately emetogenic with risk for emesis on days 1 8) whereas ASHAP is moderately emetogenic on days 1 4 but becomes highly emetogenic on day 5 due to the administration of high-dose cytarabine). Risk for acute and delayed emesis for ASHAP may last up to 10 days.

Accordingly, the panel recommends the following as general principles:

A 5-HT3 receptor antagonist should be administered prior to the first dose of moderately or highly emetogenic chemotherapy (See AE-2 and AE-3).

- Dexamethasone should be administered once daily (either orally or intravenously) for moderately or highly emetogenic chemotherapy and for 2 3 days after chemotherapy for regimens that are likely to cause significant delayed emesis (see AE-2 and AE-3). Dexamethasone should not be added when the chemotherapy regimen already includes a corticosteroid (as in ASHAP illustrated above).
- Intravenous palonosetron may be used prior to the start of a 3 day chemotherapy regimen instead of multiple daily doses of oral or intravenous 5-HT3 receptor antagonists (see AE-2 and AE-3). Repeat dosing of palonosetron 0.25 mg IV is likely to be safe, based upon the dose-ranging phase II trial where up to 30 times the FDA-approved dose (90 mcg/kg) was administered and the 3 phase III trials that evaluated palonosetron 0.75 mg as a single fixed dose. Compared to the approved dose of palonosetron 0.25 mg, these higher doses were not associated with significantly different grades or durations of adverse events. In terms of efficacy, need for repeat dosing with palonosetron, either daily or less frequently, in the setting of multiday chemotherapy is not yet known.
- Aprepitant may be used for multiday chemotherapy regimens likely to be highly emetogenic and associated with significant risk for delayed nausea and emesis (see AE-2). Category 1 evidence is available for single-day chemotherapy regimens only, and aprepitant should be administered 125 mg orally 1 hour prior to chemotherapy on day 1, along with a 5-HT3 receptor antagonist and dexamethasone. Aprepitant 80 mg should be administered daily on days 2 and 3 after the start of chemotherapy along with dexamethasone. Based upon phase II data, aprepitant 80 mg may be safely administered on days 4 and 5 after multiday chemotherapy. It is not yet known if dosing aprepitant after day 3 improves control of nausea or emesis in this clinical setting. Note that fosaprepitant dimeglumine (115 mg) may be substituted for aprepitant (125 mg) on day 1 only. Alternatively, for highly emetogenic regimens, fosaprepitant 150 mg IV with recommended dexamethasone dosing may be given on day 1 with no need for oral aprepitant on days 2 and 3.

Note: All recommendations are category 2A unless otherwise indicated.

Comprehensive NCCN Guidelines™ Version 1.2012 Cancer Network® Antiemesis

NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

PRINCIPLES FOR MANAGING BREAKTHROUGH EMESIS

- Breakthrough emesis presents a difficult situation as correction of refractory ongoing nausea/vomiting is often challenging to reverse. It is generally far easier to prevent nausea/vomiting than to treat it.
- The general principle of breakthrough treatment is to give an additional agent from a different drug class. No one drug class has been shown to be superior for the management of breakthrough emesis, and the choice of agent should be based on assessment of the current prevention strategies used. Some patients may require several agents utilizing differing mechanisms of action.
- One should strongly consider routine, around-the-clock administration rather than PRN dosing.
- The PO route is not likely to be feasible due to ongoing vomiting, therefore, rectal or IV therapy is often required.
- Multiple concurrent agents, perhaps in alternating schedules or by alternating routes, may be necessary. Dopamine antagonists (eg, metoclopramide, haloperidol), corticosteroids, and agents such as lorazepam may be required.
- Ensure adequate hydration or fluid repletion, simultaneously checking and correcting any possible electrolyte abnormalities.
- Prior to administering the next cycle of chemotherapy the patient should be reassessed, with attention given to various possible non-chemotherapy-related reasons for breakthrough emesis with the current cycle:
- ➤ Brain metastases
- ➤ Electrolyte abnormalities
- > Tumor infiltration of the bowel or other gastrointestinal abnormality
- > Other comorbidities
- Prior to the next cycle of chemotherapy, reassess both the day 1 and post-chemo antiemetic regimen which did not protect the patient during the present cycle, and consider alternatives: (Suggestions are not in order of preference)
- ➤ Add aprepitant if not previously included.
- ▶ Add other concomitant antiemetics, eg, dopamine antagonists (metoclopramide or haloperidol).
- ➤ Possibly adjust dose(s), either intensity or frequency, of the 5-HT3 antagonist. Based on the patient's experiences, the chemotherapy regimen in question may actually be more emetogenic than generally classified (eg, Hesketh method).
- ➤ Possibly switch to a different 5-HT3. Although not necessarily likely to be effective, anecdotal and limited investigational trial data suggest it may sometimes be efficacious.
- ▶ If the goal of chemotherapy is non-curative, consider other appropriate regimens, if any, which might be less emetogenic.
- > It may be beneficial to add an anxiolytic agent in combination with the antiemetic agents.
- Consider antacid therapy if patient has dyspepsia (H2 blocker or proton pump inhibitor).

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

Discussion

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Overview

Chemotherapy-induced (or radiation-therapy induced) vomiting (emesis) and nausea can significantly affect a patient's quality of life, leading to poor compliance with further chemotherapy or radiation therapy treatment. In addition, nausea and vomiting can result in metabolic imbalances, degeneration of self-care and functional ability, nutrient depletion, anorexia, decline of the patient's performance status and mental status, wound dehiscence, esophageal tears, and withdrawal from potentially useful or curative anticancer treatment.¹⁻⁴

The incidence and severity of nausea and/or vomiting in patients receiving chemotherapy or radiation therapy (or both) are affected by numerous factors, including: (1) the specific chemotherapeutic agents used; (2) dosage of the agents; (3) schedule and route of administration of the agents; (4) target of the radiation therapy (e.g., whole body,

upper abdomen); and (5) individual patient variability (e.g., age, sex, prior chemotherapy, history of alcohol use).^{5, 6} More than 90% of patients receiving highly emetogenic chemotherapy will have episodes of vomiting. However, only about 30% of these patients will vomit if they receive prophylactic (preventive) antiemetic regimens before treatment with highly emetogenic chemotherapy.^{5, 7, 8} Although vomiting can often be prevented or substantially decreased by using prophylactic antiemetic regimens, nausea is much harder to control.^{9, 10} The NCCN Clinical Practice Guidelines in Oncology (NCCN GuidelinesTM) for Antiemesis is intended to provide an overview of the treatment principles for preventing chemotherapy- (or radiotherapy-) induced vomiting and nausea, and recommendations for antiemetic prophylaxis according to the emetogenic potential of anti-tumor therapies. The NCCN Guidelines for Antiemesis is updated on a yearly basis by a multidisciplinary panel of experts.

Pathophysiology of Emesis

Vomiting results from stimulation of a multistep reflex pathway controlled by the brain.⁵ Vomiting is triggered by afferent impulses to the vomiting center (located in the medulla) from the chemoreceptor trigger zone, pharynx and gastrointestinal (GI) tract (via vagal afferent fibers), and cerebral cortex. Vomiting occurs when efferent impulses are sent from the vomiting center to the salivation center, abdominal muscles, respiratory center, and cranial nerves.¹¹

The chemoreceptor trigger zone, vomiting center, and GI tract have many neurotransmitter receptors. Activation of these receptors by chemotherapeutic agents or their metabolites may be responsible for chemotherapy-induced emesis. The principal neuroreceptors involved in the emetic response are the serotonin (5-hydroxytryptamine [5-HT3]) and dopamine receptors. ^{12, 13} Other neuroreceptors involved in emesis



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

include acetylcholine, corticosteroid, histamine, cannabinoid, opiate, and neurokinin-1 (NK-1) receptors, which are located in the vomiting and vestibular centers of the brain.¹⁴

Antiemetic agents can block different neuronal pathways, exert their effects at different points during the course of emesis, or behave synergistically with other antiemetic agents to potentiate an antiemetic effect. When used at a certain concentration, each antiemetic agent predominantly blocks one receptor type. A final common pathway for emesis has yet to be identified. Therefore, no single agent can be expected to provide complete protection from the various emetic phases of chemotherapy.

Nausea

With use of effective antiemetic regimens, patients receiving emetogenic chemotherapy often experience more nausea than vomiting. Total Vomiting and nausea are related; however, they may occur via different mechanisms. In general, younger patients are more likely to have nausea than older patients. Younger women receiving chemotherapy for breast cancer are more prone to nausea than other populations. Delayed nausea is more common than acute nausea, is often more severe, and tends to be resistant to treatment.

Types of Nausea and/or Vomiting

Chemotherapy-Induced Nausea and/or Vomiting

Nausea and/or vomiting induced by chemotherapy is commonly classified as acute, delayed, anticipatory, breakthrough, or refractory. *Acute-onset* nausea and/or vomiting usually occurs within a few minutes to several hours after drug administration and commonly resolves within the first 24 hours. The intensity of acute-onset emesis generally peaks after 5 to 6 hours. The occurrence of acute emesis is

influenced by the patient's age and gender (females and younger patients [age < 50 years] are more prone to emesis), environment in which chemotherapy is administered, whether the patient has a history of chronic alcoholism (which decreases the incidence of emesis) or motion sickness, previous episodes of nausea and vomiting, dosage of the emetogenic agent, and efficacy of the antiemetic regimen.^{18, 19}

Delayed-onset nausea and/or vomiting develops in patients more than 24 hours after chemotherapy administration. ^{18, 19} It occurs commonly with the administration of cisplatin, carboplatin, cyclophosphamide, and/or doxorubicin. For cisplatin, emesis reaches its maximal intensity 48 to 72 hours after administration and can last 6 to 7 days.

Anticipatory nausea and/or vomiting occurs before patients receive their next chemotherapy treatment. Because it is primarily considered a conditioned response, anticipatory emesis typically occurs after a negative past experience with chemotherapy. The incidence of anticipatory nausea and/or vomiting ranges from 18% to 57%, and nausea is more common than vomiting. Younger patients may be more susceptible to anticipatory nausea and vomiting, because they generally receive more aggressive chemotherapy and, overall, have poorer emesis control than older patients. Perakthrough emesis refers to vomiting that occurs despite prophylactic treatment and/or requires "rescue" with antiemetic agents. Refractory emesis refers to emesis that occurs during subsequent treatment cycles when antiemetic prophylaxis and/or rescue have failed in earlier cycles.

Radiation-Induced Nausea and/or Vomiting

Patients receiving whole body or upper abdominal radiation therapy have the greatest likelihood of developing nausea and/or vomiting. The GI tract (specifically, the small intestine) contains rapidly dividing cells that are particularly sensitive to radiation. In addition, the potential



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

for nausea and/or vomiting increases with larger daily fractional doses of radiotherapy, larger total doses, and larger amounts of irradiated tissue. Total body irradiation, when given before bone marrow transplantation, commonly induces nausea and/or vomiting.^{23, 26}

Emetogenicity of Chemotherapy

The frequency of chemotherapy-induced emesis depends primarily on the emetogenic potential of the specific chemotherapeutic agents used. Several classifications have been developed to define the emetogenicity of chemotherapy; however, none has been universally accepted. 11, 27-30

Hesketh and colleagues developed a classification of the acute emetogenicity of anticancer chemotherapeutic agents and developed an algorithm to define the emetogenicity of combination chemotherapeutic regimens. The classification was recently updated by Grunberg and colleagues; it divides chemotherapeutic agents into 4 levels according to the percentage of patients not receiving antiemetic prophylaxis who experience acute emesis. This classification, which is updated each year by the NCCN Guidelines Panel with recently introduced drugs, is used in these guidelines. Experts representing the panels of all of the published antiemetic treatment guidelines met to prepare a single consensus document. Although this process is ongoing, the consensus guidelines have been published. Although the CN Guidelines currently outline treatment using 4 categories of emetogenic potential for intravenous (IV) agents, which correspond to the Grunberg classification as follows:

- High emetic risk—90% or more of patients experience acute emesis;
- Moderate emetic risk—30% to 90% of patients experience acute emesis;
- Low emetic risk—10% to 30% of patients experience acute emesis;

 Minimal emetic risk—fewer than 10% of patients experience acute emesis.

In addition, the NCCN Guidelines attempt to define antiemetic regimens for particular chemotherapy drugs that cover the entire duration of time a patient is at risk for nausea and vomiting. Panel members were concerned that some patients may not receive adequate prophylaxis for delayed emesis; therefore, the NCCN Guidelines incorporate a dosing schedule that covers both acute and delayed emesis into a single algorithm. The NCCN Guidelines Panel members have also categorized the emetogenic potential of oral antineoplastic agents.⁹

Types of Antiemetic Therapies

In general, to provide maximal protection against chemotherapy-induced emesis, antiemetic therapy should be initiated before chemotherapy. The antiemetic therapy should also be continued for the same length of time as the duration of the emetic activity of the chemotherapeutic agent being used. However, daily use of antiemetics is not recommended for some therapeutic agents that are taken long term (e.g., imatinib, erlotinib). Antiemetic agents can be administered by the oral, rectal, IV, intramuscular, or transdermal route. Oral and IV 5-HT3 antagonists have equivalent efficacy when used at the appropriate doses. ^{8, 26} For patients unable to swallow or digest tablets because of emesis, IV antiemetics are required. In selected patients who are unable to swallow, transdermal antiemetics may be of value. Although studies may show drugs to be equally effective on a population basis, individual patients may respond differently. Therefore, some drug options may be based on a patient's individual experience.



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

Serotonin (5-HT3) Receptor Antagonists

The development of the 5-HT3–receptor antagonists (i.e., dolasetron mesylate, granisetron, ondansetron, palonosetron) represents a significant advance in antiemetic therapy. 33-35 All of these agents have been shown to be effective in controlling the acute nausea and/or vomiting associated with cancer chemotherapy. 35-49

Palonosetron is a 5-HT3 antagonist with an approximately 100-fold higher binding affinity for the 5-HT3 receptor compared to the other serotonin antagonists (i.e., ondansetron, granisetron, and dolasetron). It has a half-life of approximately 40 hours, which is significantly longer than other commercially available 5-HT3 antagonists.³⁵ Data suggest that palonosetron is associated with prolonged inhibition of the 5-HT3 receptor and thus differs from other 5-HT3 antagonists (e.g., ondansetron, granisetron).⁵⁰

Several large, multicenter, double-blind, randomized phase III trials have demonstrated the superiority of palonosetron compared with other 5-HT3 antagonists in preventing emesis associated with both moderate and high emetic risk chemotherapy regimens, particularly for delayed emesis.⁵¹⁻⁵⁴ In these studies, the primary efficacy end point was complete response (CR), defined as having no emesis and no rescue treatments. A study in patients receiving moderately emetogenic chemotherapy (N=569 evaluable) showed that a single dose of palonosetron (0.25 mg IV) was comparable to a single dose of dolasetron (100 mg IV) for the prevention of acute chemotherapyinduced nausea and emesis (CR rate 63% vs 53%, respectively); moreover, IV palonosetron was superior to dolasetron in preventing delayed emesis (CR rate 54% vs 39%; P=0.004).⁵² Approximately 60% of patients in the palonosetron arms and 70% in the dolasetron arm had received anthracycline in combination with cyclophosphamide; only 6% and 5% of patients, respectively, received concomitant

corticosteroids.⁵² In another study in patients receiving moderately emetogenic chemotherapy (N=563 evaluable), a single dose of palonosetron (0.25 mg IV) was found to be superior to a single dose of ondansetron (32 mg IV) in preventing both acute (CR rate 81% vs 69%; P<0.01) and delayed emesis (CR rate 74% vs 55%; P<0.01); no concomitant corticosteroids were given in this study 53 The safety and side-effect profiles of palonosetron were indistinguishable from the control 5-HT3 antagonists (ondansetron and dolasetron) using data submitted to the Food and Drug Administration (FDA). In a phase III randomized trial that compared palonosetron with ondansetron in patients receiving highly emetogenic chemotherapy (N=667), the majority (67%) had received dexamethasone on day 1 of antiemetic therapy.⁵¹ Among this subgroup of patients who received concomitant dexamethasone (n=447), palonosetron (0.25 mg IV) was similar to ondansetron (32 mg IV) in preventing acute emesis (CR rate 65% vs 56%), but significantly more effective in preventing delayed emesis (CR rate 41% vs 25%; P=0.021). In a more recent phase III randomized trial that compared palonosetron (at a higher dose of 0.75 mg IV) with granisetron (40 mcg/kg IV), both in combination with dexamethasone, in patients treated with highly emetogenic chemotherapy (N=1114 evaluable), palonosetron showed similar activity to granisetron in preventing acute emesis (CR rate 75% vs 73%), with superior activity in preventing delayed emesis (CR rate 57% vs 44.5%; P<0.0001).54 Intravenous palonosetron is FDA approved as a single dose on day 1 for the prevention of acute and delayed nausea and vomiting associated with moderately emetogenic chemotherapy and for the prevention of acute nausea and vomiting associated with highly emetogenic chemotherapy. It is the preferred 5-HT3 antagonist for the prevention of acute and delayed emesis associated with high emetic risk IV chemotherapy and is also recommended (category 1) for emesis prevention when using moderate emetic risk IV chemotherapy (see



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

Guidelines section on "High Emetic Risk Intravenous Chemotherapy - Acute and Delayed Emesis Prevention" and "Moderate Emetic Risk Intravenous Chemotherapy – Emesis Prevention; also see Discussion section below for "Prevention of Acute and Delayed Emesis"). It should be noted that the recommendation for palonosetron as the preferred 5-HT3 antagonist for antiemetic prophylaxis in the setting of high emetic risk chemotherapy is based upon data from randomized studies (discussed earlier) with the 2-drug combination of palonosetron and dexamethasone.

Intravenous palonosetron is superior to other 5-HT3 antagonists for preventing delayed nausea. ^{15, 51-54} Repeat dosing of palonosetron in the days after chemotherapy (i.e., days 2 or 3) is likely to be safe. However, in the setting of multiple day (i.e., multiday) chemotherapy, need for repeat dosing with palonosetron is not yet known (see Guidelines section on "Managing Multiday Emetogenic Chemotherapy Regimens").

Many of the 5-HT3 antagonists can be delivered orally or intravenously. Although oral palonosetron has been approved by the FDA for moderately emetic risk chemotherapy, it is not available in the United States. Note that the IV dolasetron is no longer recommended for the prevention of nausea and vomiting in the NCCN 2011 Antiemesis algorithm, because IV dolasetron has been associated with an increased risk for cardiac arrhythmias . Oral dolasetron is still recommended.

In addition, the FDA has approved the use of a granisetron transdermal system for chemotherapy-induced nausea and vomiting .⁵⁷ The patch containing 34.3 mg of granisetron is applied approximately 24 to 48 hours before the first dose of chemotherapy; the maximum duration of the patch is 7 days. A phase III randomized study compared the patch to oral granisetron in patients receiving either highly emetogenic or

moderately emetogenic chemotherapy. The patch proved non-inferior to repeat dosing of the oral antiemetic granisetron over 3 to 5 days.^{58, 59}

Many clinical trials directly comparing ondansetron, granisetron, dolasetron mesylate, and palonosetron have been conducted. These trials have used various doses, routes, and schedules of administration. A meta-analysis found no difference in efficacy between ondansetron, granisetron, and dolasetron mesylate. A recent meta-analysis of studies comparing ondansetron with granisetron has also confirmed the similar efficacy of these 5-HT3 antagonists in controlling acute and delayed nausea and vomiting, with similar safety profiles between these agents. The most recent meta-analysis of randomized controlled trials comparing palonosetron with other available 5-HT3 antagonists demonstrated that palonosetron was significantly more effective in preventing acute and delayed nausea and vomiting for both moderately and highly emetogenic chemotherapy agents.

The addition of dexamethasone improves the efficacy of the antiemetic regimen containing 5-HT3 antagonists; however, dexamethasone is associated with side effects (such as insomnia). A recent randomized trial suggests that the dose of dexamethasone can be decreased to 8 mg on day 1 and also eliminated on days 2-3 when used with palonosetron for moderately emetic chemotherapy.⁷⁷

Ondansetron, granisetron, and dolasetron are effective in preventing acute emesis but appear to be less effective for delayed emesis. A meta-analysis of randomized controlled trials found that adding a 5-HT3 antagonist to dexamethasone did not improve the antiemetic effect of dexamethasone for preventing delayed emesis.⁷⁸ Another study found that 5-HT3 antagonists (except palonosetron, which was not studied) were not more effective than prochlorperazine for preventing delayed



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

emesis.¹⁷ Intravenous palonosetron appears to be effective for preventing both delayed and acute emesis.

Neurokinin-1-Receptor Antagonist

Aprepitant selectively blocks the binding of substance P at the NK-1 receptor in the central nervous system. Thus, aprepitant provides a different and complementary mechanism of action to all other commercially available antiemetics. Aprepitant has been shown to augment the antiemetic activity of the 5-HT3-receptor antagonists and the corticosteroid dexamethasone to prevent both acute and delayed cisplatin-induced emesis. 79-81 In a randomized phase III trial comparing a standard antiemetic regimen (ondansetron 32 mg IV and oral dexamethasone) with or without the addition of aprepitant in patients receiving emetogenic chemotherapy with high-dose cisplatin (N=521 evaluable), the addition of aprepitant was significantly more effective than the 2-drug regimen in controlling both acute (CR rate 89% vs 78%; P<0.001) and delayed emesis (CR rate 75% vs 56%; P<0.001).80 Another similarly designed randomized phase III study (N=523 evaluable) also showed a significant benefit of adding aprepitant to ondansetron and dexamethasone compared with the 2-drug regimen alone for controlling both acute (CR rate 83% vs 68%; P<0.001) and delayed emesis (CR rate 68% vs 47%; P<0.001).81 A pooled analysis of data combined from these two phase III trials found that the aprepitant regimen was particularly beneficial in improving CR rates for patients receiving concomitant emetogenic therapy with doxorubicin and/or cyclophosphamide, along with high-dose cisplatin therapy.⁷⁹

A meta-analysis (of 7 randomized controlled trials) in patients receiving highly emetogenic chemotherapy found that NK-1 receptor antagonists used alone or with standard therapy did not significantly increase protection from acute emesis or nausea; however, for delayed emesis

and nausea, NK-1 receptor antagonists was associated with significantly increased protection compared with control.⁸²

A randomized phase III trial (N=866) showed that an aprepitant regimen was more effective than a standard regimen for preventing vomiting in patients receiving moderately emetogenic chemotherapy (non-cisplatin based) during 120 hours after initiation of chemotherapy (CR rate 51% vs 43%, *P*=0.015); However, approximately 40% of patients (receiving either regimen) still experienced significant nausea.⁸³ The aprepitant regimen included aprepitant, ondansetron, and dexamethasone; the standard regimen included ondansetron and dexamethasone.

A phase II study (N=58) found that combining palonosetron, aprepitant, and dexamethasone was effective for preventing both acute and delayed emesis and nausea when using various chemotherapeutic regimens (moderate to moderate-highly emetogenic); 78% of patients had a CR (no emetic episodes and no rescue medication) during the overall time frame, from 0 to 120 hours after initiation of emetogenic therapy.⁸⁴ A phase II study in patients with breast cancer (N=41) receiving moderately emetogenic chemotherapy also found that a single-day regimen of palonosetron (0.25 mg IV), aprepitant (285 mg oral), and dexamethasone (20 mg) was effective; 76% and 66% of patients had a CR during the acute and delayed phases, respectively.⁸⁵

Oral aprepitant is approved by the FDA for the prevention of nausea and vomiting in patients receiving highly (e.g., cisplatin-containing) and moderately emetogenic chemotherapy. ⁸⁶ The oral doses of aprepitant are 125 mg on day 1 (before chemotherapy) and then 80 mg on days 2 and 3 (after chemotherapy). ^{86, 87} An IV version of aprepitant (fosaprepitant dimeglumine), which can be given on day 1 only, is also approved by the FDA. IV fosaprepitant is given 30 minutes before



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

chemotherapy on day 1 only, per the package insert.⁸⁸ If a higher dose of fosaprepitant is used (150 mg IV) on day 1, then it is not necessary to give oral aprepitant on days 2-3.^{89, 90} Note that the dexamethasone dosing is slightly different on days 3 and 4 (8 mg PO twice daily) when using the higher dose of fosaprepitant (150 mg IV) per the package insert.⁸⁸ A single dose of 150 mg IV fosaprepitant was shown to be non-inferior to the standard regimen with 3-day oral aprepitant in a recent randomized study.⁹¹ There are no studies showing efficacy or safety of chronic dosing with aprepitant. It is possible that the drug-drug interaction profile may change with chronic dosing.

Drug Interactions

Aprepitant is simultaneously a substrate, moderate inducer, and moderate inhibitor of cytochrome P450 enzyme 3A4 (CYP3A4); aprepitant also induces CYP2C9. 92 Thus, aprepitant can alter the metabolism of certain drugs and change their plasma concentrations (i.e., AUCs [area under the curve]). These interactions are more significant with orally administered forms of these drugs than with IV forms because of first-pass metabolism. Patients should not take aprepitant with pimozide, terfenadine, astemizole, or cisapride; these combinations are contraindicated, because they may cause "serious or life-threatening reactions" (see the aprepitant package insert)⁸⁶ Chemotherapeutic agents known to be metabolized by CYP3A4 include docetaxel, paclitaxel, etoposide, irinotecan, ifosfamide, imatinib, vinorelbine, vinblastine, and vincristine. In clinical trials, aprepitant was used concurrently with etoposide, vinorelbine, or paclitaxel; although chemotherapy doses were not adjusted for potential drug interactions in phase III trials, caution is urged when using any chemotherapeutic agent that is metabolized by CYP3A4. Aprepitant has been shown to interact with several nonchemotherapeutic drugs (including warfarin, dexamethasone, methylprednisolone, oral contraceptives). Again, these interactions are more significant with orally administered forms of these drugs than with IV forms because of first-pass metabolism.

Induction of warfarin metabolism by aprepitant may lead to clinically significant reductions in INR (international normalized ratio) values, particularly for patients on therapeutic (as compared to prophylactic) warfarin regimens. These changes, although brief in duration, may require increased patient monitoring. Aprepitant decreases the AUC for patients taking oral contraceptives; thus, other methods of birth control should be used during treatment with aprepitant and for 1 month after the last dose of aprepitant. Additionally, certain drugs can affect the AUCs of aprepitant. Concomitant administration with CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, and erythromycin) may lead to increased aprepitant AUCs, whereas concomitant administration with CYP3A4 inducers (e.g., carbamazepine, rifampin, and phenytoin) may lead to decreased levels of aprepitant.

Other Non-5-HT3-Receptor Antagonist Antiemetics

Before the advent of the 5-HT3–receptor antagonists, the available antiemetic agents included phenothiazines, 93 substituted benzamides, 94, 95 antihistamines, 96 butyrophenones, 97 corticosteroids, 98-100 benzodiazepines, 101, 102 and cannabinoids. 103, 104 Most drugs used to prevent chemotherapy-induced emesis are classified as dopamine antagonists, serotonin antagonists, and other antagonists. Combination antiemetic therapy is more effective than single-agent therapy.

Olanzapine (thiobenzodiazepine) was found to be effective for preventing acute and delayed emesis in a phase II trial in patients (N=30) who received cyclophosphamide, doxorubicin, and/or cisplatin; other studies have also showed the value of olanzapine for delayed and refractory emesis and nausea. However, olanzapine should be used with caution in elderly patients (see boxed



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

warning/label indication regarding death in patients with dementiarelated psychosis and additional warnings and precautions about type II diabetes and hyperglycemia).^{110, 111}

Treatment Issues

As new data on the use of antiemetics in patients receiving chemotherapy become available, clinicians should consider these data when caring for such patients, even if the information has not been included in the guidelines. In contrast to other NCCN Guidelines in which most of the recommendations are category 2A, many of the recommendations for antiemetic management are classified as category 1, reflecting the large number of randomized controlled trials that have focused on antiemetic management.

Principles of Emesis Control

These principles are described in the algorithm and are summarized here (see Guidelines section on "Principles of Emesis Control for the Cancer Patient"). The goal of emesis control is to prevent nausea and/or vomiting. Antiemetic regimens should be chosen based on the drug with the highest emetic risk in the chemotherapy regimen, previous experience with antiemetics, and patient-specific risk factors. Patients need to be protected throughout the entire period of risk, which lasts for at least 3 days for high emetic risk and 2 days for moderate emetic risk agents after the last dose of chemotherapy.

In addition to using antiemetic regimens, patients can adjust their eating habits and adopt other lifestyle measures that may alleviate nausea and vomiting (see "Eating Hints: Before, During, and After Cancer Treatment" from the National Cancer Institute). Suggestions include eating small frequent meals, food that is "easy on the stomach", full

liquid foods, and food at room temperature; patients can also avoid foods that make them feel nauseous.

Prevention of Acute and Delayed Emesis

To prevent acute emesis, antiemetic therapy should start before the administration of chemotherapy and then should cover the first 24 hours. In the NCCN Guidelines for Antiemesis, the specific antiemetic regimens are described for highly emetogenic IV drugs, moderately emetogenic IV drugs, low, and minimally emetogenic IV drugs. Emesis prevention for oral chemotherapeutic agents is also described in the NCCN Guidelines. This section discusses prechemotherapy and postchemotherapy emesis prevention rather than primary treatment.

Prechemotherapy Emesis Prevention

The NCCN Guidelines specify different prophylactic antiemetic regimens for cancer patients receiving chemotherapy of different emetogenic potential (i.e., high, moderate, low, and minimal). Prophylactic antiemetics should be administered before chemotherapy. The recommendations for prophylactic antiemetic treatment include drug dosages. The guidelines reflect accumulating experience with the 5-HT3 antagonists, demonstrating their effectiveness in a range of doses. Unless indicated, the order of listed antiemetics in the NCCN Guidelines does not reflect preference.

Highly emetogenic IV drugs in the NCCN Guidelines include carmustine $> 250 \text{ mg/m}^2$, cisplatin $\ge 50 \text{ mg/m}^2$, cyclophosphamide $> 1500 \text{ mg/m}^2$, dacarbazine, doxorubicin $> 60 \text{ mg/m}^2$, epirubicin $> 90 \text{ mg/m}^2$, ifosfamide $\ge 10 \text{ g/m}^2$, mechlorethamine, streptozocin, or anthracycline plus cyclophosphamide (AC) combinations (e.g., doxorubicin or epirubicin with cyclophosphamide). Although most of these drugs are also considered highly emetogenic by the Multinational Association of Supportive Care in Cancer/European Society of Medical Oncology



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

(MASCC/ESMO) guidelines,⁸ the NCCN Guidelines for highly, moderately, low, and minimally emetogenic agents differ slightly based on the experience and expertise of the panel members.

The antiemetic regimen for these highly emetogenic drugs on day 1 includes aprepitant (or fosaprepitant), dexamethasone, and a 5-HT3 antagonist with or without lorazepam and with or without either an H2 blocker or a proton pump inhibitor (category 1 for the combined regimen);^{23, 26, 80} note that the regimen and doses are often modified on days 2 to 4 after chemotherapy. Although it is not recommended as a single agent, lorazepam is a useful adjuvant because it decreases anxiety.^{26, 102} Lorazepam is also recommended for patients who are at risk for anticipatory nausea and/or vomiting (see Guidelines section on "Anticipatory Emesis Prevention/Treatment"). Antacid therapy (e.g., proton pump inhibitors, H2 blockers) should be considered if patients have dyspepsia, because patients sometimes have difficulty discriminating heartburn from nausea.

For IV regimens with high emetogenic potential, aprepitant is used at an oral dosage of 125 mg on day 1 and then 80 mg on days 2 and 3. When given with aprepitant, dexamethasone is used at a dosage of 12 mg on day 1; the dose can be oral or IV. Note that IV fosaprepitant may be substituted for oral aprepitant on day 1 only. If appropriate, lorazepam (0.5-2 mg either every 4 or every 6 hours on days 1-4; either oral, IV, or sublingual) may be used with each of these regimens (i.e., high, moderate, or low). As previously discussed, a recent phase III randomized trial suggested that palonosetron is preferred over granisetron for high emetic risk chemotherapy in combination with dexamethasone. This trial has been criticized because: 1) the control arm was not adequately dosed; thus, the trial "stacked the deck" in favor of palonosetron; 2) a larger non-FDA-approved dose of palonosetron was used (i.e., 0.75 mg IV); and 3) aprepitant was not

used in this study. The superiority of palonosetron over other available 5-HT3 antagonists in preventing acute and delayed nausea and vomiting in the setting of high emetogenic chemotherapy was demonstrated in a recent meta-analysis of randomized controlled trials. Therefore, the NCCN Guidelines Panel recommends palonosetron as the preferred 5-HT3 antagonists for high emetic risk chemotherapy. As previously noted, the recommendation for palonosetron as the preferred 5-HT3 antagonist for antiemetic prophylaxis in this setting is based upon data from randomized studies (discussed earlier) with the 2-drug combination of palonosetron and dexamethasone.

A Canadian meta-analysis suggested that the use of 5-HT3 antagonists (i.e., ondansetron) on days 2 to 4 to prevent delayed emesis was not cost effective; however, ondansetron (when used alone) did protect against delayed emesis in this meta-analysis. Palonosetron was not assessed in these studies. The NCCN Guidelines Panel recommends the use of 5-HT3 antagonists as one of several options to prevent delayed emesis for moderately emetogenic agents.

The antiemetic regimen for moderately emetogenic IV drugs on day 1 includes dexamethasone and a 5-HT3 antagonist with or without lorazepam and with or without either an H2 blocker or a proton pump inhibitor. Aprepitant (or fosaprepitant) should be added (to dexamethasone and a 5-HT3 antagonist) for select patients receiving certain moderate emetic risk chemotherapy (i.e., carboplatin, cisplatin, doxorubicin, epirubicin, ifosfamide, irinotecan, or methotrexate), because these agents are more emetogenic than the other moderately emetogenic agents. IV fosaprepitant may be substituted for oral aprepitant on day 1 only. Any one of the 5-HT3 antagonists can be used, because they are all category 1 for day 1. However, as previously mentioned, palonosetron was shown in a recent meta-analysis to be



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

more effective than other available 5-HT3 antagonists in preventing acute and delayed nausea and vomiting for both high and moderately emetogenic chemotherapy agents⁷⁶; hence, the NCCN Panel recommends palonosetron as the preferred 5-HT3 antagonist in the setting of moderately emetogenic chemotherapy.

The antiemetic regimen for low emetogenic IV drugs (see the NCCN 2011 Antiemesis algorithm) includes agents such as dexamethasone, prochlorperazine, or metoclopramide, with or without lorazepam, and with or without either an H2 blocker or a proton pump inhibitor. When using prochlorperazine or metoclopramide, patients should be monitored for dystonic reactions. 114-116 Diphenhydramine can be used for dystonic reactions. 117, 118 Benztropine may be used in patients who are allergic to diphenhydramine. 115 If appropriate, lorazepam (0.5-2 mg either every 4 or every 6 hours on days 1-4; either oral, IV, or sublingual) may be used with each of these regimens (i.e., high, moderate, or low).

The emetogenic potential of oral chemotherapeutic agents is shown in the NCCN Guidelines. Antiemetic prophylaxis is recommended for the following oral agents: altretamine, busulfan (4 mg/day or more), cyclophosphamide (100 mg/m²/day or more), estramustine, etoposide, lomustine (single day), procarbazine, and temozolomide (> 75 mg/m²/day). For high or moderate emetic risk oral agents, recommended prophylaxis includes oral 5-HT3 antagonists (such as granisetron or ondansetron) with or without lorazepam and with or without either an H2 blocker or a proton pump inhibitor. For low or minimal emetic risk oral agents, recommended prophylaxis includes metoclopramide, prochlorperazine, or haloperidol with or without lorazepam and with or without either an H2 blocker or a proton pump inhibitor; prophylaxis is given before chemotherapy is started and then on an as needed basis only (i.e., PRN).

Postchemotherapy/Delayed Emesis Prevention

The best management for delayed emesis is prevention. 119 For chemotherapeutic agents with high emetogenic potential, the prophylactic treatment (i.e., dexamethasone and aprepitant) is continued through the period when delayed emesis may occur. Using this strategy, prophylaxis continues for 2 to 4 days after completion of a chemotherapy cycle. However, 5-HT3 antagonists are given on day 1 only.

For drugs with moderate emetogenic potential, postchemotherapy prevention depends on which antiemetics were used before chemotherapy. For example, palonosetron (category 1) is only administered on day 1.53 If either aprepitant or fosaprepitant was used on day 1, then aprepitant is continued on days 2 and 3.

The antiemetic regimens in the NCCN Guidelines include different options on days 2 to 3 for moderate emetic risk agents. ^{23, 26, 119} There are 3 possible regimens on days 2-3 including: 1) aprepitant; 2) dexamethasone; or 3) 5-HT3 antagonist, such as ondansetron, granisetron, or dolasetron. 119 Each of these regimens may also include the following: ± lorazepam and ± either an H2 blocker or a proton pump inhibitor. It is important to note that the doses of both aprepitant (80 mg PO) and dexamethasone (8 mg PO or IV) are decreased when used on days 2-3 (when compared with the doses given on day 1). Note that palonosetron is not given on days 2-3.

The NCCN, MASCC, and American Society of Clinical Oncology (ASCO) guidelines all recommend using aprepitant to prevent delayed nausea and/or vomiting when giving AC regimens.^{8, 26, 120} Note that the MASCC guidelines are updated on a biannual basis following the publication of the initial consensus guidelines, which were based on the



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

Perugia Consensus Conference on Antiemetic Therapy held in June 2009.

Breakthrough Treatment

Breakthrough emesis presents a difficult situation, because refractory ongoing nausea and/or vomiting is often challenging to reverse (see Guidelines section on "Principles for Managing Breakthrough Emesis"). Generally, it is much easier to prevent nausea and/or vomiting than to treat it. Thus, routine around-the-clock administration of antiemetics should be strongly considered to prevent emesis, rather than PRN (as required) dosing. The general principle of breakthrough treatment is to give an additional agent as needed from a different drug class.²³ However, no single treatment is better than another for managing breakthrough emesis. Some patients may require several agents using different mechanisms of action. The oral route may not be feasible because of ongoing vomiting; therefore, rectal or IV therapy is often required. Nasal sprays might be useful for treatment of breakthrough emesis, because they provide acute delivery of agents. 121, 122 Multiple concurrent agents, perhaps in alternating schedules or by alternating routes, may be necessary.

Miscellaneous agents (e.g., haloperidol, metoclopramide, olanzapine, scopolamine transdermal patch), corticosteroids, and agents such as lorazepam may be incorporated for breakthrough treatment. Dronabinol and nabilone (which are cannabinoids) are approved by the FDA for nausea and vomiting in patients who have not responded to conventional antiemetic agents. Adequate hydration or fluid repletion should be ensured, and any possible electrolyte abnormalities should be assessed and corrected. Before administering the next cycle of chemotherapy, the patient should be reassessed for other possible nonchemotherapy-related reasons for breakthrough emesis with the current cycle (e.g., brain metastases, electrolyte abnormalities, tumor

infiltration of the bowel or other GI abnormality, and other comorbidities; see Guidelines section on "Principles for Managing Breakthrough Emesis").

In addition, before the next cycle of chemotherapy, the antiemetic regimen (both the day 1 and postchemotherapeutic) that did not protect the patient during the present cycle should be assessed and alternatives should be considered (see Guidelines section on "Principles for Managing Breakthrough Emesis"). Because patients sometimes have difficulty discriminating heartburn from nausea, use of antacid therapy (e.g., proton pump inhibitors, H2 blockers) should be considered.

Radiation-Induced Nausea and/or Vomiting

Prophylaxis for radiation-induced nausea and/or vomiting is based on the site of radiation and whether it is combined with chemotherapy.²⁴ 123, 124 When radiation is combined with chemotherapy, prophylaxis is dictated by the emetogenic potential of the chemotherapy regimen. Recent MASCC/ESMO guidelines state that total body irradiation is associated with the highest risk for emesis and that upper abdominal radiation is associated with moderate risk.²⁴ A recent meta-analysis suggests that 5-HT3 antagonists are the preferred agents for preventing radiation-induced vomiting. 125

Patients undergoing radiation therapy to the upper abdomen may receive antiemetic prophylaxis with oral ondansetron or oral granisetron, with or without oral dexamethasone.^{8, 24} A randomized study compared oral ondansetron with placebo in patients receiving daily fractionated radiotherapy including the abdomen. In this study, 67% of patients given ondansetron had complete control of emesis compared with 45% of patients who received placebo (P<0.05). 126 A study showed that the addition of oral dexamethasone (4 mg daily) to



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

the ondansetron regimen decreases emesis and nausea, although the effect is modest. 127 Another randomized study in patients receiving radiotherapy to the upper abdomen found that oral granisetron decreased emesis and nausea when compared with placebo. 128

Patients undergoing total body irradiation may receive antiemetic prophylaxis with either ondansetron or granisetron; either agent can be given with or without oral dexamethasone. 8, 24, 129 Treatment of breakthrough radiation-induced emesis is similar to chemotherapyinduced emesis. Patients who do not receive primary prophylaxis and experience breakthrough nausea and/or vomiting may be treated with ondansetron, similar to primary prophylaxis.

Anticipatory Nausea and/or Vomiting

About 20% of patients develop anticipatory nausea and/or vomiting. However, the rate of anticipatory nausea and/or vomiting appears to be decreasing (when compared with older studies) with current use of more effective antiemetic regimens.8 The most effective way to treat anticipatory nausea and/or vomiting is to prevent it by using optimal antiemetic therapy during every cycle of treatment.^{23, 130, 131} Behavioral therapy has been used in patients with anticipatory nausea and/or vomiting. 132-137 Systematic desensitization may also be helpful. 133 Hypnosis with guided imagery is another behavioral technique that has shown some success in treating this condition. 134

The antianxiety agents lorazepam and alprazolam have been combined with antiemetics for anticipatory nausea and/or vomiting. 131, 138, 139 The usual starting dose of alprazolam for anxiety is 0.25 to 0.5 mg orally 3 times daily, beginning on the night before treatment. In elderly patients, patients with debilitating disease, and patients with advanced liver disease, the usual starting dose of alprazolam is 0.25 mg orally 2 or 3 times daily for treatment of anxiety. 140 This dose may be gradually

increased if needed. Note that the elderly are especially sensitive to the effects of benzodiazepines. The dose should be gradually reduced when decreasing or discontinuing alprazolam therapy.

Managing Multiday Emetogenic Chemotherapy Regimens

Patients receiving multiple day (i.e., multiday) chemotherapy are at risk for both acute and delayed nausea and/or vomiting based on the emetogenic potential of the individual chemotherapy agents and their sequence. 23, 141-145 It is difficult to recommend a specific antiemetic regimen for each day, especially because acute and delayed emesis may overlap after the initial day of chemotherapy until the last day of chemotherapy. The period of risk for delayed emesis following completion of chemotherapy also depends on the specific regimen and the emetogenic potential of the last chemotherapy agent administered in the regimen. For multi-drug regimens, antiemetic therapy should be selected based on the drug with the highest emetic risk. General principles for managing multiday emetogenic chemotherapy regimens recommended by the panel are described in the algorithm (see Guidelines section on "Principles for Managing Multiday Emetogenic Chemotherapy Regimens").

For antiemetic prophylaxis of multiday emetogenic chemotherapy regimens (e.g., cisplatin-containing regimens), the combination of a 5-HT3 antagonist with dexamethasone remains the standard treatment.8, ^{23, 120} Dexamethasone should be administered once daily either orally or intravenously for every day of moderately or highly emetogenic chemotherapy and for 2-3 days after chemotherapy for regimens that are likely to cause significant delayed emesis. However, dexamethasone should not be added when the chemotherapy regimen already includes a corticosteroid. Steroids should also be avoided when



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

using regimens containing interleukin-2 (IL-2, aldesleukin) and interferon. 146

A 5-HT3 receptor antagonist should be administered each day before the first dose of moderately or highly emetogenic chemotherapy. Intravenous palonosetron may be used before the start of a 3-day chemotherapy regimen instead of multiple daily doses of oral or intravenous 5-HT3 receptor antagonists. 147, 148 Repeat dosing of palonosetron (0.25 mg IV) is likely to be safe, based on the dose ranging phase II trial and the 3 phase III trials using palonosetron as a single fixed dose (0.75 mg IV). 51-53, 149 Compared to the approved dose of palonosetron of 0.25 mg IV, these higher doses were not associated with significantly different adverse events.

The need for repeat dosing with palonosetron, either daily or less frequently, in the setting of multiday chemotherapy is not yet known. In one study, patients receiving highly emetogenic multiday cisplatinbased chemotherapy for testicular cancer (N=41) received multiday dosing of palonosetron (0.25 mg IV on days 1, 3, and 5) and dexamethasone, which prevented nausea and emesis in most patients on days 1-5 (51%) and on days 6-9 (83%); the most common adverse events were mild headache and constipation. ¹⁵⁰ A recent study assessed palonosetron given for 1, 2, or 3 days in combination with dexamethasone for patients receiving multiday high-dose chemotherapy prior to stem cell transplantation for multiple myeloma (N=73); during the 7-day emesis prevention period, about 40-45% of patients had no emesis (with no differences observed between palonosetron treatment groups), and no serious adverse events were reported. However, even among the patients who received either 2 or 3 days of palonosetron, only 20% had a complete response (i.e., emesis free without rescue medication). 151 Another study found that a palonosetron/dexamethasone regimen appeared to be more effective

for multiday chemotherapy than an ondansetron/dexamethasone regimen; patients received a second dose of palonosetron for breakthrough emesis, which was effective in 67% of patients who experienced nausea or vomiting. 147 Further studies are needed to define whether a need exists for repeat dosing of palonosetron in the setting of multiday chemotherapy.

The potential role of NK-1 antagonists in the antiemetic management of multiday chemotherapy regimens remains to be defined. In one study, the addition of the NK-1 antagonist aprepitant to granisetron and dexamethasone was evaluated in patients receiving multiday high and moderate emetogenic chemotherapy (N=78); in this study, the 3-drug antiemetic regimen was given during chemotherapy, and aprepitant and dexamethasone was given for an additional 2 days following chemotherapy. 152 Complete response (during the time period from day 1 until 5 days after chemotherapy) was observed in 58% and 73% of patients who received high and moderate emetogenic regimens, respectively. 152 Aprepitant may be used for multiday chemotherapy regimens likely to be highly emetogenic and associated with significant risk for delayed nausea and emesis. As per the labeled indication, aprepitant should be administered 125 mg orally 1 hour prior to chemotherapy on day 1, along with a 5-HT3 receptor antagonist and dexamethasone. Aprepitant 80 mg should be administered daily on days 2 and 3 after the start of chemotherapy along with dexamethasone. 141 Repeated dosing of aprepitant over multiple cycles of cisplatin-based chemotherapy was shown to be feasible and well tolerated; importantly, protection from emesis and from significant nausea was maintained during the subsequent cycles of emetogenic chemotherapy. 86, 141 Based on phase II data, aprepitant 80 mg may be safely administered on days 4 and 5 after chemotherapy.⁸⁷ It is not yet known if dosing aprepitant after day 3 improves control of nausea or



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

emesis in this clinical setting. As previously noted, fosaprepitant dimeglumine (115 mg) may be substituted for aprepitant (125 mg) on day 1 only. Alternatively, for highly emetogenic chemotherapy regimens, fosaprepitant 150 mg IV with dexamethasone may be given on day 1 with no need for oral aprepitant on days 2 and 3 with recommended dosing of dexamethasone.



NCCN Guidelines Index Antiemesis Table of Contents Discussion

References

- 1. Laszlo J. Emesis as limiting toxicity in cancer chemotherapy. In: Laszlo J, ed. Antiemetics and Cancer Chemotherapy. Baltimore, Maryland: Williams & Wilkins; 1983:1-5.
- 2. Ingle RJ, Burish TG, Wallston KA. Conditionability of cancer chemotherapy patients. Oncol Nurs Forum 1984;11:97-9102. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6566342.
- 3. Mitchell EP. Gastrointestinal toxicity of chemotherapeutic agents. Semin Oncol 1992;19:566-579. Available at: http://www.ncbi.nlm.nih.gov/pubmed/1411654.
- 4. Richardson JL, Marks G, Levine A. The influence of symptoms of disease and side effects of treatment on compliance with cancer therapy. J Clin Oncol 1988;6:1746-1752. Available at: http://www.ncbi.nlm.nih.gov/pubmed/3183704.
- 5. Herrstedt J. Antiemetics: an update and the MASCC guidelines applied in clinical practice. Nat Clin Pract Oncol 2008;5:32-43. Available at: http://www.ncbi.nlm.nih.gov/pubmed/18097455.
- 6. Hesketh PJ, Grunberg SM, Herrstedt J, et al. Combined data from two phase III trials of the NK1 antagonist aprepitant plus a 5HT 3 antagonist and a corticosteroid for prevention of chemotherapy-induced nausea and vomiting: effect of gender on treatment response. Support Care Cancer 2006;14:354-360. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16450086.
- 7. Hesketh PJ, Kris MG, Grunberg SM, et al. Proposal for classifying the acute emetogenicity of cancer chemotherapy. J Clin Oncol 1997:15:103-109. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8996130.
- 8. Roila F, Herrstedt J, Aapro M, et al. Guideline update for MASCC and ESMO in the prevention of chemotherapy- and radiotherapyinduced nausea and vomiting: results of the Perugia consensus

- conference. Ann Oncol 2010;21 Suppl 5:v232-243. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20555089.
- 9. Grunberg SM, Warr D, Gralla RJ, et al. Evaluation of new antiemetic agents and definition of antineoplastic agent emetogenicity-state of the art. Support Care Cancer 2010. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20972805.
- 10. Roscoe JA, Morrow GR, Colagiuri B, et al. Insight in the prediction of chemotherapy-induced nausea. Support Care Cancer 2010;18:869-876. Available at: http://www.ncbi.nlm.nih.gov/pubmed/19701781.
- 11. Craig JB, Powell BL. The management of nausea and vomiting in clinical oncology. Am J Med Sci 1987;293:34-44. Available at: http://www.ncbi.nlm.nih.gov/pubmed/3544842.
- 12. Borison HL, Wang SC. Physiology and pharmacology of vomiting. Pharmacol Rev 1953;5:193-230. Available at: http://www.ncbi.nlm.nih.gov/pubmed/13064033.
- 13. Seigel LJ, Longo DL. The control of chemotherapy-induced emesis. Ann Intern Med 1981;95:352-359. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7023313.
- 14. Dodds LJ. The control of cancer chemotherapy-induced nausea and vomiting. J Clin Hosp Pharm 1985;10:143-166. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2862166.
- 15. Aapro MS. Palonosetron as an anti-emetic and anti-nausea agent in oncology. Ther Clin Risk Manag 2007;3:1009-1020. Available at: http://www.ncbi.nlm.nih.gov/pubmed/18516316.
- 16. Grunberg SM, Deuson RR, Mavros P, et al. Incidence of chemotherapy-induced nausea and emesis after modern antiemetics. Cancer 2004:100:2261-2268. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15139073.
- 17. Hickok JT, Roscoe JA, Morrow GR, et al. 5-Hydroxytryptaminereceptor antagonists versus prochlorperazine for control of delayed



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

nausea caused by doxorubicin: a URCC CCOP randomised controlled trial. Lancet Oncol 2005:6:765-772. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16198982.

- 18. Kris MG, Gralla RJ, Clark RA, et al. Incidence, course, and severity of delayed nausea and vomiting following the administration of highdose cisplatin. J Clin Oncol 1985;3:1379-1384. Available at: http://www.ncbi.nlm.nih.gov/pubmed/4045527.
- 19. Roila F, Boschetti E, Tonato M, et al. Predictive factors of delayed emesis in cisplatin-treated patients and antiemetic activity and tolerability of metoclopramide or dexamethasone. A randomized singleblind study. Am J Clin Oncol 1991;14:238-242. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2031511.
- 20. Moher D, Arthur AZ, Pater JL. Anticipatory nausea and/or vomiting. Cancer Treat Rev 1984;11:257-264. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6151870.
- 21. Jacobsen PB, Redd WH. The development and management of chemotherapy-related anticipatory nausea and vomiting. Cancer Invest 1988;6:329-336. Available at: http://www.ncbi.nlm.nih.gov/pubmed/3048577.
- 22. Morrow GR. Clinical characteristics associated with the development of anticipatory nausea and vomiting in cancer patients undergoing chemotherapy treatment. J Clin Oncol 1984;2:1170-1176. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6491699.
- 23. Roila F, Hesketh PJ, Herrstedt J. Prevention of chemotherapy- and radiotherapy-induced emesis: results of the 2004 Perugia International Antiemetic Consensus Conference, Ann Oncol 2006:17:20-28. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16314401.
- 24. Feyer PC, Maranzano E, Molassiotis A, et al. Radiotherapy-induced nausea and vomiting (RINV): MASCC/ESMO guideline for antiemetics in radiotherapy: update 2009. Support Care Cancer 2010. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20697746.

- 25. Harding R, Young R, Anno G. Radiotherapy-induced emesis. In: Andrews P, Sanger G, eds. Emesis in Anti-Cancer Therapy. London: Chapman & Hall Medical; 1993:163-178.
- 26. Kris MG, Hesketh PJ, Somerfield MR, et al. American Society of Clinical Oncology guideline for antiemetics in oncology: update 2006. J Clin Oncol 2006;24:2932-2947. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16717289.
- 27. Laszlo J. Treatment of nausea and vomiting caused by cancer chemotherapy. Cancer Treat Rev 1982;9 Suppl B:3-9. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6299552.
- 28. Strum SB, McDermed JE, Pileggi J, et al. Intravenous metoclopramide: prevention of chemotherapy-induced nausea and vomiting. A preliminary evaluation. Cancer 1984;53:1432-1439. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6692333.
- 29. Lindley CM, Bernard S, Fields SM. Incidence and duration of chemotherapy-induced nausea and vomiting in the outpatient oncology population. J Clin Oncol 1989;7:1142-1149. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2787840.
- 30. Aapro M. Methodological issues in antiemetic studies. Invest New Drugs 1993;11:243-253. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8157467.
- 31. Grunberg SM, Osoba D, Hesketh PJ, et al. Evaluation of new antiemetic agents and definition of antineoplastic agent emetogenicity-an update. Support Care Cancer 2005;13:80-84. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15599601.
- 32. Koeller JM, Aapro MS, Gralla RJ, et al. Antiemetic guidelines: creating a more practical treatment approach. Support Care Cancer 2002;10:519-522. Available at: http://www.ncbi.nlm.nih.gov/pubmed/12324805.



NCCN Guidelines Index Antiemesis Table of Contents Discussion

- 33. Hesketh PJ, Gandara DR. Serotonin antagonists: a new class of antiemetic agents. J Natl Cancer Inst 1991;83:613-620. Available at: http://www.ncbi.nlm.nih.gov/pubmed/1850806.
- 34. Andrews PL, Bhandari P, Davey PT, et al. Are all 5-HT3 receptor antagonists the same? Eur J Cancer 1992;28A Suppl 1:2-6. Available at: http://www.ncbi.nlm.nih.gov/pubmed/1320915.
- 35. Grunberg SM, Koeller JM. Palonosetron: a unique 5-HT3-receptor antagonist for the prevention of chemotherapy-induced emesis. Expert Opin Pharmacother 2003;4:2297-2303. Available at: http://www.ncbi.nlm.nih.gov/pubmed/14640928.
- 36. Grunberg SM, Stevenson LL, Russell CA, McDermed JE. Dose ranging phase I study of the serotonin antagonist GR38032F for prevention of cisplatin-induced nausea and vomiting. J Clin Oncol 1989;7:1137-1141. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2526864.
- 37. Chevallier B. Efficacy and safety of granisetron compared with highdose metoclopramide plus dexamethasone in patients receiving highdose cisplatin in a single-blind study. The Granisetron Study Group. Eur J Cancer 1990;26 Suppl 1:S33-36. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2169784.
- 38. Cupissol DR, Serrou B, Caubel M. The efficacy of granisetron as a prophylactic anti-emetic and intervention agent in high-dose cisplatininduced emesis. Eur J Cancer 1990:26 Suppl 1:23-27. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2169782.
- 39. De Mulder PH, Seynaeve C, Vermorken JB, et al. Ondansetron compared with high-dose metoclopramide in prophylaxis of acute and delayed cisplatin-induced nausea and vomiting. A multicenter, randomized, double-blind, crossover study. Ann Intern Med 1990;113:834-840. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2146911.

- 40. Marty M. A comparative study of the use of granisetron, a selective 5-HT3 antagonist, versus a standard anti-emetic regimen of chlorpromazine plus dexamethasone in the treatment of cytostaticinduced emesis. The Granisetron Study Group. Eur J Cancer 1990;26 Suppl 1:S28-32. Available at:
- http://www.ncbi.nlm.nih.gov/pubmed/2169783.
- 41. Marty M, Pouillart P, Scholl S, et al. Comparison of the 5hydroxytryptamine3 (serotonin) antagonist ondansetron (GR 38032F) with high-dose metoclopramide in the control of cisplatin-induced emesis. N Engl J Med 1990;322:816-821. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2137902.
- 42. Soukop M. A comparison of two dose levels of granisetron in patients receiving high-dose cisplatin. The Granisetron Study Group. Eur J Cancer 1990;26 Suppl 1:S15-19. Available at: http://www.ncbi.nlm.nih.gov/pubmed/2169779.
- 43. Roila F, Tonato M, Cognetti F, et al. Prevention of cisplatin-induced emesis: a double-blind multicenter randomized crossover study comparing ondansetron and ondansetron plus dexamethasone. J Clin Oncol 1991:9:675-678. Available at: http://www.ncbi.nlm.nih.gov/pubmed/1829757.
- 44. Fraschini G. Antiemetic activity of ondansetron in cancer patients receiving non-cisplatin chemotherapy. Semin Oncol 1992;19:41-47. Available at: http://www.ncbi.nlm.nih.gov/pubmed/1387249.
- 45. Kamanabrou D. Intravenous granisetron--establishing the optimal dose. The Granisetron Study Group. Eur J Cancer 1992;28A Suppl 1:6-11. Available at: http://www.ncbi.nlm.nih.gov/pubmed/1320919.
- 46. Sledge GW, Einhorn L, Nagy C, House K. Phase III double-blind comparison of intravenous ondansetron and metoclopramide as antiemetic therapy for patients receiving multiple-day cisplatin-based chemotherapy. Cancer 1992;70:2524-2528. Available at: http://www.ncbi.nlm.nih.gov/pubmed/1423181.



NCCN Guidelines Index Antiemesis Table of Contents Discussion

- 47. Chevallier B. The control of acute cisplatin-induced emesis--a comparative study of granisetron and a combination regimen of highdose metoclopramide and dexamethasone. Granisetron Study Group. Br J Cancer 1993:68:176-180. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8391304.
- 48. Navari RM, Kaplan HG, Gralla RJ, et al. Efficacy and safety of granisetron, a selective 5-hydroxytryptamine-3 receptor antagonist, in the prevention of nausea and vomiting induced by high-dose cisplatin. J Clin Oncol 1994;12:2204-2210. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7931490.
- 49. Riviere A. Dose finding study of granisetron in patients receiving high-dose cisplatin chemotherapy. The Granisetron Study Group. Br J Cancer 1994;69:967-971. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8180032.
- 50. Rojas C, Thomas AG, Alt J, et al. Palonosetron triggers 5-HT(3) receptor internalization and causes prolonged inhibition of receptor function. Eur J Pharmacol 2010;626;193-199. Available at: http://www.ncbi.nlm.nih.gov/pubmed/19836386.
- 51. Aapro MS, Grunberg SM, Manikhas GM, et al. A phase III, doubleblind, randomized trial of palonosetron compared with ondansetron in preventing chemotherapy-induced nausea and vomiting following highly emetogenic chemotherapy. Ann Oncol 2006;17:1441-1449. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16766588.
- 52. Eisenberg P. Figueroa-Vadillo J. Zamora R. et al. Improved prevention of moderately emetogenic chemotherapy-induced nausea and vomiting with palonosetron, a pharmacologically novel 5-HT3 receptor antagonist: results of a phase III, single-dose trial versus dolasetron. Cancer 2003;98:2473-2482. Available at: http://www.ncbi.nlm.nih.gov/pubmed/14635083.
- 53. Gralla R, Lichinitser M, Van Der Vegt S, et al. Palonosetron improves prevention of chemotherapy-induced nausea and vomiting following moderately emetogenic chemotherapy: results of a double-

blind randomized phase III trial comparing single doses of palonosetron with ondansetron. Ann Oncol 2003;14:1570-1577. Available at: http://www.ncbi.nlm.nih.gov/pubmed/14504060.

- 54. Saito M, Aogi K, Sekine I, et al. Palonosetron plus dexamethasone versus granisetron plus dexamethasone for prevention of nausea and vomiting during chemotherapy: a double-blind, double-dummy, randomised, comparative phase III trial. Lancet Oncol 2009;10:115-124. Available at: http://www.ncbi.nlm.nih.gov/pubmed/19135415.
- 55. Yang LP, Scott LJ. Palonosetron: in the prevention of nausea and vomiting. Drugs 2009;69:2257-2278. Available at: http://www.ncbi.nlm.nih.gov/pubmed/19852528.
- 56. United States Food and Drug Administration. Anzemet (dolasetron mesylate): Drug Safety Communication - Reports of Abnormal Heart Rhythms. 2010. Available at: http://www.fda.gov/Drugs/DrugSafety/ucm237081.htm. Accessed June 2011.
- 57. ProStrakan Inc. Prescribing Information. Sancuso (Granisetron Transdermal System). 2008. Available at: http://www.sancuso.com/forms/SANCUSO-Full Pl.pdf. Accessed June 2011.
- 58. Boccia RV, Gordan LN, Clark G, et al. Efficacy and tolerability of transdermal granisetron for the control of chemotherapy-induced nausea and vomiting associated with moderately and highly emetogenic multi-day chemotherapy: a randomized, double-blind, phase III study. Support Care Cancer 2010. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20835873.
- 59. Howell J, Smeets J, Drenth HJ, Gill D. Pharmacokinetics of a granisetron transdermal system for the treatment of chemotherapyinduced nausea and vomiting. J Oncol Pharm Pract 2009;15:223-231. Available at: http://www.ncbi.nlm.nih.gov/pubmed/19304880.



NCCN Guidelines Index Antiemesis Table of Contents Discussion

- 60. Bonneterre J, Hecquet B. Granisetron (IV) compared with ondansetron (IV plus oral) in the prevention of nausea and vomiting induced by moderately-emetogenic chemotherapy. A cross-over study. Bull Cancer 1995:82:1038-1043. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8745670.
- 61. Jantunen IT, Muhonen TT, Kataja VV, et al. 5-HT3 receptor antagonists in the prophylaxis of acute vomiting induced by moderately emetogenic chemotherapy--a randomised study. Eur J Cancer 1993:29A:1669-1672. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8398291.
- 62. Martoni A, Angelelli B, Guaraldi M, et al. An open randomised cross-over study on granisetron versus ondansetron in the prevention of acute emesis induced by moderate dose cisplatin-containing regimens. Eur J Cancer 1996;32A:82-85. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8695248.
- 63. Gebbia V, Cannata G, Testa A, et al. Ondansetron versus granisetron in the prevention of chemotherapy-induced nausea and vomiting. Results of a prospective randomized trial. Cancer 1994:74:1945-1952. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8082100.
- 64. Mantovani G, Maccio A, Bianchi A, et al. Comparison of granisetron, ondansetron, and tropisetron in the prophylaxis of acute nausea and vomiting induced by cisplatin for the treatment of head and neck cancer: a randomized controlled trial. Cancer 1996:77:941-948. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8608488.
- 65. Noble A, Bremer K, Goedhals L, et al. A double-blind, randomised, crossover comparison of granisetron and ondansetron in 5-day fractionated chemotherapy: assessment of efficacy, safety and patient preference. The Granisetron Study Group. Eur J Cancer 1994:30A:1083-1088. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7654434.

- 66. Ruff P. Paska W. Goedhals L. et al. Ondansetron compared with granisetron in the prophylaxis of cisplatin-induced acute emesis: a multicentre double-blind, randomised, parallel-group study. The Ondansetron and Granisetron Emesis Study Group. Oncology 1994;51:113-118. Available at:
- http://www.ncbi.nlm.nih.gov/pubmed/8265095.
- 67. Ondansetron versus granisetron, both combined with dexamethasone, in the prevention of cisplatin-induced emesis. Italian Group of Antiemetic Research. Ann Oncol 1995;6:805-810. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8589019.
- 68. Navari R, Gandara D, Hesketh P, et al. Comparative clinical trial of granisetron and ondansetron in the prophylaxis of cisplatin-induced emesis. The Granisetron Study Group. J Clin Oncol 1995;13:1242-1248. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7738628.
- 69. Stewart A, McQuade B, Cronje JD, et al. Ondansetron compared with granisetron in the prophylaxis of cyclophosphamide-induced emesis in out-patients: a multicentre, double-blind, double-dummy, randomised, parallel-group study. Emesis Study Group for Ondansetron and Granisetron in Breast Cancer Patients. Oncology 1995:52:202-210. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7715904.
- 70. Audhuy B, Cappelaere P, Martin M, et al. A double-blind, randomised comparison of the anti-emetic efficacy of two intravenous doses of dolasetron mesilate and granisetron in patients receiving high dose cisplatin chemotherapy. Eur J Cancer 1996;32A:807-813. Available at: http://www.ncbi.nlm.nih.gov/pubmed/9081358.
- 71. Fauser AA, Duclos B, Chemaissani A, et al. Therapeutic equivalence of single oral doses of dolasetron mesilate and multiple doses of ondansetron for the prevention of emesis after moderately emetogenic chemotherapy. European Dolasetron Comparative Study Group. Eur J Cancer 1996;32A:1523-1529. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8911112.



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

- 72. Hesketh P, Navari R, Grote T, et al. Double-blind, randomized comparison of the antiemetic efficacy of intravenous dolasetron mesylate and intravenous ondansetron in the prevention of acute cisplatin-induced emesis in patients with cancer. Dolasetron Comparative Chemotherapy-induced Emesis Prevention Group. J Clin Oncol 1996;14:2242-2249. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8708713.
- 73. Lofters WS, Pater JL, Zee B, et al. Phase III double-blind comparison of dolasetron mesylate and ondansetron and an evaluation of the additive role of dexamethasone in the prevention of acute and delayed nausea and vomiting due to moderately emetogenic chemotherapy. J Clin Oncol 1997;15:2966-2973. Available at: http://www.ncbi.nlm.nih.gov/pubmed/9256141.
- 74. Jordan K, Hinke A, Grothey A, et al. A meta-analysis comparing the efficacy of four 5-HT3-receptor antagonists for acute chemotherapyinduced emesis. Support Care Cancer 2007;15:1023-1033. Available at: http://www.ncbi.nlm.nih.gov/pubmed/17205281.
- 75. Billio A, Morello E, Clarke MJ. Serotonin receptor antagonists for highly emetogenic chemotherapy in adults. Cochrane Database Syst Rev 2010:CD006272. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20091591.
- 76. Botrel TE, Clark OA, Clark L, et al. Efficacy of palonosetron (PAL) compared to other serotonin inhibitors (5-HT(3)R) in preventing chemotherapy-induced nausea and vomiting (CINV) in patients receiving moderately or highly emetogenic (MoHE) treatment: systematic review and meta-analysis. Support Care Cancer 2011;19:823-832. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20495832.
- 77. Aapro M, Fabi A, Nole F, et al. Double-blind, randomised, controlled study of the efficacy and tolerability of palonosetron plus dexamethasone for 1 day with or without dexamethasone on days 2 and 3 in the prevention of nausea and vomiting induced by moderately

- emetogenic chemotherapy. Ann Oncol 2010;21:1083-1088. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20080830.
- 78. Huang JQ, Zheng GF, Deuson R, et al. Do 5-hydroxytryptamine3 receptor antagonists (5-HT3) improve the antiemetic effect of dexamethasone for preventing delayed chemotherapy-induced nausea and vomiting (CINV)? A meta-analysis of randomized controlled trials [abstract]. J Clin Oncol 2004;22(Suppl 14): Abstract 6037. Available at: http://meeting.ascopubs.org/cgi/content/abstract/22/14 suppl/6037.
- 79. Gralla RJ, de Wit R, Herrstedt J, et al. Antiemetic efficacy of the neurokinin-1 antagonist, aprepitant, plus a 5HT3 antagonist and a corticosteroid in patients receiving anthracyclines or cyclophosphamide in addition to high-dose cisplatin: analysis of combined data from two Phase III randomized clinical trials. Cancer 2005;104:864-868. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15973669.
- 80. Hesketh PJ, Grunberg SM, Gralla RJ, et al. The oral neurokinin-1 antagonist aprepitant for the prevention of chemotherapy-induced nausea and vomiting: a multinational, randomized, double-blind, placebo-controlled trial in patients receiving high-dose cisplatin--the Aprepitant Protocol 052 Study Group. J Clin Oncol 2003;21:4112-4119. Available at: http://www.ncbi.nlm.nih.gov/pubmed/14559886.
- 81. Poli-Bigelli S, Rodrigues-Pereira J, Carides AD, et al. Addition of the neurokinin 1 receptor antagonist aprepitant to standard antiemetic therapy improves control of chemotherapy-induced nausea and vomiting. Results from a randomized, double-blind, placebo-controlled trial in Latin America, Cancer 2003:97:3090-3098, Available at: http://www.ncbi.nlm.nih.gov/pubmed/12784346.
- 82. Tremont-Lukats IW, Gonzalez-Barboteo J, Bruera E, Brescia FJ. Meta-analysis of neurokinin-1 receptor antagonists (NK-1 RA) for chemotherapy-induced nausea and vomiting (CINV) [abstract]. J Clin Oncol 2004;22(Suppl 14):Abstract 8047. Available at: http://meeting.ascopubs.org/cgi/content/abstract/22/14 suppl/8047.



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

- 83. Warr DG, Hesketh PJ, Gralla RJ, et al. Efficacy and tolerability of aprepitant for the prevention of chemotherapy-induced nausea and vomiting in patients with breast cancer after moderately emetogenic chemotherapy. J Clin Oncol 2005;23:2822-2830. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15837996.
- 84. Grote T, Hajdenberg J, Cartmell A, et al. Combination therapy for chemotherapy-induced nausea and vomiting in patients receiving moderately emetogenic chemotherapy: palonosetron, dexamethasone, and aprepitant. J Support Oncol 2006;4:403-408. Available at: http://www.ncbi.nlm.nih.gov/pubmed/17004515.
- 85. Grunberg SM, Dugan M, Muss H, et al. Effectiveness of a singleday three-drug regimen of dexamethasone, palonosetron, and aprepitant for the prevention of acute and delayed nausea and vomiting caused by moderately emetogenic chemotherapy. Support Care Cancer 2009:17:589-594. Available at: http://www.ncbi.nlm.nih.gov/pubmed/19037667.
- 86. Merck & Co., Inc. . Prescribing Information. EMEND (aprepitant) Capsules. 2011. Available at: http://www.merck.com/product/usa/pi circulars/e/emend/emend pi.pdf. Accessed June 2011.
- 87. Chawla SP, Grunberg SM, Gralla RJ, et al. Establishing the dose of the oral NK1 antagonist aprepitant for the prevention of chemotherapyinduced nausea and vomiting. Cancer 2003;97:2290-2300. Available at: http://www.ncbi.nlm.nih.gov/pubmed/12712486.
- 88. Merck & Co., Inc. Prescribing Information. EMEND (fosaprepitant dimeglumine) for Injection, for intravenous use. 2011. Available at: http://www.merck.com/product/usa/pi_circulars/e/emend iv/emend iv pi.pdf. Accessed June 2011.
- 89. Marbury TC, Ngo PL, Shadle CR, et al. Pharmacokinetics of Oral Dexamethasone and Midazolam When Administered With Single-Dose Intravenous 150 mg Fosaprepitant in Healthy Adult Subjects. J Clin

Pharmacol 2011. Available at: http://www.ncbi.nlm.nih.gov/pubmed/21209230.

90. Lasseter KC, Gambale J, Jin B, et al. Tolerability of fosaprepitant and bioequivalency to aprepitant in healthy subjects. J Clin Pharmacol 2007;47:834-840. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/17525168.

- 91. Grunberg S, Chua D, Maru A, et al. Single-dose fosaprepitant for the prevention of chemotherapy-induced nausea and vomiting associated with cisplatin therapy: randomized, double-blind study protocol--EASE. J Clin Oncol 2011;29:1495-1501. Available at: http://www.ncbi.nlm.nih.gov/pubmed/21383291.
- 92. Shadle CR, Lee Y, Majumdar AK, et al. Evaluation of potential inductive effects of aprepitant on cytochrome P450 3A4 and 2C9 activity. J Clin Pharmacol 2004;44:215-223. Available at: http://www.ncbi.nlm.nih.gov/pubmed/14973304.
- 93. Wampler G. The pharmacology and clinical effectiveness of phenothiazines and related drugs for managing chemotherapy-induced emesis. Drugs 1983;25 Suppl 1:35-51. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6840019.
- 94. Gralla RJ, Itri LM, Pisko SE, et al. Antiemetic efficacy of high-dose metoclopramide: randomized trials with placebo and prochlorperazine in patients with chemotherapy-induced nausea and vomiting. N Engl J Med 1981:305:905-909. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7024807.
- 95. Gralla RJ, Tyson LB, Bordin LA, et al. Antiemetic therapy: a review of recent studies and a report of a random assignment trial comparing metoclopramide with delta-9-tetrahydrocannabinol. Cancer Treat Rep 1984;68:163-172. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/6318993.



NCCN Guidelines Index **Antiemesis Table of Contents** Discussion

- 96. Bakowski MT. Advances in anti-emetic therapy. Cancer Treat Rev 1984:11:237-256. Available at:
- http://www.ncbi.nlm.nih.gov/pubmed/6098368.
- 97. Grossman B, Lessin LS, Cohen P. Droperidol prevents nausea and vomiting from cis-platinum. N Engl J Med 1979;301:47. Available at: http://www.ncbi.nlm.nih.gov/pubmed/449910.
- 98. Aapro MS, Alberts DS. High-dose dexamethasone for prevention of cis-platin-induced vomiting. Cancer Chemother Pharmacol 1981;7:11-14. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6122510.
- 99. Aapro MS, Plezia PM, Alberts DS, et al. Double-blind crossover study of the antiemetic efficacy of high-dose dexamethasone versus high-dose metoclopramide. J Clin Oncol 1984;2:466-471. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6539363.
- 100. Cassileth PA, Lusk EJ, Torri S, et al. Antiemetic efficacy of dexamethasone therapy in patients receiving cancer chemotherapy. Arch Intern Med 1983:143:1347-1349. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6347109.
- 101. Kris MG, Gralla RJ, Clark RA, et al. Consecutive dose-finding trials adding lorazepam to the combination of metoclopramide plus dexamethasone: improved subjective effectiveness over the combination of diphenhydramine plus metoclopramide plus dexamethasone. Cancer Treat Rep 1985;69:1257-1262. Available at: http://www.ncbi.nlm.nih.gov/pubmed/3912039.
- 102. Kris MG, Gralla RJ, Clark RA, et al. Antiemetic control and prevention of side effects of anti-cancer therapy with lorazepam or diphenhydramine when used in combination with metoclopramide plus dexamethasone. A double-blind, randomized trial. Cancer 1987;60:2816-2822. Available at: http://www.ncbi.nlm.nih.gov/pubmed/3315176.
- 103. Herman TS, Einhorn LH, Jones SE, et al. Superiority of nabilone over prochlorperazine as an antiemetic in patients receiving cancer

- chemotherapy. N Engl J Med 1979;300:1295-1297. Available at: http://www.ncbi.nlm.nih.gov/pubmed/375088.
- 104. Steele N, Gralla RJ, Braun DW, Young CW. Double-blind comparison of the antiemetic effects of nabilone and prochlorperazine on chemotherapy-induced emesis. Cancer Treat Rep 1980;64:219-224. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6250699.
- 105. Navari RM, Einhorn LH, Passik SD, et al. A phase II trial of olanzapine for the prevention of chemotherapy-induced nausea and vomiting: a Hoosier Oncology Group study. Support Care Cancer 2005;13:529-534. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15700131.
- 106. Srivastava M, Brito-Dellan N, Davis MP, et al. Olanzapine as an antiemetic in refractory nausea and vomiting in advanced cancer. J Pain Symptom Manage 2003;25:578-582. Available at: http://www.ncbi.nlm.nih.gov/pubmed/12782438.
- 107. Passik SD, Navari RM, Jung S-H, et al. A phase I trial of olanzapine (Zyprexa) for the prevention of delayed emesis in cancer patients: a Hoosier Oncology Group study. Cancer Invest 2004;22:383-388. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15493359.
- 108. Passik SD, Kirsh KL, Theobald DE, et al. A retrospective chart review of the use of olanzapine for the prevention of delayed emesis in cancer patients. J Pain Symptom Manage 2003;25:485-488. Available at: http://www.ncbi.nlm.nih.gov/pubmed/12727048.
- 109. Passik SD, Lundberg J, Kirsh KL, et al. A pilot exploration of the antiemetic activity of olanzapine for the relief of nausea in patients with advanced cancer and pain. J Pain Symptom Manage 2002;23:526-532. Available at: http://www.ncbi.nlm.nih.gov/pubmed/12067777.
- 110. Eli Lilly and Company. Prescribing Information. ZYPREXA (olanzapine) 2010. Available at: http://pi.lilly.com/us/zyprexa-pi.pdf. Accessed June 2011.



NCCN Guidelines Index Antiemesis Table of Contents Discussion

- 111. Morita T, Tei Y, Shishido H, Inoue S. Olanzapine-induced delirium in a terminally ill cancer patient. J Pain Symptom Manage 2004;28:102-103. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15276190.
- 112. National Cancer Institute. Eating Hints: Before, During, and After Cancer Treatment. 2009. Available at: http://www.cancer.gov/cancertopics/coping/eatinghints/page1. Accessed June 2011.
- 113. Geling O, Eichler H-G. Should 5-hydroxytryptamine-3 receptor antagonists be administered beyond 24 hours after chemotherapy to prevent delayed emesis? Systematic re-evaluation of clinical evidence and drug cost implications. J Clin Oncol 2005;23:1289-1294. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15718327.
- 114. United States Food and Drug Administration FDA News Release: FDA Requires Boxed Warning and Risk Mitigation Strategy for Metoclopramide-Containing Drugs Agency warns against chronic use of these products to treat gastrointestinal disorders. 2009. Available at: http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/2009 /ucm149533.htm. Accessed June 2011.
- 115. Pasricha PJ, Pehlivanov N, Sugumar A, Jankovic J. Drug Insight: from disturbed motility to disordered movement--a review of the clinical benefits and medicolegal risks of metoclopramide. Nat Clin Pract Gastroenterol Hepatol 2006;3:138-148. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16511548.
- 116. Kawanishi C, Onishi H, Kato D, et al. Unexpectedly high prevalence of akathisia in cancer patients. Palliat Support Care 2007;5:351-354. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/18044412.

117. Patanwala AE, Amini R, Hays DP, Rosen P. Antiemetic therapy for nausea and vomiting in the emergency department. J Emerg Med 2010;39:330-336. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/20022195.

- 118. Vinson DR. Diphenhydramine in the treatment of akathisia induced by prochlorperazine. J Emerg Med 2004;26:265-270. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15028322.
- 119. Roila F, Warr D, Clark-Snow RA, et al. Delayed emesis: moderately emetogenic chemotherapy. Support Care Cancer 2005;13:104-108. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15549426.
- 120. Multinational Association of Supportive Care in Cancer. MASCC/ESMO Antiemetic Guideline 2011. 2011. Available at: http://data.memberclicks.com/site/mascc/MASCC Guidelines English 2011.pdf. Accessed June 2011.
- 121. Grunberg S, Clark-Snow RA, Koeller J. Chemotherapy-induced nausea and vomiting: contemporary approaches to optimal management: Proceedings from a symposium at the 2008 Multinational Association of Supportive Care in Cancer (MASCC) Annual Meeting. Support Care Cancer 2010. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20084406.
- 122. Ormrod D, Goa KL. Intranasal metoclopramide. Drugs 1999;58:315-322; discussion 323-314. Available at: http://www.ncbi.nlm.nih.gov/pubmed/10473023.
- 123. Urba S. Radiation-induced nausea and vomiting. J Natl Compr Canc Netw 2007:5:60-65. Available at: http://www.ncbi.nlm.nih.gov/pubmed/17239327.
- 124. Maranzano E, Feyer P, Molassiotis A, et al. Evidence-based recommendations for the use of antiemetics in radiotherapy. Radiother Oncol 2005:76:227-233. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16150504.
- 125. Salvo N, Doble B, Khan L, et al. Prophylaxis of radiation-induced nausea and vomiting using 5-hydroxytryptamine-3 serotonin receptor antagonists: a systematic review of randomized trials. Int J Radiat



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

Oncol Biol Phys 2010. Available at: http://www.ncbi.nlm.nih.gov/pubmed/21075553.

- 126. Franzen L, Nyman J, Hagberg H, et al. A randomised placebo controlled study with ondansetron in patients undergoing fractionated radiotherapy. Ann Oncol 1996;7:587-592. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8879372.
- 127. Wong RK, Paul N, Ding K, et al. 5-hydroxytryptamine-3 receptor antagonist with or without short-course dexamethasone in the prophylaxis of radiation induced emesis: a placebo-controlled randomized trial of the National Cancer Institute of Canada Clinical Trials Group (SC19). J Clin Oncol 2006;24:3458-3464. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16849762.
- 128. Lanciano R, Sherman DM, Michalski J, et al. The efficacy and safety of once-daily Kytril (granisetron hydrochloride) tablets in the prophylaxis of nausea and emesis following fractionated upper abdominal radiotherapy. Cancer Invest 2001;19:763-772. Available at: http://www.ncbi.nlm.nih.gov/pubmed/11768028.
- 129. Spitzer TR, Friedman CJ, Bushnell W, et al. Double-blind, randomized, parallel-group study on the efficacy and safety of oral granisetron and oral ondansetron in the prophylaxis of nausea and vomiting in patients receiving hyperfractionated total body irradiation. Bone Marrow Transplant 2000;26:203-210. Available at: http://www.ncbi.nlm.nih.gov/pubmed/10918432.
- 130. Roscoe JA, Morrow GR, Aapro MS, et al. Anticipatory nausea and vomiting. Support Care Cancer 2010. Available at: http://www.ncbi.nlm.nih.gov/pubmed/20803345.
- 131. Aapro MS, Molassiotis A, Olver I. Anticipatory nausea and vomiting. Support Care Cancer 2005;13:117-121. Available at: http://www.ncbi.nlm.nih.gov/pubmed/15599779.
- 132. Ezzo J, Vickers A, Richardson MA, et al. Acupuncture-point stimulation for chemotherapy-induced nausea and vomiting. J Clin

Oncol 2005;23:7188-7198. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16192603.

- 133. Morrow GR, Morrell C. Behavioral treatment for the anticipatory nausea and vomiting induced by cancer chemotherapy. N Engl J Med 1982;307:1476-1480. Available at: http://www.ncbi.nlm.nih.gov/pubmed/6128673.
- 134. Redd WH, Andrykowski MA. Behavioral intervention in cancer treatment: controlling aversion reactions to chemotherapy. J Consult Clin Psychol 1982;50:1018-1029. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7174969.
- 135. Ezzo J, Streitberger K, Schneider A. Cochrane systematic reviews examine P6 acupuncture-point stimulation for nausea and vomiting. J Altern Complement Med 2006;12:489-495. Available at: http://www.ncbi.nlm.nih.gov/pubmed/16813514.
- 136. Figueroa-Moseley C, Jean-Pierre P, Roscoe JA, et al. Behavioral interventions in treating anticipatory nausea and vomiting. J Natl Compr Canc Netw 2007;5:44-50. Available at: http://www.ncbi.nlm.nih.gov/pubmed/17239325.
- 137. Molassiotis A, Yung HP, Yam BM, et al. The effectiveness of progressive muscle relaxation training in managing chemotherapy-induced nausea and vomiting in Chinese breast cancer patients: a randomised controlled trial. Support Care Cancer 2002;10:237-246. Available at: http://www.ncbi.nlm.nih.gov/pubmed/11904789.
- 138. Razavi D, Delvaux N, Farvacques C, et al. Prevention of adjustment disorders and anticipatory nausea secondary to adjuvant chemotherapy: a double-blind, placebo-controlled study assessing the usefulness of alprazolam. J Clin Oncol 1993;11:1384-1390. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8315437.
- 139. Malik IA, Khan WA, Qazilbash M, et al. Clinical efficacy of lorazepam in prophylaxis of anticipatory, acute, and delayed nausea and vomiting induced by high doses of cisplatin. A prospective



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

randomized trial. Am J Clin Oncol 1995;18:170-175. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7900711.

140. Pfizer. Prescribing Information. XANAX® alprazolam tablets. 2011. Available at:

http://labeling.pfizer.com/ShowLabeling.aspx?id=547. Accessed June 2011.

- 141. de Wit R, Herrstedt J, Rapoport B, et al. The oral NK(1) antagonist, aprepitant, given with standard antiemetics provides protection against nausea and vomiting over multiple cycles of cisplatin-based chemotherapy: a combined analysis of two randomised, placebo-controlled phase III clinical trials. Eur J Cancer 2004;40:403-410. Available at: http://www.ncbi.nlm.nih.gov/pubmed/14746859.
- 142. Ellebaek E, Herrstedt J. Optimizing antiemetic therapy in multipleday and multiple cycles of chemotherapy. Curr Opin Support Palliat Care 2008;2:28-34. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/18685391.

143. Einhorn LH, Rapoport B, Koeller J, et al. Antiemetic therapy for multiple-day chemotherapy and high-dose chemotherapy with stem cell transplant: review and consensus statement. Support Care Cancer 2005;13:112-116. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/15480812.

- 144. Navari RM. Prevention of emesis from multiple-day and high-dose chemotherapy regimens. J Natl Compr Canc Netw 2007;5:51-59. Available at: http://www.ncbi.nlm.nih.gov/pubmed/17239326.
- 145. Herrstedt J, Sigsgaard TC, Nielsen HA, et al. Randomized, double-blind trial comparing the antiemetic effect of tropisetron plus metopimazine with tropisetron plus placebo in patients receiving multiple cycles of multiple-day cisplatin-based chemotherapy. Support Care Cancer 2007;15:417-426. Available at: http://www.ncbi.nlm.nih.gov/pubmed/17093916.

146. Mier JW, Vachino G, Klempner MS, et al. Inhibition of interleukin-2-induced tumor necrosis factor release by dexamethasone: prevention of an acquired neutrophil chemotaxis defect and differential suppression of interleukin-2-associated side effects. Blood 1990;76:1933-1940. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/2242421.

147. Musso M, Scalone R, Bonanno V, et al. Palonosetron (Aloxi) and dexamethasone for the prevention of acute and delayed nausea and vomiting in patients receiving multiple-day chemotherapy. Support Care Cancer 2009;17:205-209. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/18839220.

- 148. Celio L, Denaro A, Canova S, et al. Clinical update on palonosetron in the management of chemotherapy-induced nausea and vomiting. Tumori 2008;94:447-452. Available at: http://www.ncbi.nlm.nih.gov/pubmed/18822676.
- 149. Eisenberg P, MacKintosh FR, Ritch P, et al. Efficacy, safety and pharmacokinetics of palonosetron in patients receiving highly emetogenic cisplatin-based chemotherapy: a dose-ranging clinical study. Ann Oncol 2004;15:330-337. Available at: http://www.ncbi.nlm.nih.gov/pubmed/14760130.
- 150. Einhorn LH, Brames MJ, Dreicer R, et al. Palonosetron plus dexamethasone for prevention of chemotherapy-induced nausea and vomiting in patients receiving multiple-day cisplatin chemotherapy for germ cell cancer. Support Care Cancer 2007;15:1293-1300. Available at: http://www.ncbi.nlm.nih.gov/pubmed/17436025.
- 151. Giralt SA, Mangan KF, Maziarz RT, et al. Three palonosetron regimens to prevent CINV in myeloma patients receiving multiple-day high-dose melphalan and hematopoietic stem cell transplantation. Ann Oncol 2010. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/20935058.

152. Jordan K, Kinitz I, Voigt W, et al. Safety and efficacy of a triple antiemetic combination with the NK-1 antagonist aprepitant in highly



NCCN Guidelines Index
Antiemesis Table of Contents
Discussion

and moderately emetogenic multiple-day chemotherapy. Eur J Cancer 2009;45:1184-1187. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/19135359.