

General Topology Notes

Made by Leon :) *Note: Any reference numbers are to the lecture notes*

1 Topological Spaces and Examples

Definition 1.1: Topological Space

A **topological space** is a pair (X, \mathcal{T}) , where X is a nonempty set, and \mathcal{T} is a collection of subsets of X which satisfies:

- a) $\emptyset \in \mathcal{T}$ and $X \in \mathcal{T}$
- b) if $U_\lambda \in \mathcal{T}$ for each $\lambda \in \Lambda$ (where Λ is some indexing set), then $\bigcup_{\lambda \in \Lambda} U_\lambda \in \mathcal{T}$
- c) if $U_1, U_2 \in \mathcal{T}$, then $U_1 \cap U_2 \in \mathcal{T}$

The collection \mathcal{T} is called the **topology** of the topological space, and the members of \mathcal{T} are called the **open sets** of the topology

Example 1.7: Euclidean Spaces

Let \mathbb{R}^n denote the n -dimensional Euclidean vector space with elements $x = (x_1, x_2, \dots, x_n)$ and $x_i \in \mathbb{R}$, and let

$$|x| = \sqrt{\sum_{i=1}^n x_i^2} \geq 0$$

be the length of x . ($\mathbb{R}^1 = \mathbb{R}$ is the real line). A subset U of \mathbb{R}^n is **open (for the usual topology)** iff for each $a \in U$ there exists an $r > 0$ such that

$$|x - a| < r \implies x \in U.$$

The collection of open sets thus defined is called the **usual topology** on \mathbb{R}^n . Note that open balls $B(y, \rho) = \{x \in \mathbb{R}^n : |x - y| < \rho\}$ are open sets under this definition.

Example 1.8: Metric Spaces

A **metric space** (X, d) is a nonempty set X together with a function $d : X \times X \rightarrow \mathbb{R}$ with the following properties:

- a) $d(x, y) \geq 0$ and $d(x, y) = 0 \iff x = y$
- b) $d(x, y) = d(y, x)$
- c) $d(x, y) \leq d(x, z) + d(z, y)$ (Triangle Inequality)

The function d is called the **metric**.

Let (X, d) be a metric space, x be a point in X , and $r > 0$. The **open ball** with center x and radius r is defined by

$$B(x, r) = \{y, \in X : d(x, y) < r\}.$$

A subset U of X is **open (in the metric topology given by d)** iff for each $a \in U$ there is an $r > 0$ such that $B(a, r) \subseteq U$. Just like euclidean spaces, open balls are open in this sense.

Proposition 1.11: Topology Equality

Let d, d' be metrics on the same set X , and let $\mathcal{T}, \mathcal{T}'$ be the corresponding metric topologies. If for real numbers $A, B > 0$ we have

$$d(x, y) \leq Ad'(x, y), d'(x, y) \leq Bd(x, y) \text{ for all } x, y \in X,$$

then $\mathcal{T} = \mathcal{T}'$.

Definition 1.16: Subspace Topology

Let (X, \mathcal{T}) be a topological space, and let $A \subseteq X$ be any subset. Then the **subspace topology** on A consists of all sets of the form $U \cap A$ where $U \in \mathcal{T}$.

Definition 1.17: Closed Set

Let (X, \mathcal{T}) be a topological space. A subset $A \subseteq X$ is **closed** iff its complement $X \setminus A := \{x \in X \mid x \notin A\}$ is open in X . Note that a set being *closed* does not mean it isn't *open*. Sets that are both *closed* and *open* are called **clopen**.

Definition 1.20: Properties of Topological Spaces

For a subset $A \subseteq X$,

- The **closure** of A is $\bar{A} := \bigcap_{\substack{C \subseteq X \text{ closed;} \\ A \subseteq C}} C.$
- The **interior** of A is $\text{int } A = A^\circ := \bigcap_{\substack{C \subseteq X \text{ open;} \\ A \subseteq C}} C.$

- The **boundary** (or **frontier**) of A is $\partial A := \bar{A} \setminus A^\circ.$

- A is **dense** in X iff $\bar{A} = X$.
- A **limit point** of A is a point $x \in X$ s.t. for every open subset $U \subseteq X$ with $x \in U$ there exists an element $a \in A \cup U$ with $a \neq x$. Let A' be the set of limit points of A . Note that this has nothing to do with limits of sequences.

Proposition 1.22: Relating Topological Properties

- \bar{A} is closed, and contains A and is the smallest set with this property. So A is closed iff $\bar{A} = A$.
- A° is open, and is contained in A , and is the largest set with this property. So A is open iff $A^\circ = A$.
- The closure of the complement is the complement of the interior: $\overline{X \setminus A} = X \setminus (A^\circ).$
- The interior of the complement is the complement of the closure: $(X \setminus A)^\circ = X \setminus \bar{A}.$

Proposition 1.26: Union of Limit Points

Let (X, \mathcal{T}) be a topological space, and suppose $A \subseteq X$. Then

$$\bar{A} = A \cup A'$$

Corollary 1.27

A subset $A \subseteq X$ is closed iff it contains all its limit points.

Theorem 1.19: Properties of open and closed sets

Let (X, \mathcal{T}) be a topological space.

1. \emptyset and X are **closed**.
2. The union of **finitely many** closed sets is an closed set.
3. The intersection of **any collection** of closed sets is a closed set.

Lemma 1.24: Limit Points and Open Balls

An element $x \in X$ in a metric space (X, d) is a limit point of a subset $A \subseteq X$ iff for every $\epsilon > 0$ there exists $a \in A$ with $0 < d(x, a) < \epsilon$, or iff there exists a sequence a_1, a_2, a_3, \dots of elements $a_i \in A$, with $a_i \neq x$ for all i , s.t. $d(x_i, a_i) \rightarrow 0$ as $i \rightarrow \infty$. This interpretation does not extend to general topological spaces.

Theorem 1.30: Open and Closed sets in \mathbb{R}

Consider \mathbb{R} with the usual topology.

1. A nonempty set U is open iff it can be written as a countable union of disjoint nonempty open intervals I_j (shown left):
2. A set F is closed iff it can be written as a countable intersection where each F_j is a finite union of closed intervals (shown right).

$$U = \bigcup_{j=1}^{\infty} I_j, \quad F = \bigcap_{j=1}^{\infty} F_j.$$

Definition 1.32: Hausdorff Spaces

A topological space (X, \mathcal{T}) is **Hausdorff** if for each $x, y \in X$ with $x \neq y$ there exist **disjoint** open sets U and V such that $x \in U$ and $y \in V$.

Any metrisable space is Hausdorff, The trivial topology on a set with more than one element is not Hausdorff.

Definition 1.33: Convergence of a Topological space

A sequence (x_n) of members of a topological space X converges to $x \in X$ if for every open set U containing x , there exists an N such that $n \geq N \implies x_n \in U$

Proposition 1.34: Convergence of Hausdorff Spaces

Suppose (X, \mathcal{T}) is Hausdorff. Then a sequence (x_n) can converge to at most one limit.

Definition 1.36: Cauchy and Completeness

Let (X, d) be a metric space.

1. A **Cauchy sequence** is a sequence (x_n) with each $x_n \in X$ with the property that for each $\epsilon > 0$, there exists an N such that $m, n \geq N \implies d(x_m, x_n) < \epsilon$
2. (X, d) is **complete** if every Cauchy sequence converges.

Definition 1.37: Topology Basis

A **basis for a topology** on a set X is a collection \mathcal{B} of subsets $B \subseteq X$ such that:

1. $X = \bigcup_{B \in \mathcal{B}} B$
2. The intersection of sets $B_1, B_2 \in \mathcal{B}$ is a set $B_1 \cap B_2 \in \mathcal{B}$.

The **topology \mathcal{T} generated by a basis \mathcal{B}** has open sets the arbitrary unions of basis elements $B_\lambda \in \mathcal{B}$:

$$U = \bigcup_{\lambda \in \Lambda} B_\lambda$$

(Don't forget to check that this really is a topology)

Example 1.38: Finite Intersections of open balls

For any metric space (X, \mathcal{T}) the finite intersections of open balls

$$B(x, r) = \{y \in X \mid d(x, y) < r\} \subseteq X \quad (r > 0, x \in X)$$

constitute a basis for the metric topology on X

$$\mathcal{B} = \{B(x_1, r_1) \cap B(x_2, r_2) \cap \dots \cap B(x_k, r_k) \mid x_1, x_2, \dots, x_k \in X, r_1, r_2, \dots, r_k > 0\}$$

2 Continuous functions and Homeomorphisms

Definition 2.1: Continuity

Let (X, \mathcal{T}) , (Y, \mathcal{U}) be topological spaces. A function $f : X \rightarrow Y$ is **continuous** iff

$$U \in \mathcal{U} \text{ implies } f^{-1}(U) \in \mathcal{T}.$$

That is, **inverse** images of open sets are open. Continuous functions are often called **maps** or **mappings** of topological spaces.

Proposition 2.6: Topological and Analytic Continuity

Let (X, d) and (Y, ρ) be metric spaces with their induced topologies \mathcal{T} and \mathcal{U} respectively. A function $f : X \rightarrow Y$ is continuous (topologically) iff it is continuous analytically: for every $a \in X$ and every $\epsilon > 0$ there exists $\delta > 0$ such that

$$d(x, a) < \delta \implies \rho(f(x), f(a)) < \epsilon$$

Definition 2.7: Homeomorphism

Let (X, \mathcal{T}) and (Y, \mathcal{U}) be topological spaces. A **homeomorphism** is a bijection $f : X \rightarrow Y$ such that both f and f^{-1} are continuous. Two topological spaces are **homeomorphic** if there is a homeomorphism between them.

Proposition 2.18: The Punctured Sphere

Consider the n -dimensional sphere

$$\mathbb{S}^n = \{x \in \mathbb{R}^{n+1} : |x| = 1\}$$

with the metric topology inherited from \mathbb{R}^{n+1} . Let $x_0 \in \mathbb{S}^n$. Then $\mathbb{S}^n \setminus \{x_0\}$ is homeomorphic to \mathbb{R}^n .

3 Subspaces Revisited

Definition 3.65: Disjoint Unions

Let (X, \mathcal{T}) and (Y, \mathcal{U}) be topological spaces. Their **disjoint union** $X + Y$ is the set $(X \times \{0\}) \cup (Y \times \{1\})$ with the topology consisting of all sets of the form

$$(T \times \{0\}) \cup (U \times \{1\}) \text{ such that } T \in \mathcal{T}, U \in \mathcal{U}$$

Definition 3.8: Product Topology

Let (X, \mathcal{T}) , (Y, \mathcal{U}) be topological spaces. The **product topology** on their product $X \times Y$ consists of all sets of the form

$$T = \bigcup_{\alpha \in A} (U_\alpha \times V_\alpha)$$

where A is an arbitrary indexing set, and $U_\alpha \in \mathcal{U}$ and $V_\alpha \in \mathcal{V}$.

Lemma 3.10

The product topology is indeed a topology. (lol)

Lemma 3.9: Openness in Product Topologies

Let (X, \mathcal{T}) , (Y, \mathcal{U}) be topological spaces. Then $T \subseteq X \times Y$ is open in the product topology if and only if for all $t \in T$ there exists $U \in \mathcal{U}$ and $V \in \mathcal{V}$ such that $t \in U \times V$ and $U \times V \subseteq T$.

Definition 3.11.5: Projection Maps

Let (X, \mathcal{T}) and (Y, \mathcal{U}) be topological spaces, and consider their product $X \times Y$ with the product topology. There are two natural maps Π_X and Π_Y , the projections of $X \times Y$ onto X and Y respectively, given by

$$\Pi_X : X \times Y \rightarrow X, \quad (x, y) \mapsto x$$

$$\Pi_Y : X \times Y \rightarrow Y, \quad (x, y) \mapsto y.$$

Definition 3.14: Weak Topology

Suppose that X is a set. $(X_\lambda, \mathcal{T}_\lambda)$ is a family of topological spaces, and that $f_\lambda : X \rightarrow X_\lambda$ are functions. The **weak topology generated by $\{f_\lambda\}$** is the smallest topology on X making all the f_λ continuous.

Thus the product topology on $X \times Y$ is the weak topology generated by the two maps Π_X and Π_Y .

Definition 3.15: Cartesian Product Topology

If X_λ is a topological space, (with λ in some arbitrary indexing set Λ), the product topology on the cartesian product $\prod_{\lambda \in \Lambda} X_\lambda$ is defined to be the weak topology generated by the projections

$$\Pi_\lambda : \prod_{\mu \in \Lambda} X_\mu \rightarrow X_\lambda$$

Definition 3.5.0: Equivalence Relation

An **equivalence** relation on a set X is a binary operation \sim on X which is:

1. **Reflexive**: $x \sim x$ for all $x \in X$.
2. **Symmetric**: if $x \sim y$ then $y \sim x$.
3. **Transitive**: if $x \sim y$ and $y \sim z$ then $x \sim z$.

The **equivalence class** of any element $x \in X$ is the set

$$[x] = \{y \in X \mid x \sim y\},$$

and the set of equivalence classes is denoted by X/\sim . The function which assigns to each $x \in X$ the equivalence class $[x] \in X/\sim$ is a surjection

$$p : X \rightarrow X/\sim; \quad x \mapsto [x]$$

Definition 3.17: Quotient Space

Given a topological space (X, \mathcal{T}) , and an equivalence relation \sim on X , the **quotient space** or **identification space** is the set of equivalence classes X/\sim together with the topology

$$\{U \subseteq X/\sim : p^{-1}(U) \in \mathcal{T}\}$$

Definition 3.25: Generated Topological Spaces

Let X be a topological space, and let $Y_0, Y_1 \subseteq X$ be subspaces related by a continuous function $f : Y_0 \rightarrow Y_1$. Let \sim_f be the equivalence relation on X **generated by f** , the intersection of all the equivalence relations on X (regarded as subsets of $X \times X$) containing the pairs $(y_0, f(y_0))$ with $y_0 \in Y_0$. The identification space

$$X/\sim_f = X/\{y_1 = f(y_0)\}$$

is obtained by identifying each $y_0 \in Y_0 \subseteq X$ with $y_1 = f(y_0) \in Y_1 \subseteq X$.

Proposition 3.34: Homeomorphisms of Relations

Given a continuous function $f : X \rightarrow Y$ let \sim be the equivalence relation defined on X by $x \sim x'$ if $f(x) = f(x') \in Y$. The function

$$g : X/\sim \rightarrow Y; [x] \mapsto f(x)$$

is continuous and injective, with

$$f = g \circ p : X \xrightarrow{p} X/\sim \xrightarrow{g} Y.$$

If f is onto, and such that $f(U) \subseteq Y$ is open for every open subset $U \subseteq X$ then g is a homeomorphism.

4 Compact Spaces

Definition 4.1: Open Covers and Compact Spaces

1. An **open cover** of a topological space X is a collection $\{U_\lambda \mid \lambda \in \Lambda\}$ of open subsets U_λ of X such that

$$\bigcup_{\lambda \in \Lambda} U_\lambda = X$$

2. A topological space X is **compact** if every open cover $\{U_\lambda \mid \lambda \in \Lambda\}$ of X has a finite subcover, i.e. there exists $\lambda_1, \dots, \lambda_n \in \Lambda$ such that

$$X = \bigcup_{j=1}^n U_{\lambda_j}.$$

Definition 4.2: Open Covers as Collections

1. If $A \subseteq X$ is a subset of a topological space X , an **open cover** of A is a collection $\{V_\lambda \mid \lambda \in \Lambda\}$ of subsets V_λ which are open in X such that

$$X = \bigcup_{\lambda \in \Lambda} V_\lambda$$

2. A subset A of a topological space X is **compact** if it is compact as a subspace of X .

Proposition 4.7: Boundedness of Compact Spaces

A compact metric space (X, d) is bounded, i.e. there exists a number $K \geq 0$ such that $d(x, y) \leq K$ for all $x, y \in X$.

Proposition 4.8: Compactness of Products

A product of closed bounded intervals $R = [a_1, b_1] \times \dots \times [a_n, b_n] \subseteq \mathbb{R}^n$ is compact in the usual topology. A collection of subsets of a set X has the **finite intersection property** if every finite intersection of their members is nonempty.

Corollary 4.12: Limit Property of Compactness

Suppose that $f : X \rightarrow \mathbb{R}^n$ is a continuous map and that X is compact. Then there exists an M such that

$$|f(x)| \leq M \text{ for all } x \in X.$$

Moreover, there exists an $x \in X$ such that

$$|f(x)| = \sup_{y \in X} |f(y)|.$$

If $n = 1$ there are x_0 and $x_1 \in X$ such that

$$f(x_0) = \min_{x \in X} f(x) \quad \text{and} \quad f(x_1) = \max_{x \in X} f(x).$$

Theorem 4.13: Compact, Hausdorff, Bijection, Oh My!

Suppose X is compact, Y is Hausdorff, and that $f : X \rightarrow Y$ is a continuous bijection. Then it is a homeomorphism.

Theorem 4.14: Lebesgue Numbers

Let X be a compact metric space and $\{U_\lambda \mid \lambda \in \Lambda\}$ an open cover of X . Then there exists a positive number $\delta > 0$ (the **Lebesgue number** of the cover) such that for all $x \in X$, $B(x, \delta)$ lies *entirely inside some single* U_λ .

Corollary 4.17: Compactness of Identification Spaces

1. An identification space X/\sim of a compact space X is compact.
2. If $f : X \rightarrow Y$ is a map from a compact space X to a Hausdorff space Y and \sim is the equivalence relation on X defined by $x \sim x'$ if $f(x) = f(x') \in Y$, then the continuous bijection
$$g : X/\sim \rightarrow f(X); \quad [x] \mapsto f(x)$$
 is a homeomorphism.

Lemma 4.20: Open sets in Product spaces

Let X be a topological space, Y a compact space, $x \in X$, N an open set in $X \times Y$ such that $\{x\} \times Y \subseteq N$. Then there is an open set $W \subseteq X$ such that $x \in W$ and $W \times Y \subseteq N$.

Lemma 4.22 - 4.23: Collections and Intersections

- 4.22) Let X be a set, and suppose \mathcal{C} is a collection of subsets of X which has the finite intersection property. Then there is a collection \mathcal{B} of subsets of X , with $\mathcal{C} \subseteq \mathcal{B}$, such that \mathcal{B} has the finite intersection property, and such that \mathcal{B} is maximal with respect to this property: i.e. no collection containing \mathcal{B} as a proper subcollection has the finite intersection property.
- 4.23) Let X be a set, and suppose that \mathcal{B} is a collection of subsets of X which is maximal with respect to the finite intersection property. Then \mathcal{B} is closed under finite intersections, and any set which meets all members of \mathcal{B} is also in \mathcal{B} .

Definition 4.24: Compactifications

1. A **compactification** of a topological space X is a compact space Y which contains a homeomorphic copy of X as a subspace, i.e. such that there is a one-one map $f : X \rightarrow Y$ such that $X \rightarrow f(X); \quad x \mapsto f(x)$ is a homeomorphism.
2. A compactification Y is **dense** if X is dense in Y , i.e. $\overline{X} = Y$.

Definition 4.27: One-point compactification

The **one-point compactification** of a topological space X is the set

$$X^\infty = X \cup \{\infty\}$$

obtained by adjoining a “point at infinity” ∞ , where ∞ is a symbol *not* in X , with open sets of the form either

1. U , where $U \subseteq X$ is open, or
2. $X^\infty \setminus K$, where $K \subseteq X$ is compact and closed.

Lemma 4.28

1. The collection of open sets just defined does form a topology
2. The subspace topology on X induced by this topology coincides with its original topology.

Definition 4.32: Local Compactness

A topological space X is **locally compact** if for each $x \in X$, there exists an open subset $U \subseteq X$ and a compact C such that $x \in U \subseteq C$.

Remark 4.33

When X is Hausdorff, it is locally compact iff for each $x \in X$ there exists an open subset $U \subseteq X$ and a compact $x \in U$ and the closure \overline{U} is compact.

Definition 4.35: Normal Space

A topological space (X, \mathcal{T}) is **normal** if for every pair of disjoint closed subsets C and $D \subseteq X$, there are disjoint open subsets $U, V \subseteq X$ such that $C \subseteq U$ and $D \subseteq V$.

Lemma 4.37: Normal Complements

A space X is normal iff for every closed $F \subseteq X$ and open $G \subseteq X$ with $F \subseteq G$, there exist open G' and closed F' such that

$$F \subseteq G' \subseteq F' \subseteq G.$$

Theorem 4.38: Urysohn's Lemma

Suppose that X is a normal topological space, and that C, D are disjoint closed subsets of X . Then there is a continuous function $f : X \rightarrow \mathbb{R}$ such that

- $f(x) = 0$ for all $x \in C$
- $f(x) = 1$ for all $x \in D$
- $0 \leq f(x) \leq 1$ for all $x \in X$

Theorem 4.39: Tietze extension theorem

Suppose that X is a normal topological space, and that C is a closed subset of X . Suppose that $f : C \rightarrow \mathbb{R}$ is continuous. Then there is a continuous function $\bar{f} : X \rightarrow \mathbb{R}$ such that

- $\bar{f}(x) = f(x)$ for all $x \in C$
- If $a \leq f(x) \leq b$ for all $x \in C$, then $a \leq \bar{f}(x) \leq b$ for all $x \in X$.

Theorem 4.40: Stone-Weierstrass Theorem

The algebra A is dense in the normed space $C(X)$, i.e. $\overline{A} = C(X)$, i.e. for all $f \in C(X)$ and for all $\epsilon > 0$ there is $g \in A$ such that $\sup_{x \in X} |f(x) - g(x)| < \epsilon$.

5 Connected Spaces

Definition 5.1: Connected Spaces

1. A topological space X is **connected** if it *cannot* be written as a union

$$X = A \cup B$$

where A and B are disjoint nonempty open subsets of X

2. A topological space X is **disconnected** if it is not connected, i.e. if it *can* be expressed as a union

$$X = A \cup B$$

where A and B are disjoint nonempty open subsets of X

Connectedness is a **Topological Property** (See P6).

Remark 5.8: Connected Homeomorphisms

- If X is a compact connected metric space with exactly two points x such that $X \setminus \{x\}$ is connected, then X is homeomorphic to $[0, 1]$
- If X is a compact connected space, where for every pair of distinct points $x, y \in X$ the complement $X \setminus \{x, y\}$ is disconnected, then X is homeomorphic to the circle \mathbb{S}_1

Proposition 5.11: Connectedness other properties

The following statements about a nonempty subset $A \subseteq \mathbb{R}$ are equivalent:

1. A is connected
2. A has the interval property
3. A is an interval

Theorem 5.12: Intermediate Value Theorem

Let I be a closed bounded interval and suppose $f : I \rightarrow \mathbb{R}$ is continuous. Then the image $f(I)$ is a closed bounded interval

$$f(I) = [a, b] \subseteq \mathbb{R} (a \leq b).$$

Definition 5.13: Fixed Points of Maps

A **fixed point** of a map $f : X \rightarrow X$ is an $x \in X$ s.t. $f(x) = x$.

Theorem 5.15: 1-D Brouwer Fixed Point Theorem

Every continuous map $f : [0, 1] \rightarrow [0, 1]$ has a fixed point, i.e. there exists $x \in [0, 1]$ such that $f(x) = x$.

General Case: Every continuous map $f : \mathbb{D}^n \rightarrow \mathbb{D}^n$ has a fixed point

Definition 5.16: Path

A **path** in a topological space X is a continuous map $\alpha : I = [0, 1] \rightarrow X$. Its **initial point** is $\alpha(0) \in X$ and its **terminal point** is $\alpha(1) \in X$.

Definition 5.18: Path Connectedness

A topological space X is **path-connected** if for any two points $x_0, x_1 \in X$ there exists a path $\alpha : I \rightarrow X$ from $\alpha(0) = x_0$ to $\alpha(1) = x_1$.

Theorem 5.24: Homeomorphisms of Real Spaces

If $n \geq 2$, the spaces \mathbb{R}^n and \mathbb{R} are not homeomorphic. Additionally, there is no bijection $f : \mathbb{R} \rightarrow \mathbb{R}^n$ which is continuous.

Definition 5.35: Connected Components

We define an equivalence relation \sim on a topological space x by $x \sim y$ iff there is a connected subset of X which contains both x and y . The resulting equivalence classes are called the **components** or **connected components** of X . For two homeomorphic topological spaces there will be a bijection between the sets of their components. So spaces with differing numbers of components cannot be homeomorphic. The components are the maximal connected subsets of a topological space and by Prop 5.9 are always closed in X . If we take $U \subseteq \mathbb{R}$ an open set, its connected component decomposition is its canonical representation as in Thm 1.30 as a countable disjoint union of open intervals.

Lemma 5.31.5: Path Components

Define a path (equivalence) relation

$$x_0 \sim x_1 \text{ if there exists a path } \alpha : I \rightarrow X \text{ from } \alpha(0) = x_0 \in X \text{ to } \alpha(1) = x_1 \in X.$$

5.32) The **constant path** at $x \in X$ is the path

$$\alpha_x : I \rightarrow X; \quad t \mapsto x$$

from $\alpha_x(0) = x \in X$ to $\alpha_x(1) = x \in X$

5.33) The **reverse** of a path $\alpha : I \rightarrow X$ is the path

$$-\alpha : I \rightarrow X; \quad t \mapsto \alpha(1 - t)$$

retracting α backwards, with

$$\begin{array}{ccc} -\alpha(0) = \alpha(1) & & -\alpha(1) = \alpha(0) \\ \bullet & \xrightarrow{-\alpha} & \bullet \end{array}$$

5.34) The **concatenation** of paths $\alpha : I \rightarrow X$, $\beta : I \rightarrow X$ with

$$\alpha(1) = \beta(0) \in X$$

is the path

$$\alpha \bullet \beta : I \rightarrow X, \quad t \mapsto \begin{cases} \alpha(2t) & \text{if } 0 \leq t \leq 1/2 \\ \beta(2t - 1) & \text{if } 1/2 \leq t \leq 1 \end{cases}$$

which starts at $\alpha(0)$, follows along α at twice the speed in the first half, switching at $\alpha(1) = \beta(0)$ to follow β at twice the speed in the second half.

$$\begin{array}{ccccc} \alpha \bullet \beta(0) = \alpha(0) & & \alpha(1) = \beta(0) & & \beta(1) = \alpha \bullet \beta(1) \\ \bullet & \xrightarrow{\alpha} & \bullet & \xrightarrow{\beta} & \bullet \end{array}$$

Lemma 5.31: Connected Components and Openness

Let X be a topological space and C a connected component of X . Then C is open iff for all $x \in C$ there is an open connected V such that $x \in V \subseteq C$.

Proposition 5.35: Equivalence of Path Relations

The path relation defined on a space X by $x_0 \sim x_1$ if there exists a path $\alpha : I \rightarrow X$ from $\alpha(0) = x_0$ to $\alpha(1) = x_1$ is an equivalence relation.

Definition 5.36: Path Components Formally

Let X be a topological space.

1. The **path components** of X are the equivalence classes of the path equivalence relation \sim , i.e. the subspaces

$$\begin{aligned} [x] &= \{y \in X \mid y \sim x\} \\ &= \{y \in X \mid \exists \alpha : I \rightarrow X \text{ from } \alpha(0) = x \text{ to } \alpha(1) = y\} \end{aligned}$$

2. The **set of path components** (which may be infinite) is denoted by

$$X / \sim = \pi_0(X)$$

3. The function

$$X \rightarrow \pi_0(X), \quad x \mapsto [x] = \{\text{equivalence class of } x\}$$

is surjective.

Lemma 5.39: Open Condition of Path Components

Let X be a topological space and P a path component of X . Then P is open iff for all $x \in P$ there is an open path connected V such that $x \in V \subseteq P$.

Lemma 5.40: Openness and Singular Components

Let C be a connected component of a topological space X . If every path component $P \subseteq C$ is open, then C consists of a single path component. Note that the converse of this is not true.

6 Relations between Top Props

Proposition A: Topological Invariants

A **topological property** of a topological space is one which is **invariant** under homeomorphism. Let $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{U})$ be a homeomorphism. The following properties are true:

2.8) \mathcal{U} is open in Y iff $f^{-1}(\mathcal{U})$ is open in X .

- X is Hausdorff iff Y is Hausdorff.

3.6) $X \setminus \{x_0\}$ is homeomorphic to $Y \setminus \{f(x_0)\}$.

4.11) X is compact, iff Y is compact.

5.6) X is connected iff Y is connected.

5.21) X is path-connected iff Y is path-connected.

5.37) There exists a bijection between the set of path components $\pi_0(X)$ and $\pi_0(Y)$. However, existence of a bijection between $\pi_0(X)$ and $\pi_0(Y)$ does *not* necessarily imply that X and Y are homeomorphic.

Proposition B: Hausdorff if...

3.4) Suppose (X, \mathcal{T}) is a Hausdorff topological space and suppose A is a subspace. Then A is Hausdorff.

4.34) The one-point compactification X^∞ of a space X is Hausdorff iff X is Hausdorff and locally compact.

Proposition C: Compact if...

4.3) Let X be a topological space and $A \subseteq X$. Then A is compact iff every open cover of A has a finite subcover.

4.5) **Heine-Borel Theorem:** A subset $F \subseteq \mathbb{R}^n$ is compact iff it is closed and bounded.

4.6) Let X be a topological space and $A \subseteq X$.

1. If X is compact and A is closed, then A is compact
2. If X is Hausdorff and A is compact, then A is closed.

4.10) Let $f : X \rightarrow Y$ be a continuous map between topological spaces. If X is compact, so is $f(X)$.

4.18) **Tychonoff's Theorem:** Suppose X and Y are compact spaces. Then their product $X \times Y$ is compact. The converse is also true.

4.21) **Tychonoff's Theorem (General):** Suppose that \mathcal{A} is an indexing set and that for each $\alpha \in \mathcal{A}$, X_α is a compact topological space. Then the product $\prod_{\alpha \in \mathcal{A}} X_\alpha$ is compact.

4.30) Suppose $X^\infty = X \cup \{\infty\}$ is the *one-point compactification* of X . Then either X^∞ is compact, or X is dense in X^∞

Proposition D: Continuous if..

2.14) Suppose (X, \mathcal{T}) and (Y, \mathcal{U}) are topological spaces and that $f : X \rightarrow Y$. Then f is continuous iff for every closed subset $F \subseteq Y$ its inverse image $f^{-1}(F)$ is closed in X .

2.14) f is continuous iff the image of the closure of every subset $A \subseteq X$ is contained in the closure of the image, i.e.,
 $\forall A \subseteq X,$

$$f(\overline{A}) \subseteq \overline{f(A)}$$

3.5) Suppose (X, \mathcal{T}) and (Y, \mathcal{U}) are topological spaces and suppose A is a subspace of X . Let $f : X \rightarrow Y$ be continuous. Then $f|_A : A \rightarrow Y$ is continuous.

3.12) Let (X, \mathcal{T}) and (Y, \mathcal{U}) be topological spaces and \mathcal{T} the product topology on $X \times Y$. Then the projection maps Π_X and Π_Y are continuous. Moreover, \mathcal{T} is the smallest topology on $X \times Y$ such that the projection maps are continuous.

3.13) Let X, Y, Z be topological spaces. Endow $X \times Y$ with the product topology. A function $f : Z \rightarrow X \times Y$ is continuous iff the functions $\Pi_X \circ f : Z \rightarrow X$ and $\Pi_Y \circ f : Z \rightarrow Y$ are both continuous.

Let X be a topological space with an equivalence relation \sim .

1. The function $p : X \rightarrow X/\sim; \quad x \mapsto [x]$ is continuous.
2. A continuous function $f : X \rightarrow Y$ such that $f(x) = f(x') \in Y$ for all $x, x' \in X$ with $x \sim x'$ determines a continuous function

$$g : X/\sim \rightarrow Y; \quad [x] \mapsto f(x)$$

such that

$$f = g \circ p : X \xrightarrow{p} X/\sim \xrightarrow{g} Y$$

$f = g \circ p$ is best described by a commutative triangle:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow p \quad \nearrow g & \\ & X/\sim & \end{array}$$

In fact, every continuous function on X determines an equivalence relation.

Proposition E: Connected if...

5.2) X is connected iff the only subsets of X which are clopen are \emptyset and X

5.4) \mathbb{R} with the usual topology is connected.

5.5) If $f : X \rightarrow Y$ is continuous and X is connected, then $f(X)$ (with the subspace topology) is connected.

5.9) Let A be a connected subsets of a topological space X and suppose $A \subseteq B \subseteq \overline{A}$. Then B is connected.

5.10) Every nonempty interval $I \subseteq \mathbb{R}$ is connected.

5.25) If a topological space X is path-connected, then it is also connected. Note that the converse need not be true.

5.30) Let $A_\lambda \subseteq X, (\lambda \in \Lambda)$ be a family of connected subsets of a topological space X . Suppose $\bigcup_{\lambda \in \Lambda} A_\lambda \neq \emptyset$. Then $\bigcup_{\lambda \in \Lambda} A_\lambda$ is connected.

Proposition F: Path-Connected if...

- Suppose $f : X \rightarrow Y$ is a continuous map between topological spaces and that X is path-connected. Then $f(X)$ is path-connected as a subspace of Y .

- For any equivalence relation \sim on a path-connected space X the identification space $Y = X/\sim$ is path-connected.

- Any connected open subset $\Omega \subseteq \mathbb{R}^n$ is also path-connected.

- Let X be a topological space. Then X is path connected iff X is connected *and* for all $x \in X$ there is an open path connected V such that $x \in V$.

7 Examples

Example 7.0.1: Other Topologies and Metrics

If (X, \mathcal{T}) is a topological space, and X admits a metric whose metric topology is precisely \mathcal{T} , then we say that (X, \mathcal{T}) is **metrisable**.

- Euclidean spaces with their usual topologies are metrisable.

1.9) The **Discrete Topology** is the topology of all subsets of a set X . We can define the **discrete metric** of X to be

$$d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{otherwise} \end{cases}$$

1.10) The **Trivial** or **Indiscrete Topology** is the topology $\mathcal{T} := \{\emptyset, X\}$ for a set X . This is a non-metrisable topology when X has more than one member.

1.14) Let $X = \{a, b, c\}$, where a, b, c are distinct. Then

$$\mathcal{T} = \{\emptyset, X, \{a\}, \{a, b\}, \{a, c\}\}$$

is a topology on X

1.15) Give \mathbb{R} the topolgy whose open subsets $U \subseteq \mathbb{R}$ are precisely the subsets with finite complement $\mathbb{R} \setminus U$, or $U = \emptyset$. Then \mathbb{R} with this topology is not metrisable. This is an example of a **Zariski Topology**

Example B: The Punctured Sphere




Figure 1: Homeomorphism of S^2 to \mathbb{R}^n

Example C: Topological Objects




Figure 2: Standard Topological Objects




Fig 3: Möbius Strip

Fig 4: Torus

Fig 5: Klein Bottle

Fig 6: \mathbb{RP}^2

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accu-

msan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent feugiat sapien aliquet odio. Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at, consectetur sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget lectus. Proin eu metus. Sed porttitor. In hac habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui. Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis, ultrices a, dui.

Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam. Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetur. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros

tellus scelerisque quam, pellentesque hendrerit ipsum dolor sed augue. Nulla nec lacus.

Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui. Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros pede, suscipit ac, varius

vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem dictum tortor, vel consectetur odio sem sed wisi.

Sed feugiat. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Ut pellentesque augue sed urna. Vestibulum diam eros, fringilla et, consectetur eu, nonummy id, sapien. Nullam at lectus. In sagittis ultrices mauris. Curabitur malesuada erat sit amet massa. Fusce blandit. Aliquam erat volutpat. Aliquam euismod. Aenean vel lectus. Nunc imperdiet justo nec dolor.

Etiam euismod. Fusce facilisis lacinia dui. Suspendisse potenti. In mi erat, cursus id, nonummy sed, ullamcorper eget, sapien. Praesent pretium, magna in eleifend egestas, pede pede pretium lorem, quis consectetur tortor sapien facilisis magna. Mauris quis magna varius nulla scelerisque imperdiet. Aliquam non quam. Aliquam porttitor quam a lacus. Praesent vel arcu ut tortor cursus volutpat. In vitae pede quis diam bibendum placerat. Fusce elementum convallis neque. Sed dolor orci, scelerisque ac, dapibus nec, ultricies ut, mi. Duis nec dui quis leo sagittis commodo.