

Algebraic Topology Notes

Made by Leon :) *Note: Any reference numbers are to the lecture notes*

1 Introduction

Recall 1.1.1: Topology

An **(open) topology** on X is a collection of subsets $\tau \subset P(X)$ such that

- $\emptyset \in \tau$ and $X \in \tau$
- τ is closed under finite intersections: If $\{U_1, \dots, U_n\} \subset \tau$ then
- τ is closed under arbitrary unions: If $\{U_1, \dots, U_n\} \subset \tau$ is a family of open subsets then

$$\bigcap_{i=1, \dots, n} U_i \in \tau \qquad \bigcup_{i=1, \dots, n} U_i \in \tau$$

The subsets $U \in \tau$ are called **open** and their complements in X define **closed subsets**.

Two examples of a topology on a set X are the following:

- The **Trivial Topology**: $\tau_{\text{triv}} = \{\emptyset, X\}$
- The **Discrete Topology**: $\tau_{\text{dis}} = P(X)$

A subset $A \subset X$ is **clopen** if it is both closed and open

Definition 1: Connected Spaces

A topological space X is **connected** if $X = A \amalg B$ with $A, B \subset X$ open implies that $A = \emptyset$ or $A = X$.

Proposition 1: Connectedness and Clopens

A topological space X is *connected* iff the only clopens are \emptyset and X .

Example 1: Examples of Connected Topologies

- Every X with the trivial topology is connected.
- Every X with the discrete topology isn't connected unless $X = \emptyset$ or $X = \{*\}$ (in which it coincides with the trivial topology).
- The real line \mathbb{R} with the standard topology is connected.

Proposition 2: Continuous Maps

Let $f : X \rightarrow Y$ be a continuous map of topological spaces and let X be connected. Then $f(X)$ is connected.

Proposition 3: Connected Equivalence Relation

For a topological space X , define $x \sim y$ if there exists some connected subset that contains both. The relation $x \sim y$ is an equivalence relation.

Definition 2: Connected Components

The equivalence classes of this relation are called **connected components**. In particular, a space X is connected iff it only has a single connected component.

Definition 3: Path

Let I denote the closed unit interval $[0, 1]$. A **path** in X is a continuous map $\alpha : I \rightarrow X$. The points $\alpha(0) \in X$ and $\alpha(1) \in X$ will be called **start** and **end** points respectively.

We define a path relation between points in X by declaring $x \sim y$ if there exists some path $\alpha : I \rightarrow X$ that starts at x and ends in y , i.e. $\alpha(0) = x$ and $\alpha(1) = y$. This is an equivalence relation from the following properties:

1. **Constant Path**: For all $x \in X$ there exists the constant path $c_x : I \rightarrow X$ defined by $c_x(t) = x$ for all $t \in I$
2. **Path reversal**: Let $\alpha : I \rightarrow X$ be a path in X . Define its reversed path by

$$\bar{\alpha} : I \rightarrow X, \quad t \mapsto \alpha(1 - t) \tag{1}$$

3. **Path Concatenation**: Let $\alpha, \beta : I \rightarrow X$ be two paths in X s.t. $\alpha(1) = \beta(0)$. Their concatenated path is defined by:

$$\alpha * \beta(t) := \begin{cases} \alpha(2t), & 0 \leq t \leq 1/2 \\ \beta(2t - 1) & 1/2 \leq t \leq 1 \end{cases} \tag{2}$$

Definition 4: Path-Connected Components

The equivalence classes are called **path-connected components** and their set is denoted by $\pi_0(X)$. A space X is called **path-connected** if $\pi_0(X)$ is a one-point set, i.e. any two points x, y can be related by a path in X .


Remark 1: Random examples

The following statements are true:

- A homeomorphism $X \cong Y$ induces a bijection $\pi_0(X) \cong \pi_0(Y)$.
- If X is path-connected, it is also connected.
- The *topologist's sine curve* defined by $X = \{0\} \times [-1, 1] \times \{(x, \sin(1/x)) \mid 0 < x\}$ is connected but not path-connected.

Definition 5: Homotopy

A **homotopy** of maps $f, g : X \rightarrow Y$ is a continuous map $h : X \times I \rightarrow Y$ such that $h(-, 0) = f$ and $h(-, 1) = g$.



If such a homotopy exists, f is **homotopic** to g . This defines an equivalence relation $f \simeq g$ on the space of maps $\text{Map}(X, Y)$.

Example 2: Paths as Homotopies

Points in X are the same as maps $* \rightarrow X$ from the one-point set $*$ to X . A path $\alpha : I \rightarrow X$ corresponds to a homotopy $* \times I \rightarrow X$.

Remark 1.5: Composition of Homotopies

- **Vertical Composition**: Let $h, h' : X \times I \rightarrow Y$ be two homotopies in X such that $h(-, 1) = h'(-, 0) : X \rightarrow Y$. Their concatenated homotopy is defined by


$$h * h'(-, t) := \begin{cases} h(-, 2t) & 0 \leq t \leq 1/2 \\ h'(-, 2t - 1) & 1/2 \leq t \leq 1 \end{cases} \tag{4}$$

- **Horizontal Composition**: Let $h : X \times I \rightarrow Y, k : Y \times I \rightarrow Z$ be two homotopies on maps from X to Y , and Y to Z . Then

$$k \circ h := [X \times I \xrightarrow{\text{id} \times \Delta} X \times I^2 \xrightarrow{h \times \text{id}} Y \times I \xrightarrow{k} Z] \tag{5}$$

where $\Delta : I \rightarrow I^2, t \mapsto (t, t)$ is the diagonal map, or explicitly,

$$k \circ h(x, t) = k(h(x, t), t)$$



Lemma 1: Concatenation Relation

Let $f, f' : X \rightarrow Y$ and $g, g' : Y \rightarrow Z$ be maps such that $f \simeq f'$ and $g \simeq g'$. Then $g \circ f \simeq g' \circ f'$ as maps from X to Z . In particular, $g' \circ f \sim g \circ f$ and $g \circ f' \sim g \circ f$.

Definition 6: Homotopy Equivalence

A map $f : X \rightarrow Y$ is called a **homotopy equivalence** if there exists a map $g : Y \rightarrow X$ and homotopies $f \circ g \simeq \text{id}_Y, g \circ f \simeq \text{id}_X$. In other words, g satisfies the properties of an inverse up to homotopy. It is called a **homotopy inverse** of f .

Example 3: Circle to \mathbb{R}^2

The inclusion map $\mathbb{S}^1 \hookrightarrow \mathbb{R}^2$ is not a homotopy equivalence, but the inclusion $\mathbb{S}^1 \hookrightarrow \mathbb{R}^2 \setminus \{0\}$ is a homotopy equivalence.

Proposition 4: Unique Inverses of Homotopy

Homotopy inverses are unique up to homotopy.

Definition 7: Homotopic Spaces

Two spaces X and Y are called **homotopy equivalent**, or of the **same homotopy type**, and denoted by $X \simeq Y$ if there exists a homotopy equivalence $f : X \rightarrow Y$.

Note: \cong for homeomorphisms and \simeq for homotopy equivalence.

Lemma 2: Composition of Inverses

Let $f : X \rightarrow y, g : Y \rightarrow Z$ with homotopy inverses $\bar{f} : Y \rightarrow X$ and $\bar{g} : Z \rightarrow Y$ respectively. Then $\bar{f} \circ \bar{g} : Z \rightarrow X$ is a homotopy inverse of $g \circ f : X \rightarrow Z$. In particular, $X \simeq Y, Y \simeq Z$ implies $X \simeq Z$.

2 Contractible Spaces

Definition 8: Contractible Space

A space X is called **contractible** if it is homotopy equivalent to a point, i.e. $X \simeq *$.

The **terminal map** is the unique map $X \rightarrow *$. Contractibility requires that there is a homotopy inverse of that map, i.e. a map $*$ \rightarrow x along with homotopies

$$h : [* \rightarrow X \rightarrow *] \simeq \text{id}_*, \quad k : [X \rightarrow * \rightarrow X] \simeq \text{id}_X \quad (6)$$

Example 4: Examples of Contractible Spaces

- \mathbb{R}^n is contractible. Let x_0 be a fixed point in \mathbb{R}^n and define the (straight line) homotopy $h : c_{x_0} \simeq \text{id}_{\mathbb{R}^n}$ by

$$h(x, t) = (1 - t)x_0 + tx.$$

- $\mathbb{S}^{n-1} \simeq \mathbb{R}^n \setminus \{0\}$. The inclusion $\mathbb{S}^{n-1} \hookrightarrow \mathbb{R}^n \setminus \{0\}$ and the shrinking map

$$\mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{S}^{n-1}, \quad x \mapsto \frac{x}{|x|}$$

are homotopy inverses.

Remark 3: Remarks about Contractible Spaces

- Contractible spaces are path-connected. Let x_0 be the point where the space X contracts to. In particular, we are given with a homotopy $h : c_{x_0} \simeq \text{id}_X$. For any $x \in X$, the map $h(x, -) : I \rightarrow X$ defines a path from x_0 to x and thus every element $x \in X$ is path-connected to x_0 .
- The converse does not hold, for example $X = \mathbb{S}^1$.
- A contractible space X is contractible at any point x_0 . X is path-connected, so a path x to x' defines a homotopy $c_x \simeq c_{x'}$.
- Any two maps $f, g : X \rightarrow Y$ are homotopic if Y is contractible.

Definition 9: Retracts and Deformation Retracts

- A **retract** of X onto a subspace $A \subset X$ is a map $r : X \rightarrow A$ such that $r|_A = \text{id}_A$. Equivalently, this is a map $r : X \rightarrow X$ such that $r^2 = r$ and $r(X) = A$. Or, s.t. $ri = 1 : A \rightarrow A$, where $i : A \rightarrow X$ is the inclusion map.
- A **deformation retract** of X onto A is the additional datum of a homotopy $h : \text{id}_X \simeq i \circ r$.

In other words, a deformation retract is a homotopy $h : X \times I \rightarrow X$ such that $h(x, 0) = x$ and $h(x, 1) \in A$ for all $x \in X$ and $h(a, 1) = a$ for all $a \in A$. Not all retracts can form deformation retracts. For instance, the retract X onto a point $\{x_0\}$ can be a deformation retract iff X is contractible.

Remark 4: Strong vs Weak Deformation Retracts

This notion is called **weak** deformation retract. A **strong** deformation retract has the condition $h(a, t) = a$ for all $t \in I$, $a \in A$. Our notion of a (weak) deformation retract deforms X into A while allowing to deform A to do so, while a strong deformation retract deforms X into A while keeping A fixed at all times

Proposition 5: Deformation Retracts and Homotopies

A deformation retract of X onto A induces a homotopy equivalence $X \simeq A$.

Recall 2: Quotient Space

Let X be a topological space and let \sim be an equivalence relation on X . Then, X/\sim is equipped with the quotient topology and called a **quotient space**. If Z is a closed subset in X , then we can also define the quotient space X/Z .

Another form of quotient spaces: Let $f : Z \rightarrow Y$ be a continuous map between a closed subset $Z \subset X$ and Y . Then

$$X \amalg_f Y = X \amalg Y / z \sim f(z).$$

Example 5: Examples of Quotient Spaces

- The quotient of the n -dimensional closed disk by its boundary is the n -sphere, i.e. $\mathbb{D}^n / \partial \mathbb{D}^n \cong \mathbb{S}^n$.
- The 2-torus: $\mathbb{R}^2 / \mathbb{Z}^2$.
- The projective space: $\mathbb{RP}^n = \mathbb{R}^{n+1} \setminus \{0\} / \sim$ by the relation $x \sim y$ iff there exists some $\lambda \in \mathbb{R}^\times$ such that $x = \lambda y$. This corresponds to the space of lines through the origin in \mathbb{R}^{n+1} .

Definition 10: Mapping Quotients

Let $f : X \rightarrow Y$ be a continuous map.

- Its **mapping cylinder** is defined as the topological space

$$M_f := (X \times I) \amalg Y / \sim$$

where the quotient identifies $(x, 0) \sim f(x)$ for any $x \in X$.

- Its **cone** is the further quotient: $C_f = M_f / X \times \{1\}$.
- The **cone** of a topological space X is $C_X := C_{\text{id}_X} = X \times I / X \times \{1\}$.

In other words, the mapping cylinder of $f : X \rightarrow Y$ is the pushout of the diagram:

$$\begin{array}{ccc} X \times \{0\} & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ X \times I & \longrightarrow & M_f \end{array}$$

Example 5.5: Spheres

For \mathbb{S}^n with the standard embedding $\mathbb{R}^{n+1} \setminus \{0\}$, the following map is a retract, because if x has norm $|x| = 1$, then $r(x) = x$.

$$r : \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{S}^n, \quad x \mapsto \frac{x}{|x|}$$

For a deformation retract one needs to find a homotopy $h : i \circ r \simeq \text{id}_X$. We use the following straight-line homotopy:

$$h : \mathbb{R}^{n+1} \setminus \{0\} \times I \rightarrow \mathbb{R}^{n+1} \setminus \{0\}, \quad (x, t) \mapsto (1 - t) \frac{x}{|x|} + tx.$$

Indeed, $h(x, 0) = r(x)$ and $h(x, 1) = x$ for all x .

Definition 11: Star-Shaped Spaces

A subset $S \subset \mathbb{R}^n$ is called **star-shaped** at a point $x_0 \in S$, if for any $x \in S$ the line segment from x_0 to x is contained in S , i.e.

$$\{(1 - t)x_0 + tx \mid t \in [0, 1]\} \subset S$$

If S is star-shaped at every point, then it is called **convex**.

Example 5.6: Star-Shaped Spaces are Contractible

Let S be star-shaped at x_0 and $i : \{x_0\} \hookrightarrow S : r$ be the inclusion and constant maps. Define the straight line homotopy

$$h : S \times I \rightarrow S, \quad (x, t) \mapsto (1 - t)x_0 + tx$$

which is well-defined by the star-shaped condition. Moreover, $h(x, 0) = x_0 = r(x)$ and $h(x, 1) = x$ for all x . Hence, star-shaped, and in particular convex spaces, are contractible.

Example 5.7: Möbius band

The Möbius band M can be defined as

$$M = I^2 / \sim$$

where \sim identifies the two vertical edges of I^2 by flipping one, i.e. $(0, b) \sim (1, 1 - b)$ for $b \in I$. Its core $C \subset M$ is the line $\{[a, 1/2] \mid a \in I\}$. Thus, the core is homeomorphic to \mathbb{S}^1 . The Möbius band deformation retracts onto its core, e.g. the retract $r : M \rightarrow C$ defined by $r([a, b]) := [(a, 1/2)]$ and the homotopy

$$h : M \times I \rightarrow M, \quad ([a, b], t) \mapsto \left[\left(a, (1 - t) \frac{1}{2} + \right) \right].$$

In particular, $M \simeq \mathbb{S}^1$.

Proposition 6: Retracts of the Mapping Cylinder

Via Definition 10, the mapping cylinder is formed by the cylinder of X by gluing Y onto the bottom with the map f . The mapping cylinder M_f strongly deformation retracts onto Y .

Proof. Consider the retract:

$$r : M_f \rightarrow Y$$

defined by $r([x, s]) := [(x, 0)] = [f(x)]$ on the class of $(x, s) \in X \times I$ and $r([y]) = y$ for $y \in Y$. This is well-defined and by definition a retract on Y . Define the homotopy

$$h : M_f \times I \rightarrow M_f$$

by $h([x, s], t) := [(x, st)]$ for $(x, s) \in X \times I$ and $t \in I$, and by $h([y], t) := y$ for $y \in Y$. In particular, $h(-, 0) = i \circ r$ and $h(-, 1) = \text{id}_{M_f}$. This forms a strong deformation retract. \square

Remark 6: Continuous Maps are Homotopic

Any continuous $f : X \rightarrow Y$ can be replaced up to homotopy equivalence by the closed inclusion $X \hookrightarrow M_f$, $x \mapsto [(x, 1)]$. More precisely, it factorises through an inclusion and a homotopy equivalence as the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow & \nearrow \simeq \\ & M_f & \end{array}$$

Definition 12: Relative Homotopy

Let X, Y be topological spaces and $A \subset X$ a subset in X . A homotopy $h : X \times I \rightarrow Y$ is called **relative to A** if $h(a, t)$ is independent of t for all $a \in A$. In particular, this defines homotopies between maps $f, g : X \rightarrow Y$ such that $f|_A = g|_A$.

This definition generalises ordinary homotopies, as an ordinary homotopy is the same as a homotopy relative to \emptyset .

Example 6: Relative Homotopies and Retracts

A strong deformation retract of X onto A is a deformation retract such that the homotopy $h : i \circ r \simeq \text{id}_X$ is relative to A .

Definition 13: Homotopic Path

Let $\alpha, \beta : I \rightarrow X$ be paths in X such that $\alpha(0) = \beta(0)$ and $\alpha(1) = \beta(1)$. A relative homotopy from α to β is a homotopy $h : I \times I \rightarrow X$ relative to $\partial I = \{0, 1\}$, i.e.

$$h(-, 0) = \alpha, \quad h(-, 1) = \beta \quad (7)$$

and

$$h(0, t) = \alpha(0) = \beta(0), \quad h(1, t) = \alpha(1) = \beta(1), \quad \forall t \in I. \quad (8)$$

In particular, at any point $t \in I$ a relative homotopy h defines a path $h_t := h(-, t) : I \rightarrow X$ with start $\alpha(0) = \beta(0)$ and end $\alpha(1) = \beta(1)$. If one omits the relative condition, the start and end points of h_t would be allowed to vary.

Remark 7: Ordinary Homotopies and Paths

Ordinary homotopies are not well suited for paths: Any path $\alpha : I \rightarrow X$ is homotopic (rel. \emptyset) to a constant - as the homotopy

$$h : I \times I \rightarrow X, \quad (s, t) \mapsto \alpha(st)$$

defines a homotopy from the constant path $c_{\alpha(0)}$ on $\alpha(0)$ to α , i.e. $c_{\alpha(0)} \simeq \alpha$. Hence, (ordinary) homotopy classes of paths in X are in 1-to-1 correspondence with path-connected components of X .

Proposition 7: Homotopic Properties of Paths

Path concatenation is **unital**, **associative**, and **invertible** up to homotopy in the following sense: Let $\alpha, \beta, \gamma : I \rightarrow X$ be paths such that $\alpha(1) = \beta(0)$ and $\beta(1) = \gamma(0)$. Then there exists homotopies relative to $\{0, 1\}$:

1. **Left Unitality:** $c_{\alpha(0)} * \alpha \simeq \alpha$
2. **Right Unitality:** $\alpha \simeq c_{\alpha(0)} * \alpha$
3. **Associativity:** $(\alpha * \beta) * \gamma \simeq \alpha * (\beta * \gamma)$
4. **Right Inverse:** $\alpha * \bar{\alpha} \simeq c_{\alpha(0)}$
5. **Left Inverse:** $\bar{\alpha} * \alpha \simeq c_{\alpha(1)}$

where c_x for some $x \in X$ denotes the constant path on x and $\bar{\alpha}$ is the reversed path.

Lemma 3:

Let $\alpha : I \rightarrow X$ be a path and $\lambda : I \rightarrow I$ a boundary preserving map, i.e. $\lambda(0) = 0$ and $\lambda(1) = 1$. Then,

$$\alpha \circ \lambda \simeq \alpha, \quad \text{rel. } \partial I.$$

Definition 14: Fundamental Group

Let X be a topological space and $x_0 \in X$ some fixed point. The **fundamental group** of X at x_0 is the group of homotopy classes of paths in X that start and end on x_0 . i.e. $\alpha : I \rightarrow X$ such that $\alpha(0) = \alpha(1) = x_0$, i.e.

$$\pi_1(X, x) = \{\alpha : I \rightarrow X \mid \alpha(0) = \alpha(1)\} / \sim.$$

Theorem 1: Defining the Fundamental Group

The fundamental group $\pi_1(X, x_0)$ is a well-defined group with:

- **Multiplication:** $[\alpha] \cdot [\beta] := [\alpha * \beta]$
- **Unit:** $1 = [c_{x_0}]$
- **Inverse:** $[\alpha]^{-1} = [\bar{\alpha}]$

Lemma 4: Relative Concatenated Homotopic Paths

Let $\alpha \simeq \alpha' : I \rightarrow X$ and $\beta \simeq \beta' : I \rightarrow X$ be two pairs of relative homotopic paths such that $\alpha(1) = \alpha'(1) = \beta(0) = \beta'(0)$. Then the concatenations are relative homotopic, i.e.

$$\alpha * \beta \simeq \alpha' * \beta', \quad \text{rel. } \{0, 1\}.$$

Proposition 8: Fundamental Group is Point Independent

Let $\gamma : I \rightarrow X$ be a path from $\gamma(0) = x$ to $\gamma(1) = x'$. Then it induces a group isomorphism:

$$(\gamma)_\# : \pi_1(X, x) \rightarrow \pi_1(X, x'), \quad [\alpha] \mapsto [\bar{\gamma} * \alpha * \gamma].$$

We abuse notation to say that for a path-connected space X , $\pi_1(X)$ is the fundamental group omitting the choice of base point.

Example 7: Examples of Fundamental Groups

- **Euclidean:** $\pi_1(\mathbb{R}^n) \cong 1$.
- **n -Sphere,** $n \geq 2$: $\pi_1(\mathbb{S}^n) \cong 1$.
- **Circle:** $\pi_1(\mathbb{S}^1) \cong \mathbb{Z}$.
- **Torus:** $\pi_1(\mathbb{S}^1 \times \mathbb{S}^1) \cong \mathbb{Z} \oplus \mathbb{Z}$.
- **Projective Spaces:** $\pi_1(\mathbb{RP}^n) \cong \mathbb{Z}_2$ for $n \geq 2$.

Definition 15: Pointed Space and Loop Space

- A **pointed space** is a pair (X, x) consisting of a topological space X and a point $x \in X$.
- A **map of pointed spaces** $f : (X, x) \rightarrow (Y, y)$ is a continuous map $f : X \rightarrow Y$ such that $f(x) = y$.
- The **space of pointed maps** from (X, x) to (Y, y) is denoted

$$\text{Map}_*((X, x), (Y, y)) \subset \text{Map}(X, Y).$$

With the (pointed) homeomorphism $(\mathbb{S}^1, 1) \cong (I/\partial I, [0])$, closed paths (where $\alpha(0) = \alpha(1) = x$) are the same as pointed maps

$$(\mathbb{S}^1, 1) \rightarrow (X, x)$$

The space of such **loops** based at x is called the **loop space at x** .

$$\Omega X := \text{Map}_*((\mathbb{S}^1, 1), (X, x))$$

It is itself a pointed space with the compact-open topology, and the constant map c_x as the base point. Path concatenation is the operation $* : \Omega X \times \Omega X \rightarrow \Omega X$ which is associative, unital, invertible up to path-connectedness, which gives a group structure

$$\pi_0(\Omega X).$$

Proposition 9: Loop Space Isomorphism

We have a group isomorphism: $\pi_1(X, x) \cong \pi_0(\Omega X)$.

Iteratively defining the n -fold loop space:

$$\Omega^n X := \Omega \Omega^{n-1} X = \Omega \cdot \Omega X$$

There is a homeomorphism: $\Omega^n X \cong \text{Map}_*((\mathbb{S}^n, 1), (X, x))$

Definition 16: n -th Homotopy Group

The n -th homotopy group $\pi_n(X, x)$ is defined by:

$$\pi_n(X, x) := \pi_0(\Omega^n X) \cong \pi_0(\text{Map}_*(\mathbb{S}^n, (X, x))).$$

Definition 17: Simply Connected Space

A path-connected space X is **simply connected** if its fundamental group is trivial, i.e. $\pi_1(X) = 1$.

Some examples are \mathbb{R}^n , \mathbb{S}^n for $n > 1$, and some non-examples are \mathbb{S}^1 , $\mathbb{S}^1 \times \mathbb{S}^1$ and \mathbb{RP}^2 .

Theorem 2: Fundamental Group Isomorphism

Let $f : X \rightarrow Y$ be a homotopy equivalence and $x \in X$ an arbitrary base point. Then, the following map is a group isomorphism:

$$f_* : \pi_1(X, x) \rightarrow \pi_1(Y, f(x)), \quad [\alpha] \mapsto [f \circ \alpha]$$

In particular, for homotopy equivalent spaces $X \simeq Y$ which are path-connected, we get $\pi_1(X) \cong \pi_1(Y)$.

A map of pointed spaces $f : (X, x) \rightarrow (Y, y)$ is a **homotopy equivalence of pointed spaces** or **homotopy equivalence relative $\{x\}$** if there exists a map of pointed spaces $g : (Y, y) \rightarrow (X, x)$ along with relative homotopies

$$h : f \circ g \simeq \text{id}_Y \quad \text{rel. } \{y\} \quad \text{and} \quad k : g \circ f \simeq \text{id}_X \quad \text{rel. } \{x\}$$

Example 9: Strong Deformation Retracts Homotopies

A strong deformation retract of X onto a subspace A gives a homotopy equivalence of pointed spaces $(x, a) \rightarrow (A, a)$ for any choice of $a \in A$. In particular, a contractible space $X \simeq *$ determines a homotopy equivalence of pointed spaces $(X, x) \rightarrow *$ for any choice of base point x .

Lemma 5: Pointed Space Isomorphism

Let $f : (X, x) \rightarrow (Y, y)$ be a homotopy equivalence of pointed spaces. Then the map

$$f_* : \pi_1(X, x) \rightarrow \pi_1(Y, y)$$

is a group isomorphism.

Corollary 1: Let $r : X \rightarrow A$ be a strong deformation retract of X onto $A \subset X$. Then for any $a \in A$,

$$\pi_1(X, a) \cong \pi_1(A, a)$$

In particular, contractible spaces are simply connected.

Lemma 6: Identity Homomorphic Isomorphism

Let $f : X \rightarrow X$ be a cts. map homotopic to id_X . Then, the map

$$f_* : \pi_1(X, x_0) \rightarrow \pi_1(X, f(x_0))$$

is a group isomorphism for any choice of base point $x_0 \in X$.

Definition 18: Homotopy Lifting Property

A continuous map $p : E \rightarrow X$ satisfies the **homotopy lifting property** (HLP) with respect to a topological space Y if for any commuting diagram:

$$\begin{array}{ccc} Y \times \{0\} & \xrightarrow{H_0} & E \\ \downarrow & \searrow \exists & \downarrow p \\ Y \times I & \xrightarrow{h} & X \end{array}$$

There exists a map $H : Y \times I \rightarrow E$ s.t. both triangles commute, i.e. $H|_{Y \times \{0\}} = H_0$ and $p \circ H = h$.

The map $p : E \rightarrow X$ has the HLP if for any homotopy $h : Y \times I \rightarrow X$ of maps $h(-, 0) := f_0$ and $h(-, 1) := f_1$ of maps $Y \rightarrow X$ and a choice of lift H_0 of f_0 , then the homotopy h lifts to a homotopy $H : Y \times I \rightarrow E$. In particular, if $f_0 \simeq f_1 : Y \rightarrow X$ and H_0 is a lift of f_0 , we find $H_0 \simeq H_1$ where H_1 lifts f_1 .

Ex. 10: The identity map $\text{id}_X : X \rightarrow X$ has the HLP with respect to any space Y .

Definition 19: Covering Space

A **covering space** of X is a topological space \bar{X} along with a continuous map $p : \bar{X} \rightarrow X$ s.t. for any point $x \in X$ there exists an open nbhd $U \subset X$ whose preimage $p^{-1}(U) = \bigcup_{j \in J} V_j$ and the opens $V_j \subset \bar{X}$ map homeomorphically to U under p . A covering space of X looks locally like a product of X with a discrete space.

Example 11: Example of a Covering Space

1. The projection map $p : X \times Z \rightarrow X$ is a covering map if Z is a discrete topological space. If Z is not discrete, then this is not a covering map in general.
2. The identity map $\text{id}_X : X \rightarrow X$ is trivially a covering map.
3. While the projection of $p : X \times I \rightarrow X$ from the cylinder is not a covering map, its restriction to the boundary $\partial(X \times I) = X \times \{0, 1\} =: \bar{X}$ gives a trivial (2-fold) cover of X .
4. Recall that the Möbius band M deformation retracts onto its core S^1 . Restricting to the boundary $\partial M = S^1$, one obtains a (non-trivial) covering map $S^1 \rightarrow S^1$. This map coincides with $z \mapsto z^2$ if we identify S^1 as the unit circle in \mathbb{C} .

Theorem 3: Unique HLPs from Covering Maps

Let $p : \bar{X} \rightarrow X$ be a covering map and Y any topological space. Then p satisfies the HLP uniquely: i.e. the lift H not only exists, but it is also unique.

$$\begin{array}{ccc} Y \times \{0\} & \xrightarrow{H_0} & \bar{X} \\ \downarrow & \searrow \exists! & \downarrow p \\ Y \times I & \xrightarrow{h} & X \end{array}$$

Corollary 2:

1. Let $\gamma : I \rightarrow X$ be a path and fix a point $\tilde{x}_0 \in \bar{X}$ such that $p(\tilde{x}_0) = \gamma(0)$. Then, there exists a unique path $\tilde{\gamma} : I \rightarrow \bar{X}$ which starts at \tilde{x}_0 and lifts γ i.e. $p \circ \tilde{\gamma} = \gamma$
2. Let $h : I \times I \rightarrow X$ be a (relative) homotopy of paths $h(-, 0) =: \gamma_0$ and $h(-, 1) =: \gamma_1$, and fix a point \tilde{x}_0 such that $p(\tilde{x}_0) = h(0, t) = \gamma_0(0) = \gamma_1(0)$. Suppose $\tilde{\gamma}_0 : I \rightarrow \bar{X}$ is a lift of γ_0 starting at $\tilde{\gamma}_0(0) = \tilde{x}_0$. Then, there exists a unique homotopy of paths $\tilde{h} : I \times I \rightarrow \bar{X}$ which lifts h and $\tilde{h}(-, 0) = \tilde{\gamma}_0$

Theorem 4-7: Fundamental Groups

- **Theorem 4:** The fundamental group of the circle is $\pi_1(S^1) \cong \mathbb{Z}$. It is generated by the class of

$$\alpha : I \rightarrow S^1, \quad t \mapsto e^{2\pi i t}.$$

- **Theorem 5 (Brouwer's Fixed Point Theorem):** Any continuous map $f : \mathbb{D}^2 \rightarrow \mathbb{D}^2$ has a fixed point, i.e. there exists $x \in \mathbb{D}^2$ such that $f(x) = x$.
- **Theorem 6 (Fundamental Theorem of Algebra):** Every non-constant complex polynomial $p \in \mathbb{C}[z]$ has at least one root, i.e. $p(z_0) = 0$ for some z_0 .
- **Theorem 7:** The fundamental group of S^n is trivial for $n \geq 2$, i.e. $\pi_1(S^2) \cong 1$ for $n \geq 2$

Lemma 7: Closed Paths Homotopic to Loops

Let (X, x_0) be a topological space with an open cover $\{U_j\}_{j \in J}$ such that U_j are path-connected neighbourhoods of x_0 and $U_j \cap U_{j'}$ is path-connected for any $j, j' \in J$. Then, any closed path γ based at x_0 is homotopic to a concatenation $\gamma_1 * \gamma_2 * \dots * \gamma_n$ of loops at x_0 each of them contained in a single U_j .

Corollary 3: Homomorphisms between \mathbb{R}^2 and \mathbb{R}^n

There is no homeomorphism between \mathbb{R}^2 and \mathbb{R}^n for $n \neq 2$.

Recall 4: Defining the Real Projective Space

1. The space \mathbb{RP}^2 is the quotient space:

$$\mathbb{RP}^n := (\mathbb{R}^{n+1} \setminus \{0\}) / \sim$$

where $x \sim y$ if there exists $\lambda \in \mathbb{R}$ s.t. $x = \lambda y$. i.e., the real projective n -space represents the lines in \mathbb{R}^{n+1} through the origin.

2. Picking representatives that lie in the unit n -sphere $S^n \subset \mathbb{R}^{n+1} \setminus \{0\}$, we obtain $\mathbb{RP}^n \cong S^n / \sim$ where $x \sim -x$ for all $x \in S^n$, i.e. identifying antipodal points on the n -sphere.
3. Further restricting to the upper half $\mathbb{D}^n \subset S^n$ we obtain:

$$\mathbb{RP}^n \cong \mathbb{D}^n / \sim$$

where $x \sim -x$ for any boundary points $x \in \partial \mathbb{D}^n \cong S^{n-1}$

For example, \mathbb{RP}^0 is a one point space, $\mathbb{RP}^1 \cong S^1$, while \mathbb{RP}^n are different than spheres for larger n .

Definition 20: Lift of a Path

- A lift of a path $\alpha : I \rightarrow \mathbb{RP}^n$ is a path $\tilde{\alpha} : I \rightarrow S^n$ s.t. $p \circ \tilde{\alpha} = \alpha$
- If α is a closed path, then $[\tilde{\alpha}(0)] = [\tilde{\alpha}(1)]$ which implies $\tilde{\alpha}(0) = \pm \tilde{\alpha}(1)$. The **sign** of α is defined by

$$\text{sgn}(\alpha) \mapsto \begin{cases} +1 & \tilde{\alpha}(0) = \tilde{\alpha}(1) \\ -1 & \tilde{\alpha}(0) = -\tilde{\alpha}(1) \end{cases}$$

Theorem 8: The Fundamental Group of \mathbb{RP}^2

The sign induces a surjective group homomorphism $\text{sgn} : \pi_1(\mathbb{RP}^n) \rightarrow \mathbb{Z}_2$, $[\alpha] \mapsto \text{sgn}(\alpha)$ which is an isomorphism for $n \geq 2$.

3 Covering Theory

Definition 21: Right Lifting Property

A map $p : X \rightarrow Y$ satisfies the **right lifting property** (RLP) w.r.t. a map $i : A \rightarrow B$ if any commutative square has a solution to the lifting problem making both triangles commute.

$$\begin{array}{ccc} A & \xrightarrow{\quad} & X \\ \downarrow i & \nearrow \exists! & \downarrow p \\ B & \xrightarrow{\quad} & Y \end{array}$$

Explicitly, if $f : B \rightarrow Y$ and $g : A \rightarrow X$ such that $f \circ i = p \circ g$, then there exists a map $l : B \rightarrow X$ satisfying $l \circ i = g$ and $p \circ l = f$. Dually, the map $i : A \rightarrow B$ is said to satisfy the **left lifting property** (LLP) with respect to $p : X \rightarrow Y$.

Example 13: Homotopy Lifting Property WRT Spaces

1. A map $p : X \rightarrow Y$ satisfies the **homotopy lifting property** w.r.t. a space Z iff it has the RLP with respect to the inclusion map $i : Z \times \{0\} \hookrightarrow Z \times I$, i.e. solves the following lifting problem:

$$\begin{array}{ccc} Z \times \{0\} & \xrightarrow{\quad} & X \\ \downarrow & \nearrow \exists & \downarrow p \\ Z \times I & \xrightarrow{\quad} & Y \end{array}$$

In other words, given a homotopy $h : Z \times I \rightarrow Y$ and a lift $\tilde{f} : Z \rightarrow X$ of $h(-, 0) =: f$, there is a homotopy lift $\tilde{h} : Z \times I \rightarrow X$ with $\tilde{h}(-, 0) = \tilde{f}$.

2. Dually, a map $i : A \rightarrow b$ satisfies the **homotopy extension property** (HEP) with w.r.t. a space Z iff it has the LLP w.r.t. the map

$$p : Z^I \rightarrow Z, \quad \gamma \mapsto \gamma(0)$$

$$\begin{array}{ccc} A & \xrightarrow{\quad} & Z^I \\ \downarrow i & \nearrow \exists & \downarrow p \\ B & \xrightarrow{\quad} & Z \end{array}$$

Where $Z^I := \text{Map}(I, Z)$ is the space of paths in Z . In other words, one can solve the following lifting problem.

Note that a map $A \rightarrow Z^I$ is the same datum as a homotopy $h : A \times I \rightarrow Z$. Given an extension $\tilde{f} : B \rightarrow Z$ of $h(-, 0)$ along i , the existence of a map $B \rightarrow Z^I$ which makes both triangles commute provides an extension of the homotopy h to a homotopy $\tilde{h} : B \times I \rightarrow Z$ along i .

Example 15: Covering Spaces

1. The projection map $p : X \times D \rightarrow X$ where D is a discrete space. Note that $X \times D$ cannot be path-connected unless D is a one-point set.
2. The covering map $\mathbb{R} \rightarrow S^1, t \mapsto e^{2\pi i t}$ which we can use to compute the fundamental group of S^1 .
3. The degree- n map $F_n : S^1 \rightarrow S^1, z \mapsto z^n$ provides an n -fold covering of S^1 by itself.
4. The product of two covering maps $p_i : \tilde{X}_i \rightarrow X_i$ is also a covering map $p_1 \times p_2 : \tilde{X}_1 \times \tilde{X}_2 \rightarrow X_1 \times X_2$
5. The product of F_n and F_m in the third example also provides a self covering of the torus:

$$T^2 = S^1 \times S^1 \rightarrow T^2, \quad (z, w) \mapsto (z^n, w^m)$$
6. Similarly, there is a covering $\mathbb{R}^2 \rightarrow T^2$.
7. The 2-fold covering $S^n \rightarrow \mathbb{RP}^n$ which was used to compute the fundamental group of \mathbb{RP}^n

Theorem 10: Homomorphism of Covering Maps

Let $p : \tilde{X} \rightarrow X$ be a covering map. The induced group homomorphism

$$p_* : \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$$

is injective for any $\tilde{x}_0 \in p^{-1}(x_0)$. Its image consists of (classes of) loops in X based at x_0 that lift to loops in \tilde{X} based at \tilde{x}_0

Remark 5: Notation for Covers

Fix a covering map $p : \tilde{X} \rightarrow X$ and $x_0 \in X$ a fixed point. Write $G := \pi_1(X, x_0)$ for the fundamental group of X at x_0 and $H := p_*\pi_1(\tilde{X}, \tilde{x}_0) \subset G$ for the subgroup determined by the covering map.

The subgroup H depends on the choice of fiber point $\tilde{x}_0 \in p^{-1}(x_0)$, and we shall see that its subgroups for different fiber points are conjugate to each other. Finally, the fiber over x_0 will be denoted by

$$F_{x_0} := p^{-1}(x_0)$$

Lemma 8: Transitive Actions

If \tilde{X} is path-connected, then the G -action on F_{x_0} is transitive, i.e. for any $\tilde{x}, \tilde{x}' \in F_{x_0}$, there exists an $\alpha \in G$ such that $\tilde{x} \cdot \alpha = \tilde{x}'$.

Theorem 11: Path-Connected Correspondence

If \tilde{X} is path-connected, then there is a one-to-one correspondence between right cosets and fiber points, i.e. a bijection

$$G\tilde{x} \backslash G \rightarrow F_{x_0}, \quad G\tilde{x} \cdot g \mapsto \tilde{x} \cdot g$$

Thus the index of $G_{\tilde{x}}$ in G coincides with the cardinality of the fiber F_{x_0} :

$$[G : G_{\tilde{x}}] = |F_{x_0}| \quad (20)$$

Corollary 4: If \tilde{X} is simply-connected, then there is a bijection

$$G \rightarrow F_{x_0}$$

Equation (20) becomes

$$|G| = |F_{x_0}| \quad (21)$$

4 Deck Transformations, Further Cover Theory

Definition 23: Deck Transformation

A **deck transformation** of a covering map $p : \tilde{X} \rightarrow X$ is a self-homeomorphism $D : \tilde{X} \xrightarrow{\cong} \tilde{X}$ such that $p \circ D = p$.

Deck transformations form a group $\text{Deck}(p)$. For any two deck transformations D, D' their composite is also a deck transformation since $p \circ D \circ D' = p \circ D' = p$. If D is a deck transformation then so is its inverse D^{-1} as $p \circ D^{-1} = p \circ D \circ D' = p$. For example, deck transformations of the covering map $\mathbb{R} \rightarrow \mathbb{S}^1$ are precisely translations by integers:

$$D_n : \mathbb{R} \rightarrow \mathbb{R}, \quad t \mapsto t + n$$


In particular, the group of deck transformations is \mathbb{Z} . More generally, the group of deck transformations $\text{Deck}(p)$ of a universal covering p is isomorphic to the fundamental group G . From now on $p : \tilde{X} \rightarrow X$ will be a covering with \tilde{X} path-connected and X path-connected and locally path-connected.

Example 16: Topologist's Sine Curve

Recall that the topologist's sine curve

$$X = \{0\} \times [-1, 1] \cup \left\{ \left(x, \sin\left(\frac{1}{x}\right) \right) \mid 0 < x \leq \frac{1}{2\pi} \right\} \subset \mathbb{R}^2$$

is an example of a connected, but not path-connected. Let Z be the quotient of X by identifying the points $(0, 0) \sim (\frac{1}{2\pi}, 0)$. Z is a path-connected space but not locally path-connected.



Theorem 12: Solutions to the Lifting Problem

Let $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ be a covering with X path-connected and locally path-connected, and let $g : (Z, z_0) \rightarrow (X, x_0)$ be a pointed map. Then, there exists a solution to the lifting problem

$$\begin{array}{ccc} & (\tilde{X}, \tilde{x}_0) & \\ \exists \tilde{g} \nearrow & \downarrow p & \\ (Z, z_0) & \xrightarrow{g} & (X, x_0) \end{array} \quad \text{iff } g_*\pi_1(Z, z_0) \subset p_*\pi_1(\tilde{X}, \tilde{x}_0)$$

If a solution to the lifting problem exists, then it is unique.

Remark 10: If g is a covering map, then so is its lift \tilde{g} . In particular, homomorphisms of covering maps are also covering maps.

Corollary 5: Universal Coverings

Let $p : \tilde{X} \rightarrow X$ be a covering with \tilde{X} simply connected. Then, for any covering $p' : X' \rightarrow X$ there exists a covering $\tilde{p} : \tilde{X} \rightarrow X'$ such that the following diagram commutes:

$$\begin{array}{ccc} & X' & \\ \tilde{p} \nearrow & \downarrow p' & \\ \tilde{X} & \xrightarrow{p} & X \end{array}$$

In other words, if a simply connected covering \tilde{X} of X exists, then it covers all other possible coverings. This is why such a covering is called the **universal covering** of X . For example, we have seen \mathbb{R} as the universal covering of \mathbb{S}^1 or \mathbb{S}^n as the universal covering of \mathbb{RP}^n .

Theorem 24: Covering Isomorphism

Let $p : \tilde{X} \rightarrow X$ be a covering with X path-connected and locally path-connected. Let $H \subset \pi_1(X, x_0)$ denote the subgroup determined by the covering map. Then, there exists a group isomorphism:

$$\text{Deck}(p) \cong N(H)/H$$

where $N(H)$ denotes the normalizer.

Definition 24: Normal Coverings

A covering $p : \tilde{X} \rightarrow X$ is **normal** if the subgroup H is normal

Trivially, universal coverings are always normal. All the examples so far were normal since all the fundamental groups we have seen so far were abelian.

Corollary 6: Let \tilde{X} be simply-connected. Then

$$\text{Deck}(p) \cong \pi_1(X, x_0).$$

Example 19: The Figure Eight Space


Let X be the figure eight space, $X = \mathbb{S}^1 \vee \mathbb{S}^1$.

Consider an oriented bicolored graph \tilde{X} whose vertices are all 4-valent with one incoming edge of each color and one outgoing edge of each color. Bicolored means each edge is labelled by a or b .

Such a graph determines a covering map

$$p : \tilde{X} \rightarrow X$$

by sending all vertices to the unique vertex of the figure-eight graph and the edges are sent to one of the loops. A universal covering is obtained by the following graph:



Vertical edges are oriented upwards and labelled by b , horizontal edges are oriented to the right and labelled by a . Deck transformations are freely generated by either D_a or D_b , where D_a (resp. D_b) acts on the graph by shifting all edges once to the right, rescaling them appropriately. In other words..

Theorem 14: Fundamental Group of $\mathbb{S}^1 \vee \mathbb{S}^1$

The fundamental group of $\mathbb{S}^1 \vee \mathbb{S}^1$ is the free group generated by two elements, i.e.

$$\pi_1(\mathbb{S}^1 \vee \mathbb{S}^1) \cong \langle a, b \rangle$$

Example 20.1: Covering The Möbius Band

Consider $M : \mathbb{R} \times I / \sim$ where $(x, y) \sim (x+1, 1-y)$. We obtain the homotopy equivalence using covering theory. The quotient map

$$q : \mathbb{R} \times I \rightarrow M$$

is the universal covering, since $\mathbb{R} \times I$ is simply-connected. For some $n \in \mathbb{Z}$, let D_n be the deck transformation:

$$D_n : \mathbb{R} \times I \rightarrow \mathbb{R} \times I, \quad (x, y) \mapsto (x+n, y_n)$$

where $y_n = y$ if n is even and $y_n = 1-y$ for n odd. These are all deck transformations and $\text{Deck}(p)$ is generated by D_1 since

$$D_n = (D_1)^n$$

For odd n , there are n -fold self-coverings $M \rightarrow M$. For even n , there are n -fold coverings by the cylinder $S^1 \rightarrow I$.

Example 20.2: Covering the Klein Bottle

Consider $K = \mathbb{R}^2 / \sim$, where $(x, y) \sim (x+1, 1-y) \sim (x, y+1)$ for all $(x, y) \in \mathbb{R}^2$. The quotient map of the Klein bottle

$$q : \mathbb{R}^2 \rightarrow K$$

is the universal covering map. Consider the deck transformation

$$D_a : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad (x, y) \mapsto (x, y+1)$$

and

$$D_b : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad (x, y) \mapsto (x+1, 1-y)$$

These two deck transformations generate the deck transformation group $\text{Deck}(q)$ and satisfy the relation:

$$D_b \circ D_a \circ D_b^{-1} \circ D_a = \text{id}.$$

Proposition 10: Fundamental Group of the Klein Bottle

The fundamental group of the Klein Bottle is:

$$\pi_1(K) = \langle a, b \rangle / \langle aba^{-1}b \rangle$$

5 Free Space and Examples

5.1.1 Indecipherable proof of \mathbb{S}^1

WTS: $\pi_1(\mathbb{S}^1) \cong \mathbb{Z}$, and generated by $\alpha : I \rightarrow \mathbb{S}^1, t \mapsto 2^{2\pi it}$. Let $\alpha_n : I \rightarrow \mathbb{S}^1, t \mapsto e^{2\pi int}$ be the loop based at 1 and wraps n times around \mathbb{S}^1 . The map

$$f : \mathbb{Z} \rightarrow \pi_1(\mathbb{S}^1, 1), \quad n \mapsto [\alpha_n]$$

is a group iso. It is a homo. as $f(0) = [\alpha_0] = [c_1], c_1$ constant path at 1 and $f(n+m) = f(n) \cdot f(m)$ follows from $a_{n+m} \simeq \alpha_n * \alpha_m$. Let $\gamma : I \rightarrow \mathbb{S}^1$ a loop based at 1 $\in \mathbb{S}^1$. $\exists n \in \mathbb{Z}$ s.t. $\gamma \simeq \alpha_n$, i.e. $[a] = [\alpha_n]$. Consider the covering map

$$p : \mathbb{R} \rightarrow \mathbb{S}^1, \quad t \mapsto e^{2\pi it}$$

and note the fiber of $1 \in \mathbb{S}^1$ is $p^{-1}(1) = \mathbb{Z} \subset \mathbb{R}$. By Corollary 2, $\exists \tilde{\gamma} : I \rightarrow \mathbb{R}$ s.t. $\tilde{\gamma}(0) = 0$ and $p \circ \tilde{\gamma} = \gamma$. In particular, $p \circ \tilde{\gamma} = \gamma$ and $p \circ \tilde{\gamma}(1) = \gamma(1) = 1$, therefore $\tilde{\gamma}(1) =: n$ is some integer. Similarly

$$\tilde{\alpha}_n : I \rightarrow \mathbb{R}, \quad t \mapsto nt$$

is a lift of α_n and also starts at 0, ends n . By contract. of \mathbb{R} , both $\tilde{\gamma}$ and $\tilde{\alpha}_n$ are homotopic as paths in \mathbb{R} . In particular, $p \circ \tilde{g} = \gamma$ and $p \circ \tilde{\alpha}_n = \alpha_n$ homotopic. Show that n is uniquely determined, i.e. $[\alpha_n] = [\alpha_m]$ iff $n = m$. Suppose \exists homotopy $h : I \times I \rightarrow \mathbb{S}^1$ paths $h(-, 0) = \alpha_n$ and $h(-, 1) = \alpha_m$. By crl2 \exists rel. homotopy $\tilde{h} : I \times I \rightarrow \mathbb{R}$ s.t. $\tilde{h}(-, 0) = \tilde{\alpha}_n$. In particular, we get a lift $\tilde{h}(-, 1)$ of α_m which by uniqueness must be $\tilde{h}(-, 1) = \tilde{\alpha}_m$. Comparing end points of $\tilde{\alpha}_n$ and $\tilde{\alpha}_m$ and noting \tilde{h} homotopy of paths, $n = m$ We use: **Covering space** $p : \mathbb{R} \rightarrow \mathbb{S}^1, t \mapsto e^{2\pi it}$ and its HLP to lift a closed path $\alpha : I \rightarrow \mathbb{S}^1$ at 1 to a path $\tilde{\alpha} : I \rightarrow \mathbb{R}$. **Angle map** at α : $\alpha(t) = e^{2\pi i \tilde{\alpha}(t)}$. If $\tilde{\alpha}$ and α' are both angle maps of α s.t. $\tilde{\alpha}(0) = \alpha'(0)$, then $\tilde{\alpha} = \alpha'$, i.e. unique up to starting point. **Degree** of α : $\deg(\alpha) := \tilde{\alpha}(1) - \tilde{\alpha}(0)$ and is indep. of choice of angle map. Thm 4 states that $\pi_1(\mathbb{S}^1) \rightarrow \mathbb{Z}, [\alpha] \mapsto \deg(\alpha)$ is a grp. iso.

5.1.2 Proof of results of \mathbb{S}^1

Brouwer: Suppose \exists cts. map $f : \mathbb{D}^2 \rightarrow \mathbb{D}^2$ s.t. $f(x) \neq x \forall x \in \mathbb{D}^2$. Consider the half line $L_x = \{tx + (1-t)f(x) \mid t \in \mathbb{R}_{>0}\}$ starts at $f(x)$, passes thru x and defines $r(x) \in \mathbb{S}^1$ by the intersection $r(x) := (L_x \setminus \{f(x)\}) \cap \mathbb{S}^1$ This defines a retract of \mathbb{D}^2 onto $\mathbb{S}^1, r : \mathbb{D}^2 \rightarrow \mathbb{S}^1, x \mapsto r(x)$ cts by cts of $f(x)$. However, the existence of the retract implies we have a surj. grp.homo. $r_* : \pi_1(\mathbb{D}^2) \rightarrow \pi_1(\mathbb{S}^1)$ which is a contradiction since $\pi_1(\mathbb{D}^2)$ trivial while $\pi_1(\mathbb{S}^1)$ non-trivial.

FTA: Suppose $p(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + c_0$ noncon. poly. $a_n \neq 0, n \geq 1$ w/ no roots, i.e. $p(z) \neq 0 \forall z \in \mathbb{C}$. WLOG Take $a_n = 1$ as p/a_n also sats. Consider

$$h : \mathbb{S}^1 \times I \rightarrow \mathbb{S}^1, (z, t) \mapsto \frac{t^n p((1-t)z/t)}{|t^n p((1-t)z/t)|}$$

well defined cts as p has no roots and

$$t^n p((1-t)z/t) = (1-t)^n z^n + (1-t)^{n-1} t a_{n-1} z^{n-1} + \dots + t^n a_0$$

In particular, $h(z, 0) = z^n$ and $h(z, 1) = \frac{a_0}{|a_0|}$. Hence, h defines a homotopy between the degree n map $\mathbb{S}^1 \times \mathbb{S}^1, z \mapsto z^n$ and the constant map $\mathbb{S}^1 \times \mathbb{S}^1, z \mapsto \frac{a_0}{|a_0|}$, contradiction.

High n -sphere: Let $\mathbb{S}^n \subset \mathbb{R}^{n+1}$ unit sphere, consider north pole $x_1 = (0, \dots, 0, 1)$ and south pole $x_2 = (0, \dots, 0, -1)$. Their associated complements $U_i = \mathbb{S}^n \setminus \{x_i\}$ are homeomorphic to \mathbb{R}^n and cover \mathbb{S}^n . Their intersection $U_1 \cap U_2$ homeomorphic to $\mathbb{S}^{n-1} \times \mathbb{R}$ and thus path connected for $n \neq 2$. By lemma 7, the class of a closed loop γ in \mathbb{S}^2 is the product of $[\gamma_1]$ and $[\gamma_2]$ each of the representatives contained in U_i . By contradictability of U_i we get $[\gamma]$ trivial. \mathbb{R}^2 and \mathbb{R}^n : Suppose $\exists f : R_2 \xrightarrow{\cong} \mathbb{R}^n$.

In particular we get a homeomorphism $f : \mathbb{R}^2 \setminus \{0\} \xrightarrow{\cong} \mathbb{R}^n \setminus \{f(0)\}$ which gives rise to a homotopy equivalence $S_1 \simeq \mathbb{R}^2 \setminus \{0\} \simeq^f \mathbb{R}^n \setminus \{f(0)\} \simeq \mathbb{S}^{n-1}$. Homotopy eqs induce a bij. of path-conn components $\pi_0(\mathbb{S}^1) \cong \pi_0(\mathbb{S}^{n-1})$ which leads to contradiction for $n = 1$ and they induce a grp iso. $\pi_1(\mathbb{S}^1) \cong \pi_1(\mathbb{S}^{n-1})$ contradiction for $n \geq 2$.

5.1.3 Real Projective Space

Recall Def20: A lift of a path $\alpha : I \rightarrow \mathbb{RP}^n$ is a path $\tilde{\alpha} II \rightarrow \mathbb{S}^n$ s.t. $p \circ \tilde{\alpha} = \alpha$. If α is a closed path, then $[\tilde{\alpha}(0)] = [\tilde{\alpha}(1)]$ which implies $\tilde{\alpha}(0) = \pm \tilde{\alpha}(1)$. The **sign** of α is defined by

$$\text{sgn}(\alpha) \mapsto \begin{cases} +1 & \tilde{\alpha}(0) = \tilde{\alpha}(1) \\ -1 & \tilde{\alpha}(0) = -\tilde{\alpha}(1) \end{cases}$$

The sign is indep. of the choice of lift: The fiber of $\alpha(0)$ consists of two antipod. points $x, -x$ in the n -sphere. By uniq. of lifts, $\exists!$ lift $\tilde{\alpha}$ starting at $\tilde{\alpha}(0) = x$. Such a lift also dets. a uniq. lift starting at antipod. point $-x$ by its antipod path $t \mapsto -\tilde{\alpha}(t)$. Both lifts give the same sign for α . In partic. a loop α has trivial sign iff its lift is a closed loop in \mathbb{S}^n . Example of loop with nontriv sign: For $n < m$, the inclusion $\mathbb{R}^{n+1} \hookrightarrow \mathbb{R}^{m+1}, (x_1, \dots, x_{n+1}) \mapsto (x_1, \dots, x_{n+1}, 0, \dots, 0)$ induces an inclusion of proj. spaces:

$$i_{n,m} : \mathbb{RP}^n \hookrightarrow \mathbb{RP}^m, \quad [x_1 : \dots : x_{n+1}] \mapsto [x_1 : \dots : x_{n+1} : 0 : \dots : 0]$$

The square root loop in \mathbb{RP}^n is the comp. $\mathbb{S}^1 \rightarrow \mathbb{RP}^1, z \mapsto \sqrt{z}$ with $i_{1,n} : \mathbb{RP}^1 \hookrightarrow \mathbb{RP}^n$. Explicitly, $\sigma : I \rightarrow \mathbb{RP}^n, t \mapsto [\cos(\pi t) : \sin(\pi t) : 0 : \dots : 0]$ and its sign is $\text{sgn}(\alpha) = -1$.

Proof: Let $h : \alpha \simeq \beta$ be a rel. homop. of loops in \mathbb{RP}^n and fix a lift $\tilde{\alpha}$ of α . Then $\exists!$ rel. homotop. lift $\tilde{h} : \tilde{\alpha} \simeq \tilde{\beta}$ of h . In partic. $\tilde{\alpha}(0) = \tilde{\beta}(0)$ and $\tilde{\alpha}(1) = \tilde{\beta}(1)$ which

implies $\text{sgn}(\alpha) = \text{sgn}(\beta)$. Hence well def. **sgn grp homo:** Let $\alpha, \beta : I \rightarrow \mathbb{RP}^n$ closed loops based at $[x]$ and fix lifts $\tilde{\alpha}, \tilde{\beta} : I \rightarrow \mathbb{S}^n$ s.t. $\tilde{\alpha}(1) = \tilde{\beta}(0)$. This is possible from ext. of lifts. In partic, $\tilde{\alpha} * \tilde{\beta}$ is a lift of $\alpha * \beta$. Via lifts, $\text{sgn}(\alpha * \beta) = \text{sgn}(\alpha) \text{sgn}(\beta)$. Surj. follows from \exists of sqrt loop $\alpha : I \rightarrow \mathbb{RP}^n$. Suppose $\alpha : I \rightarrow \mathbb{RP}^n$ loop with triv. sign. In partic, its lift $\tilde{\alpha}$ is a closed loop in \mathbb{S}^n . For $n \geq 2$, the n -sphere \mathbb{S}^n simply-connn and thus every loop in \mathbb{S}^n null homotop. Hence $\tilde{\alpha} \simeq c_x$ implying $\alpha \simeq [c_x]$, showing injectivity for $n \geq 2$.

5.1.4 Winding Number

The **winding number** $W(\omega)$ of a loop $\omega : \mathbb{S}^1 \rightarrow \mathbb{C} \setminus \{0\}$ is defined as the degree of $r \circ \omega : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ where $r : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{S}^1, z \mapsto \frac{z}{|z|}$ is the deformation retract. $\omega_1(z) = \underline{z} : -1$ which follows from $e^{2\pi it} = e^{-2\pi it}$. $\omega_2(z) = z^{154} - \frac{3}{2}i : 0$ since the image ω_2 is contractible in $\mathbb{C} \setminus \{0\}$. It's the unit circle centered at $-\frac{3}{2}i$. $w_3(z) = 1/2z^6, -6. w_4 = (z^5 - 1/2)/z^6 : W(z^5 - 1/2) + W(z^{-6}) = 5 - 6 = -1$.

5.1.5 Angle Maps again

An angle map of $\alpha : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ is a lift $\theta : I \rightarrow \mathbb{R}$ along the covering map $p : \mathbb{R} \rightarrow \mathbb{S}^1, t \mapsto e^{2\pi it}$. We have used the identification $\mathbb{S}^1 \cong I/\partial I$, i.e. it satisfies $\alpha(t) = e^{2\pi i \theta(t)}$. The degree of the map α is defined as $\deg(\alpha) := \theta(1) - \theta(0)$. For loops $f, g : \mathbb{S}^1 \rightarrow \mathbb{S}^1$, $\deg(f) = n$, $\deg(g) = m$, then we have homotopies $f \simeq F_n, g \simeq F_m$ where $F_n : \mathbb{S}^1 \rightarrow \mathbb{S}^1, z \mapsto z^n$. $f \circ g \simeq F_n \circ F_m = F_{nm} = \deg(f) \cdot \deg(g)$ and $f \cdot g \simeq F_n \cdot F_m = F_{n+m} = \deg(f) + \deg(g)$.

5.1.6 Isomorphism

Let X, Y be topological spaces with fixed points $x \in X$ and $y \in Y$. Show there is a group iso. $\pi_1(X \times Y, (x, y)) \cong \pi_1(X, x) \times \pi_1(Y, y)$. Denote $p_X : X \times Y \rightarrow X$ and $p_Y : X \times Y \rightarrow Y$ proj. maps. Consider $\Phi := (p_X)_* \times (p_Y)_* : \pi_1(X \times Y) \rightarrow \pi_1(X) \times \pi_1(Y)$

$$[\alpha] \mapsto ([p_Y \circ \alpha], [p_X \circ \alpha])$$

This is a grp. homo, as a product of grp. homos (proj. maps are cts and therefore w.def grp. homos.) Its inverse

$$\Psi : \pi_1(X) \times \pi_1(Y) \rightarrow \pi_1(X \times Y); ([a], [b]) \mapsto (\alpha \times \beta) \circ \Delta$$

where $\Delta : I \rightarrow I \times I, t \mapsto (t, t)$ diag. map. i.e. $(\alpha \times \beta)(t) = (\alpha(t), \beta(t))$ is a path in $X \times Y$ (as comp. of cts maps.). Ψ w.def: Let $h : \alpha \simeq \alpha'$ homotop. loops in X (based x) and $k : \beta \simeq \beta'$ homot. loops in Y (based y). Then we define homotop. $h \times k$ as the composite

$$H : X \times Y \times I \xrightarrow{\text{id} \times \Delta} X \times Y \times I \times I \xrightarrow{h \times k} X \times Y$$

or, $H((a, b), t) = (h(a, t), k(b, t))$. cts. as a comp. of cts. maps, and homotop. between $(\alpha \times \beta) \circ \Delta$ and $(\alpha' \times \beta') \circ \Delta$. Ψ is a grp. homo.: we have following eqs for paths: $c_{(x,y)} = (c_x, c_y) \circ \Delta : I \rightarrow X \times Y$ and $((\alpha * \alpha') \times (\beta * \beta')) \circ \Delta = ((\alpha \times \beta) \circ \Delta) * ((\alpha' \times \beta') \circ \Delta)$. Finally, Φ and Ψ are inverses, since for any map $f : Z \rightarrow X \times Y$ we have $f = (p_X \circ f, p_Y \circ f)$ and for paths $\alpha : I \rightarrow X$ and $\beta : I \rightarrow Y$ we have $p_X \circ (\alpha \times \beta) \circ \Delta = \alpha$ and $p_Y \circ (\alpha \times \beta) \circ \Delta = \beta$

5.1.7 Contractible Map

Let $f, g : X \rightarrow Y$ be any two cts. maps and Y contractible. Show $f \simeq g$ and give such a homotopy.

Y contractible, therefore $\text{id}_Y \simeq c_{y_0}$ for some $y_0 \in Y$ ($c : Y \rightarrow Y, y \mapsto y_0$). Let H be the homotopy between id_Y and c_{y_0} . Let $H' : X \times I \rightarrow Y, H'(x, t) = H(f(x), t)$ which is cts as the comp. of cts. maps. Therefore, we have $f \simeq c_{x_0}$. Similarly, $g \simeq c_{x_0}$ and therefore as eq. classes partition, $f \simeq g$. Homotopy between f and g : $h(x, t) = (1-t)f(x) + tg(x)$

5.1.8 Deformation Retract

Give a deformation retract of $\mathbb{D}^2 \setminus \{0\}$ onto \mathbb{S}^1 . Let $r : \mathbb{D}^2 \setminus \{0\} \rightarrow \mathbb{S}^1$ be defined by $r(x, t) = ((1-t) + t/\|x\|) \cdot x$. We have: $r(x, 0) = (1+0) \cdot x = x$, $r(x, 1) = (0 + x/\|x\|) = x/\|x\| \in \mathbb{S}^1, r(a, 1) = (0 + a/\|1\|) = a \in \mathbb{S}^1$

5.1.9 Deck transformations

Let $p : \tilde{X} \rightarrow X$ be a cover map. Let $D \in \text{Deck}(p), \tilde{x} \in p^{-1}(x_0)$ and $\alpha \in \pi_1(X, x_0)$. Show $D(\tilde{x} \cdot \alpha) = D(\tilde{x}) \cdot \alpha$ where we used the right $\pi_1(X, x_0)$ -action on the fiber $p^{-1}(x_0)$. **Sol:** Let γ be a loop at x_0 repr. $\alpha \in \pi_1(X, x_0)$ and let $\tilde{\gamma}$ be the uniq. path lift of γ in \tilde{X} that starts at \tilde{x} . By def. of the action, $\tilde{x} \cdot \alpha = \tilde{\gamma}(1)$. On the other hand, the path $D \circ \tilde{\gamma} : I \rightarrow \tilde{X}$ starts at $D(\tilde{x})$ and is also a lift of γ since $p \circ D \circ \tilde{\gamma} = p \circ \tilde{\gamma} = \gamma$. Therefore $D(\tilde{x}) \cdot \alpha = (D \circ \tilde{\gamma})(1) = D(\tilde{\gamma}(1)) = D(\tilde{x} \cdot \alpha)$

5.1.10 Trivial RP loops

The loop $\alpha_1(t) = [\cos(2\pi t) : \sin(2\pi t) : 0]$ has trivial class as its lift is closed in \mathbb{S}^2 . The loops $[\cos(\pi t) : \sin(\pi t) : 0], [2t - 1 : t^3 - t^5 : 1 - 2t^4]$ have antipodal start and end points and thus have nontrivial classes. In general, $[\cos(\lambda t), \sin(\lambda t), 0]$ closed path if $\alpha(0) = [\cos(\lambda 0), \sin(\lambda 0), 0] = [1, 0, 0] = \alpha(1) = [\cos(\lambda), \sin(\lambda), 0]$ i.e. $\sin(\lambda) = 0, \cos(\lambda) = \pm 1$, i.e. $\lambda = \pi \mathbb{Z}$.

5.1.11 Classification of Covering spaces

Covering spaces of (X, x_0) are (up to homeo. of cover spaces) 1-to-1 correspondence with subgroups of $\pi_1(X, x_0)$. Using that determine all connected covering spaces. \mathbb{S}^1 : All subgroups of \mathbb{Z} are $n\mathbb{Z} \subset \mathbb{Z}$ for $n \in \mathbb{Z}$. In particular, we have to determine coverings inducing those subgroups, these are $\mathbb{S}^1 \rightarrow \mathbb{S}^1, z \mapsto z^n$ for $n \neq 0$ and $R \rightarrow \mathbb{S}^1$ for $0 \subset \mathbb{Z}$. \mathbb{RP}^2 : The only subgroups of \mathbb{Z}_2 are the trivial and itself. The two assoc. coverings are $\mathbb{S}^2 \rightarrow \mathbb{RP}^2$ and the id. cover. T (2-torus): All subgroups are of the form $n\mathbb{Z} \times m\mathbb{Z}$ for $n, m \in \mathbb{Z}$. The assoc. coverings are products of the ones for \mathbb{S}^1 . M (Möbius band): FG is \mathbb{Z} as it is homotop. equiv. to its core. Hence we only need to find coverings for each $n\mathbb{Z} \subset \mathbb{Z}$. We start with its universal cover $\mathbb{R} \times [0, 1] \rightarrow M = \mathbb{R} \times [0, 1]/(x, y) \sim (x+1, 1-y)$. For each n , $\tilde{X} = \mathbb{R} \times [0, 1]/(x, y) \sim (x+n, y_n)$, where $y = y$ for n even and $y_n = 1-y$ for n odd, give all the covering spaces. In particular for even n we get n -fold coverings by $\mathbb{S}^1 \times [0, 1]$ and odd n we get n -fold coverings by Möbius bands.

5.1.12 More circles

Describe $f_* : \pi_1(\mathbb{S}^1 \times \mathbb{S}^1, (1, 1)) \rightarrow \pi_1(\mathbb{S}^1, 1)$ associated to the map $f : \mathbb{S}^1 \times \mathbb{S}^1 \rightarrow \mathbb{S}^1$ given by $f(w, z) = w^p z^q$ for $p, q \in \mathbb{Z}$. Let $\alpha \in \pi_1(T_2)$ be a single loop around the first circle: $t \mapsto (e^{2\pi it}, 1)$ and $\beta \in \pi_1(T_2)$ be a single loop around the second: $t \mapsto (1, e^{2\pi it})$. Then $\langle \alpha, \beta \rangle \cong \mathbb{Z} \times \mathbb{Z}$ generates $\pi_1(T^2)$. $f_*(\alpha)$ is the eq.class of $f(e^{2\pi it}, 1) \mapsto e^{2\pi itp} \cdot 1^q = e^{2\pi itp}$, or the loop that wraps \mathbb{S}^1 p times. $f_*(\beta)$ is the eq.class of $f(1, e^{2\pi it}) \mapsto 1^p \cdot e^{2\pi itq} = e^{2\pi itq}$, or the loop that wraps \mathbb{S}^1 q times. Therefore $f(w, z)$ is the homo. that wraps around $e^{2\pi it(p+q)}$, or $p+q$ times. As α and β are the generators, this means $f_*(m, n) = pm + qn$.

Compute the degree of cts $w_1 : \mathbb{S}^1 \rightarrow \mathbb{S}^1; (x, y) \mapsto (x, -y)$ - takes points on \mathbb{S}^1 and reflects on x -axis. Therefore, $(x, y) \mapsto (x, -y)$ corresponds to $z \mapsto \bar{z}$ on \mathbb{S}^1 , i.e. $\deg(w_1) = -1$ - a map that reverses orientation and wraps around the circle once. Compute the sign of the cts $w_3 : \mathbb{S}^1 \rightarrow \mathbb{RP}^2; (x, y) \mapsto [x : 0 : -y]$. This sends (x, y) to the projective point on the line spanned by $(x, 0, -y) \in \mathbb{R}^3$, i.e. image lies in $\{[x : 0 : z] \in \mathbb{RP}^2 \mid x^2 + z^2 \neq 0\} \cong \mathbb{S}^1 \subset \mathbb{RP}^2$ since the points $[x, 0, -y]$ form a circle in \mathbb{RP} when we vary (x, y) over the unit circle. Antipodal points $w_3(x, y) = [x : 0 : -y], w_3(-x, -y) = [-x : 0 : y] = [x : 0 : -y]$. Therefore w_3 factors through the quotient map $\mathbb{S}^1 \rightarrow \mathbb{RP}^1$, and the image is homeo to \mathbb{RP}^1 , therefore as it is nontrivial the sign is -1 "inclusion $\mathbb{RP}^1 \hookrightarrow \mathbb{RP}^2$ maps gen. of $\pi_1(\mathbb{RP}^1) \cong \mathbb{Z}$ to the nontriv. elem. of $\mathbb{Z}/2\mathbb{Z}$."

5.1.13 Retracts

Show that if Y is a retract of \mathbb{D}^2 then every map $g : Y \rightarrow Y$ has a fixed point (assume brouwer). If A is a retract of \mathbb{D}^2 then $\exists r : \mathbb{D}^2 \rightarrow A$ with $rj = 1_A$. jfr has a fixed point $x \in \mathbb{D}^2$. Now since $rjfr = (rf)fr = 1_A fr = fr$ then $f(r(x)) = fr(x) = rjfr(x) = r(jfr(x)) = r(x)$ and so f fixed $r(x)$.

Does the inclusion $i : \mathbb{RP}^1 \rightarrow \mathbb{RP}^2; [x : y] \mapsto [x : y : 0]$ admit a retraction? No. Consider the fundamental groups $\pi_1(\mathbb{RP}^1) \cong \mathbb{S}^1 \cong \mathbb{Z}$, and $\pi_1(\mathbb{RP}^2) \cong \mathbb{Z}/2\mathbb{Z}$, along with the induced inclusion map $i_* : \pi_1(\mathbb{RP}^1) \rightarrow \pi_1(\mathbb{RP}^2)$ which sends \mathbb{Z} to $\mathbb{Z}/2\mathbb{Z}$. If there is a retract, i.e. $r \circ i = \text{id}_{\mathbb{RP}^2}$ then we would need to have $r_* \circ i_* = \text{id}_{\pi_1(\mathbb{RP}^2)}$. However there is no such map $r_* : \mathbb{Z}/2\mathbb{Z} \rightarrow \mathbb{Z}$, therefore no such retraction can exist.


6 Unexamined Material

Definition 22: Fibration

1. A map $p : X \rightarrow Y$ is a **fibration** if it satisfies the HLP w.r.t. all spaces Z . i.e., it has the RLP w.r.t. the set of maps $\{i : Z \times \{0\} \hookrightarrow Z \times I\}_Z$ where Z runs over all topo. spaces.
2. Dually, a map $i : A \rightarrow B$ is a **cofibration** if it satisfies the HEP with respect to all spaces Z .


Theorem 9: Covering Maps are Fibrations

A covering map $p : \tilde{X} \rightarrow X$ is a fibration. Additionally, the homotopy lifts are unique:



Example 14: Examples of Fibrations

1. By Theorem 9, fibrations include all covering maps
2. The projection map $p : X \times F \rightarrow X$ is always a fibration. However this map is a covering map iff F is a discrete space. Hence, this includes examples of fibrations that are not coming from covering maps.
3. An important example of a cofibration is the inclusion $i : X \rightarrow M_f$ where M_f is the mapping cylinder of $f : X \rightarrow Y$. We have seen that any continuous map $f : X \times Y$ factors through the mapping cylinder:



In particular, every map factors through a cofibration and a homotopy equivalence.

Theorem 15: Seifert-Vam Kampen Theorem

Let X be a topological space with a fixed point x_0 . Let $\{U_\alpha\}_\alpha$ be an open cover of X consisting of path-connected open sets U_α containing the fixed point x_0 . The inclusions $U_\alpha \subset X$ induce a group homomorphism:

$$\Phi : *_\alpha \pi_1(U_\alpha) \rightarrow \pi_1(X).$$

1. If $U_\alpha \cap U_\beta$ is path-connected for any α, β , then Φ is surjective.
2. If $U_\alpha \cap U_\beta \cap U_\gamma$ is path-connected for any α, β, γ , then the kernel of Φ is generated by elements $i_{\alpha\beta}(w)i_{\beta\alpha}(w)^{-1}$ where $w \in \pi_1(U_\alpha \cap U_\beta)$ and $i_{\alpha\beta} : \pi_1(U_\alpha \cap U_\beta) \rightarrow \pi_1(U_\alpha)$ is the induced homomorphism from the inclusion $U_\alpha \cap U_\beta \subset U_\alpha$.

The assumption $U_\alpha \cap U_\beta$ are path-connected ensures that words $\pi_1(U_\alpha)$ generate $\pi_1(X)$. The assumption $U_\alpha \cap U_\beta \cap U_\gamma$ is path-connected gives a presentation for the group $\pi_1(X)$.

Example 17: Sifert-Vam Kampen on $\mathbb{S}^1 \vee \mathbb{S}^1$

Consider the figure eight $\mathbb{S}^1 \vee \mathbb{S}^1 = \mathbb{S}^1 \times \{1\} \cup \{1\} \times \mathbb{S}^1$ and let

$$U_i := \mathbb{S}^1 \vee \mathbb{S}^1 \setminus \{x_i\}$$

be the complements of the points $x_1 = (-1, 1)$ and $x_2 = (1, -1)$. The sets U_1 and U_2 are open path-connected and cover $\mathbb{S}^1 \vee \mathbb{S}^1$. In fact, they are both homotopy equivalent to the circle $U_i \simeq \mathbb{S}^1$. Their intersection $U_1 \cap U_2$ is contractible, and applying SVK we find,

$$\pi_1(\mathbb{S}^1 \vee \mathbb{S}^1) \cong \mathbb{Z} * \mathbb{Z}.$$

Example 18: Fundamental Group of Wedged Circles

Let (X_α, x_α) be a fIYL of path-connected pointed spaces and consider their wedge sum

$$X := \bigvee_{\alpha} X_{\alpha}.$$

suppose that each $x_\alpha := X_\alpha \vee \bigvee_{\beta \neq \alpha} U_\beta \subset X$. By contractibility of the U_α 's we have homotopy equivalences $A_\alpha \simeq X_\alpha$. Moreover, the intersection $A_\alpha \cap A_\beta$ is contractible for any $\alpha \neq \beta$. Applying SVK we obtain

$$\pi_1(X) \cong *_\alpha \pi_1(X_\alpha).$$

In particular, the fundamental group of the n -th wedge sum of circles is the free group on n -generators:

$$\pi_1\left(\bigwedge^n \mathbb{S}^1\right) \cong \mathbb{Z} * \dots * \mathbb{Z} \cong \langle \alpha_1, \dots, \alpha_n \rangle. \quad (23)$$

Definition 25: CW Complexes

A special class of topological spaces which are constructed inductively attaching n -dimensional disks or n -cells are called **CW complexes**. They are described as follows:

1. A set X^0 of **vertices** or 0-cells
2. Inductively construct the n -skeleton X^n from X^{n-1} by attaching n -dimensional disks \mathbb{D}_α^n by attaching maps $\phi_\alpha : \partial D_\alpha^n = \mathbb{S}_\alpha^{n-1} \rightarrow X^{n-1}$. In other words,

$$X^n = X^{n-1} \amalg_{\phi_\alpha} \coprod_{\alpha} D_\alpha^n.$$

Equivalently, a **CW Complex** is a space X along with a filtration of subspaces

$$X^0 \subset \dots \subset X^n \subset X^{n+1} \subset \dots \subset X$$

such that $X^n \setminus X^{n-1}$ is homeomorphic to a disjoint union of n -dimensional open disks, and X^0 is discrete.

Example 19: Examples of CW Complexes

1. The Torus $T^2 = I^2 / \sim$ can be made into a CW complex with: $X^0 = \{[(0, 0)]\}$, $X^1 = \{[(a, 0)] \mid a \in I\} \cup \{[(0, b)] \mid b \in I\}$ and $X^2 = T^2$. In particular, it has one 0-cell, two 1-cell, and one 2-cell.
2. The real projective plane \mathbb{RP}^2 can be made into a CW complex with $X^0 = *$, $X^1 = \mathbb{RP}^1 = \mathbb{S}^1$ and X^2 obtained by attaching a 2-disk to \mathbb{S}^1 along the quotient map $\mathbb{S}^1 \rightarrow \mathbb{RP}^1$