

Algebraic Topology Notes

Made by Leon :) *Note: Any reference numbers are to the lecture notes*

1 Introduction

Recall 1.1.1: Topology

An **(open) topology** on X is a collection of subsets $\tau \subset P(X)$ such that

- $\emptyset \in \tau$ and $X \in \tau$
- τ is closed under finite intersections: If $\{U_1, \dots, U_n\} \subset \tau$ then

$$\bigcap_{i=1, \dots, n} U_i \in \tau$$

- τ is closed under arbitrary unions: If $\{U_1, \dots, U_n\} \subset \tau$ is a family of open subsets then

$$\bigcup_{i=1, \dots, n} U_i \in \tau$$

The subsets $U \in \tau$ are called **open** and their complements in X define **closed subsets**.

Two examples of a topology on a set X are the following:

- The **Trivial Topology**: $\tau_{\text{triv}} = \{\emptyset, X\}$
- The **Discrete Topology**: $\tau_{\text{dis}} = P(X)$

A subset $A \subset X$ is **clopen** if it is both closed and open

Definition 1: Connected Spaces

A topological space X is **connected** if $X = A \cup B$ with $A, B \subset X$ open implies that $A = \emptyset$ or $A = X$.

Proposition 1: Connectedness and Clopens

A topological space X is *connected* iff the only clopens are \emptyset and X .

Example 1: Examples of Connected Topologies

- Every X with the trivial topology is connected.
- Every X with the discrete topology is not connected unless $X = \emptyset$ or $X = \{*\}$ (in which it coincides with the trivial topology).
- The real line \mathbb{R} with the standard topology is connected.

Proposition 2: Continuous Maps

Let $f : X \rightarrow Y$ be a continuous map of topological spaces and let X be connected. Then $f(X)$ is connected.

Proposition 3: Connected Equivalence Relation

For a topological space X , define $x \sim y$ if there exists some connected subset that contains both. The relation $x \sim y$ is an equivalence relation.

Definition 2: Connected Components

The equivalence classes of this relation are called **connected components**. In particular, a space X is connected iff it only has a single connected component.

Definition 3: Path

Let I denote the closed unit interval $[0, 1]$. A **path** in X is a continuous map $\alpha : I \rightarrow X$. The points $\alpha(0) \in X$ and $\alpha(1) \in X$ will be called **start** and **end** points respectively. We define a path relation between points in X by declaring $x \sim y$ if there exists some path $\alpha : I \rightarrow X$ that starts at x and ends in y , i.e. $\alpha(0) = x$ and $\alpha(1) = y$. This is an equivalence relation from the following properties:

1. **Constant Path**: For all $x \in X$ there exists the constant path $c_x : I \rightarrow X$ defined by $c_x(t) = x$ for all $t \in I$
2. **Path reversal**: Let $\alpha : I \rightarrow X$ be a path in X . Define its reversed path by

$$\bar{\alpha} : I \rightarrow X, \quad t \mapsto \alpha(1 - t) \quad (1)$$

3. **Path Concatenation**: Let $\alpha, \beta : I \rightarrow X$ be two paths in X s.t. $\alpha(1) = \beta(0)$. Their concatenated path is defined by:

$$\alpha * \beta(t) := \begin{cases} \alpha(2t), & 0 \leq t \leq 1/2 \\ \beta(2t - 1) & 1/2 \leq t \leq 1 \end{cases} \quad (2)$$

Definition 4: Path-Connected Components

The equivalence classes are called **path-connected components** and their set is denoted by $\pi_0(X)$. A space X is called **path-connected** if $\pi_0(X)$ is a one-point set, i.e. any two points x, y can be related by a path in X .

Remark 1: Random examples

The following statements are true:

- A homeomorphism $X \cong Y$ induces a bijection $\pi_0(X) \cong \pi_0(Y)$.
- If X is path-connected, it is also connected.
- The *topologist's sine curve* defined by $X = \{0\} \times [-1, 1] \times \{(x, \sin(1/x)) \mid 0 < x\}$ is connected but not path-connected.

Definition 5: Homotopy

A **homotopy** of maps $f, g : X \rightarrow Y$ is a continuous map $h : X \times I \rightarrow Y$ such that $h(-, 0) = f$ and $h(-, 1) = g$.

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \Downarrow h & \\ X & \xrightarrow{g} & Y \end{array} \quad (3)$$

If such a homotopy exists, f is called **homotopic** to g . This defines an equivalence relation $f \simeq g$ on the space of maps $\text{Map}(X, Y)$.

Example 2: Paths as Homotopies

Points in X are the same as maps $*$ \rightarrow X from the one-point set $*$ to X . A path $\alpha : I \rightarrow X$ corresponds to a homotopy $*$ \times $I \rightarrow X$.

Remark 1.5: Composition of Homotopies

- **Horizontal Composition**: Let $h, h' : X \times I \rightarrow Y$ be two homotopies in X such that $h(-, 1) = h'(-, 0) : X \rightarrow Y$. Their concatenated homotopy is defined by

$$h * h'(-, t) := \begin{cases} h(-, 2t) & 0 \leq t \leq 1/2 \\ h'(-, 2t - 1) & 1/2 \leq t \leq 1 \end{cases} \quad (5)$$

- **Vertical Composition**: Let $h : X \times I \rightarrow Y$ and $k : Y \times I \rightarrow Z$ be two homotopies on maps from X to Y , and Y to Z . Then

$$k \circ h := [X \times I \xrightarrow{\text{id} \times \Delta} X \times I^2 \xrightarrow{h \times \text{id}} Y \times I \xrightarrow{k} Z] \quad (6)$$

where $\Delta : I \rightarrow I^2$, $t \mapsto (t, t)$ is the diagonal map, or explicitly, $k \circ h(x, t) = k(h(x, t), t)$

$$\begin{array}{ccccc} & \begin{array}{c} f \\ \Downarrow h \\ \end{array} & & \begin{array}{c} f \\ \Downarrow k' * h \\ \end{array} & \\ X & \xrightarrow{\quad} & Y \sim X & \xrightarrow{\quad} & Y \\ & \begin{array}{c} \Downarrow h' \\ l \end{array} & & \begin{array}{c} \Downarrow l \end{array} & \\ & \begin{array}{c} f \\ \Downarrow h \\ \end{array} & & \begin{array}{c} f' \\ \Downarrow k \\ \end{array} & \\ X & \xrightarrow{\quad} & Y & \xrightarrow{\quad} & Z \sim X & \xrightarrow{\quad} & Z \\ & \begin{array}{c} \Downarrow h \\ g \end{array} & & \begin{array}{c} \Downarrow k \\ g' \end{array} & & \begin{array}{c} \Downarrow k \circ h \\ g' \circ g \end{array} & \end{array}$$

Lemma 1: Concatenation Relation

Let $f, f' : X \rightarrow Y$ and $g, g' : Y \rightarrow Z$ be maps such that $f \simeq f'$ and $g \simeq g'$. Then $g \circ f \simeq g' \circ f'$ as maps from X to Z . In particular, $g' \circ f \sim g \circ f$ and $g \circ f' \sim g \circ f$.

Definition 6: Homotopy Equivalence

A map $f : X \rightarrow Y$ is called a **homotopy equivalence** if there exists a map $g : Y \rightarrow X$ and homotopies $f \circ g \simeq \text{id}_Y$, $g \circ f \simeq \text{id}_X$. In other words, g satisfies the properties of an inverse up to homotopy. It is called a **homotopy inverse** of f .

Example 3: Circle to \mathbb{R}^2

The inclusion map $\mathbb{S}^1 \hookrightarrow \mathbb{R}^2$ is not a homotopy equivalence, but the inclusion $\mathbb{S}^1 \hookrightarrow \mathbb{R}^2 \setminus \{0\}$ is a homotopy equivalence.

Proposition 4: Unique Inverses of Homotopy

Homotopy inverses are unique up to homotopy.

Definition 7: Homotopic Spaces

Two spaces X and Y are called **homotopy equivalent**, or **of the same homotopy type**, and denoted by $X \simeq Y$ if there exists a homotopy equivalence $f : X \rightarrow Y$.

Notation: We use \cong for homeomorphisms and \simeq for homotopy equivalence.

Lemma 2: Composition of Inverses

Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ with homotopy inverses $\bar{f} : Y \rightarrow X$ and $\bar{g} : Z \rightarrow Y$ respectively. Then $\bar{f} \circ \bar{g} : Z \rightarrow X$ is a homotopy inverse of $g \circ f : X \rightarrow Z$. In particular, $X \simeq Y$ and $Y \simeq Z$ implies $X \simeq Z$.

Definition 8: Contractible Space

A space X is called **contractible** if it is homotopy equivalent to a point, i.e. $X \simeq *$.

The **terminal map** is the unique map $X \rightarrow *$. Contractibility requires that there is a homotopy inverse of that map, i.e. a map $* \rightarrow X$ along with homotopies

$$h : [* \rightarrow X \rightarrow *] \simeq \text{id}_*, \quad k : [X \rightarrow * \rightarrow X] \simeq \text{id}_X \quad (7)$$

Example 4: Examples of Contractible Spaces

- \mathbb{R}^n is contractible. Let x_0 be a fixed point in \mathbb{R}^n and define the (straight line) homotopy $h : c_{x_0} \simeq \text{id}_{\mathbb{R}^n}$ by

$$h(x, t) = (1 - t)x_0 + tx.$$

- $\mathbb{S}^{n-1} \simeq \mathbb{R}^n \setminus \{0\}$. The inclusion $\mathbb{S}^{n-1} \hookrightarrow \mathbb{R}^n \setminus \{0\}$ and the shrinking map

$$\mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{S}^{n-1}, \quad x \mapsto \frac{x}{|x|}$$

are homotopy inverses.

Remark 3: Remarks about Contractible Spaces

- Contractible spaces are path-connected. Let x_0 be the point where the space X contracts to. In particular, we are given with a homotopy $h : c_{x_0} \simeq \text{id}_X$. For any $x \in X$, the map $h(x, -) : I \rightarrow X$ defines a path from x_0 to x and thus every element $x \in X$ is path-connected to x_0 .
- The converse does not hold, for example $X = \mathbb{S}^1$.
- A contractible space X is contractible at any point x_0 . Since X is path-connected, a path from x to x' defines a homotopy $c_x \simeq c_{x'}$.
- Any two maps $f, g : X \rightarrow Y$ are homotopic if Y is contractible.

Definition 9: Retracts and Deformation Retracts

- A **retract** of X onto a subspace $A \subset X$ is a map $r : X \rightarrow A$ such that $r|_A = \text{id}_A$. Equivalently, this is a map $r : X \rightarrow X$ such that $r^2 = r$ and $r(X) = A$.
- A **deformation retract** of X onto A is the additional datum of a homotopy $h : \text{id}_X \simeq i \circ r$.

In other words, a deformation retract is a homotopy $h : X \times I \rightarrow X$ such that $h(x, 0) = x$ and $h(x, 1) \in A$ for all $x \in X$ and $h(a, 1) = a$ for all $a \in A$. Not all retracts can form deformation retracts. For instance, the retract X onto a point $\{x_0\}$ can be a deformation retract iff X is contractible.

This notion is called **weak** deformation retract. A **strong** deformation retract has the condition $h(a, t) = a$ for all $t \in I$, $a \in A$. i.e. Our notion of a (weak) deformation retract deforms X into A while allowing to deform A to do so, while a strong deformation retract deforms X into A while keeping A fixed at all times

Proposition 5: Deformation Retracts and Homotopies

A deformation retract of X onto A induces a homotopy equivalence $X \simeq A$.

Recall 2: Quotient Space

Let X be a topological space and let \sim be an equivalence relation on X . Then, X/\sim is equipped with the quotient topology and called a **quotient space**. If Z is a closed subset in X , then we can also define the quotient space X/Z .

Another form of quotient spaces: Let $f : Z \rightarrow Y$ be a continuous map between a closed subset $Z \subset X$ and Y . Then

$$X \cup_f Y = X \cup Y/z \simeq f(z).$$

Example 5: Examples of Quotient Spaces

- The quotient of the n -dimensional closed disk by its boundary is the n -sphere, i.e. $\mathbb{D}^n/\partial\mathbb{D}^n \cong \mathbb{S}^n$.
- The 2-torus: $\mathbb{R}^2/\mathbb{Z}^2$. The projective space:

$\mathbb{RP}^n = \mathbb{R}^{n+1} \setminus \{0\} / \sim$ by the relation $x \sim y$ iff there exists some $\lambda \in \mathbb{R}^\times$ such that $x = \lambda y$. This corresponds to the space of lines through the origin in \mathbb{R}^{n+1} .

Definition 10: Mapping Quotients

Let $f : X \rightarrow Y$ be a continuous map.

- Its **mapping cylinder** is defined as the topological space

$$M_f := (X \times I) \cup Y / \sim$$

where the quotient identifies $(x, 0) \sim f(x)$ for any $x \in X$.

- Its **cone** is the further quotient:

$$C_f = M_f / X \times \{1\}.$$

- The **cone** of a topological space X is

$$C_X := C_{\text{id}_X} = X \times I / X \times \{1\}.$$

In other words, the mapping cylinder of $f : X \times Y$ is the pushout of the diagram

$$\begin{array}{ccc} X \times \{0\} & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ X \times I & \longrightarrow & M_f \end{array}$$

Example 5.5: Spheres

Consider the n -sphere \mathbb{S}^n with the standard embedding $\mathbb{R}^{n+1} \setminus \{0\}$. Then the map

$$r : \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{S}^n, \quad x \mapsto \frac{x}{|x|}$$

is a retract. Indeed, if x has norm $|x| = 1$, then $r(x) = x$. For a deformation retract one needs to find a homotopy $h : i \circ r \simeq \text{id}_X$. This can easily be realised by following straight-line homotopy:

$$h : \mathbb{R}^{n+1} \setminus \{0\} \times I \rightarrow \mathbb{R}^{n+1} \setminus \{0\}, \quad (x, t) \mapsto (1 - t)\frac{x}{|x|} + tx.$$

Indeed, $h(x, 0) = r(x)$ and $h(x, 1) = x$ for all x .

Definition 11: Star-Shaped Spaces

A subset $S \subset \mathbb{R}^n$ is called **star-shaped** at a point $x_0 \in S$, if for any $x \in S$ the line segment from x_0 to x is contained in S , i.e.

$$\{(1 - t)x_0 + tx \mid t \in [0, 1]\} \subset S$$

If S is star-shaped at every point, then it is called **convex**.

Example 5.6: Star-Shaped Spaces are Contractible

Let S be star-shaped at x_0 and $i : \{x_0\} \hookrightarrow S : r$ be the inclusion and constant maps. Define the straight line homotopy

$$h : S \times I \rightarrow S, \quad (x, t) \mapsto (1 - t)x_0 + tx$$

which is well-defined by the star-shaped condition. Moreover, $h(x, 0) = x_0 = r(x)$ and $h(x, 1) = x$ for all x . Hence, star-shaped, and in particular convex spaces, are contractible.

Example 5.7: Möbius band

The Möbius band M can be defined as

$$M = I^2 / \sim$$

where the equivalence relation \sim identifies the two vertical edges of I^2 by flipping one, i.e. $(0, b) \sim (1, 1 - b)$ for $b \in I$. Its core $C \subset M$ is the line $\{[a, 1/2] \mid a \in I\}$. Thus, the core is homeomorphic to \mathbb{S}^1 .

The Möbius band deformation retracts onto its core. Indeed, consider the retract $r : M \rightarrow C$ defined by $r([a, b]) := [a, 1/2]$ and the homotopy

$$h : M \times I \rightarrow M, \quad ([a, b], t) \mapsto \left[\left(a, (1 - t)\frac{1}{2} + \right) \right].$$

In particular, $M \simeq \mathbb{S}^1$.

Proposition 6: Retracts of the Mapping Cylinder

Via Definition 1.1.10, the mapping cylinder is formed by the cylinder of X by gluing Y onto the bottom with the map f . The mapping cylinder M_f strongly deformation retracts onto Y .

Proof. Consider the retract:

$$r : M_f \rightarrow Y$$

defined by $r([x, s]) := [(x, 0)] = [f(x)]$ on the class of $(x, s) \in X \times I$ and $r([y]) = y$ for $y \in Y$. This is well-defined and by definition a retract on Y . Define the homotopy

$$h : M_f \times I \rightarrow M_f$$

by $h([(x, s)], t) := [(x, st)]$ for $(x, s) \in X \times I$ and $t \in I$, and by $h([y], t) := y$ for $y \in Y$. In particular, $h(-, 0) \circ i \circ r$ and $h(-, 1) = \text{id}_{M_f}$. This forms a strong deformation retract. \square

Remark 6: Continuous Maps are Homotopic

Any continuous $f : X \rightarrow Y$ can be replaced up to homotopy equivalence by the closed inclusion $X \hookrightarrow M_f$, $x \mapsto [(x, 1)]$. More precisely, it factorises through an inclusion and a homotopy equivalence as the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow & \nearrow \simeq \\ & M_f & \end{array}$$

Definition 12: Relative Homotopy

Let X, Y be topological spaces and $A \subset X$ be a subset in X . A homotopy $h : X \times I \rightarrow Y$ is called **relative to A** if $h(a, t)$ is independent of t for all $a \in A$. In particular, this defines homotopies

between maps $f, g : X \rightarrow Y$ such that $f|_A = g|_A$.

This definition generalises ordinary homotopies, as an ordinary homotopy is the same as a homotopy relative to \emptyset .

Example 6: Relative Homotopies and Retracts

A strong deformation retract of X onto A is a deformation retract such that the homotopy $h : i \circ r \simeq \text{id}_X$ is relative to A .

Definition 13: Homotopic Path

Let $\alpha, \beta : I \rightarrow X$ be paths in X such that $\alpha(0) = \beta(0)$ and $\alpha(1) = \beta(1)$. A relative homotopy from α to β is a homotopy $h : I \times I \rightarrow X$ relative to $\partial I = \{0, 1\}$, i.e.

$$h(-, 0) = \alpha, \quad h(-, 1) = \beta \quad (8)$$

and

$$h(0, t) = \alpha(0) = \beta(0), \quad h(1, t) = \alpha(1) = \beta(1), \quad \forall t \in I. \quad (9)$$

In particular, at any point $t \in I$ a relative homotopy h defines a path $h_t := h(-, t) : I \rightarrow X$ with start $\alpha(0) = \beta(0)$ and end $\alpha(1) = \beta(1)$. If one omits the relative condition, the start and end points of h_t would be allowed to vary.

Remark 7: Ordinary Homotopies and Paths

Observe that ordinary homotopies are not well suited for paths: Any path $\alpha : I \rightarrow X$ is homotopic (relative \emptyset) to a constant. Indeed, the homotopy

$$h : I \times I \rightarrow X, \quad (s, t) \mapsto \alpha(st)$$

defines a homotopy from the constant path $c_{\alpha(0)}$ on $\alpha(0)$ to α , i.e. $c_{\alpha(0)} \simeq \alpha$. Hence, (ordinary) homotopy classes of paths in X are in one-to-one correspondence with path-connected components of X .

Proposition 7: Homotopic Properties of Paths

Path concatenation is **unital**, **associative**, and **invertible** up to homotopy in the following sense: Let $\alpha, \beta, \gamma : I \rightarrow X$ be paths such that $\alpha(1) = \beta(0)$ and $\beta(1) = \gamma(0)$. Then there exists homotopies relative to $\{0, 1\}$:

1. **Left Unitality:** $c_{\alpha(0)} * \alpha \simeq \alpha$
2. **Right Unitality:** $\alpha \simeq c_{\alpha(0)} * \alpha$
3. **Associativity:** $(\alpha * \beta) * \gamma \simeq \alpha * (\beta * \gamma)$
4. **Right Inverse:** $\alpha * \bar{\alpha} \simeq c_{\alpha(0)}$
5. **Left Inverse:** $\bar{\alpha} * \alpha \simeq c_{\alpha(1)}$

where c_x for some $x \in X$ denotes the constant path on x and $\bar{\alpha}$ is the reversed path.

Lemma 3:

Let $\alpha : I \rightarrow X$ be a path and $\lambda : I \rightarrow I$ a boundary preserving map, i.e. $\lambda(0) = 0$ and $\lambda(1) = 1$. Then,

$$\alpha \circ \lambda \simeq \alpha, \quad \text{rel. } \partial I.$$

Definition 14: Fundamental Group

Let X be a topological space and $x_0 \in X$ some fixed point. The **fundamental group** of X at x_0 is the group of homotopy classes of paths in X that start and end on x_0 . i.e. $\alpha : I \rightarrow X$ such that $\alpha(0) = \alpha(1) = x_0$, i.e.

$$\pi_1(X, x) = \{\alpha : I \rightarrow X \mid \alpha(0) = \alpha(1)\} / \sim.$$

Theorem 1: The Fundamental Group is Well Defined

The fundamental group $\pi_1(X, x_0)$ is a well-defined group with:

- **Multiplication:** $[\alpha] \cdot [\beta] := [\alpha * \beta]$
- **Unit:** $1 = [c_{x_0}]$
- **Inverse:** $[\alpha]^{-1} = [\bar{\alpha}]$

Lemma 4: Relative Concatenated Homotopic Paths

Let $\alpha \simeq \alpha' : I \rightarrow X$ and $\beta \simeq \beta' : I \rightarrow X$ be two pairs of relative homotopic paths such that $\alpha(1) = \alpha'(1) = \beta(0) = \beta'(0)$. Then the concatenations are relative homotopic, i.e.

$$\alpha * \beta \simeq \alpha' * \beta', \quad \text{rel. } \{0, 1\}.$$

Proposition 8: Fundamental Group is Point Independent

Let $\gamma : I \rightarrow X$ be a path from $\gamma(0) = x$ to $\gamma(1) = x'$. Then it induces a group isomorphism:

$$(\gamma)_\# : \pi(X, x) \rightarrow \pi(X, x'), \quad [\alpha] \mapsto [\bar{\gamma} * \alpha * \gamma].$$

We abuse notation to say that for a path-connected space X , $\pi_1(X)$ is the fundamental group omitting the choice of base point.

Example 7: Examples of Fundamental Groups

- **Euclidean:** $\pi_1(\mathbb{R}^n) \cong 1$.
- **Circle:** $\pi_1(\mathbb{S}_1) \cong \mathbb{Z}$.
- **n -Spheres:** $\pi_1(\mathbb{S}^n) \cong 1$ for $n \geq 2$.
- **Torus:** $\pi_1(\mathbb{S}^1 \times \mathbb{S}^1) \cong \mathbb{Z} \oplus \mathbb{Z}$.
- **Projective Spaces:** $\pi_1(\mathbb{RP}^n) \cong \mathbb{Z}_2$ for $n \geq 2$.

Definition 15: Pointed Space

- A **pointed space** is a pair (X, x) consisting of a topological space X and a point $x \in X$.
- A **map of pointed spaces** $f : (X, x) \rightarrow (Y, y)$ is a continu-

ous map $f : X \rightarrow Y$ such that $f(x) = y$.

- The **space of pointed maps** from (X, x) to (Y, y) is denoted by

$$\text{Map}_*((X, x), (Y, y)) \subset \text{Map}(X, Y).$$

Proposition 9: Point and Path Space Isomorphism

We have a group isomorphism:

$$\pi_1(X, x) \cong \pi_0(\Omega X).$$

Similarly, one can iteratively define the n -fold loop space

$$\Omega^n X := \Omega \Omega^{n-1} X = \Omega \cdots \Omega X$$

There is a homeomorphism

$$\Omega^n X \cong \text{Map}_*((\mathbb{S}^k, 1), (X, x))$$

Definition 16: n -th Homotopy Group

The n -th homotopy group $\pi_n(X, x)$ is defined by:

$$\pi_n(X, x) := \pi_0(\Omega^n X) \cong \pi_0(\text{Map}_*(\mathbb{S}^n, (X, x))).$$

Definition 17: Simply Connected Space

A path-connected space X is called **simply connected** if its fundamental group is trivial, i.e.

$$\pi_1(X) = 1$$

Some examples are \mathbb{R}^n , \mathbb{S}^n for $n > 1$, and some non-examples are \mathbb{S}^1 , $\mathbb{S}^1 \times \mathbb{S}^1$ and $\mathbb{R}\mathbb{P}^2$.

Theorem 2: Fundamental Group Isomorphism

Let $f : X \rightarrow Y$ be a homotopy equivalence and $x \in X$ an arbitrary base point. Then, the map:

$$f_* : \pi_1(X, x) \rightarrow \pi_1(Y, f(x)), \quad [\alpha] \mapsto [f \circ \alpha]$$

In particular, for homotopy equivalent spaces $X \simeq Y$ which are path-connected, we get $\pi_1(X) \cong \pi_1(Y)$.

A map of pointed spaces $f : (X, x) \rightarrow (Y, y)$ is a **homotopy equivalence of pointed spaces** or **homotopy equivalence relative $\{x\}$** if there exists a map of pointed spaces $g : (Y, y) \rightarrow (X, x)$ along with relative homotopies

$$h : f \circ g \simeq \text{id}_Y \quad \text{rel. } \{y\}$$

and

$$k : g \circ f \simeq \text{id}_X \quad \text{rel. } \{x\}$$

Example 9: Strong Deformation Retracts Homotopies

A strong deformation retract of X onto a subspace A gives a homotopy equivalence of pointed spaces $(x, a) \rightarrow (A, a)$ for any

choice of $a \in A$. In particular, a contractible space $X \simeq *$ determines a homotopy equivalence of pointed spaces $(X, x) \rightarrow *$ for any choice of base point x .

Lemma 5: Pointed Space Isomorphism

Let $f : (X, x) \rightarrow (Y, y)$ be a homotopy equivalence of pointed spaces. Then the map

$$f_* : \pi_1(X, x) \rightarrow \pi_1(Y, y)$$

is a group isomorphism.

Corollary 1: Let $r : X \rightarrow A$ be a strong deformation retract of X onto $A \subset X$. Then

$$\pi_1(X, a) \cong \pi_1(A, a)$$

for any $a \in A$. In particular, contractible spaces are simply connected.

Lemma 6: Identity Homomorphic Isomorphism

Let $f : X \rightarrow X$ be a continuous map which is homotopic to id_X . Then, the map

$$f_* : \pi_1(X, x_0) \rightarrow \pi_1(X, f(x_0))$$

is a group isomorphism for any choice of base point $x_0 \in X$.

Definition 18: Homotopy Lifting Property

A continuous map $p : E \rightarrow X$ satisfies the **homotopy lifting property** (HLP) with respect to a topological space Y if for any commuting diagram:

$$\begin{array}{ccc} Y \times \{0\} & \xrightarrow{H_0} & E \\ \downarrow & \nearrow \exists! & \downarrow p \\ Y \times I & \xrightarrow{h} & X \end{array}$$

There exists a map $H : Y \times I \rightarrow E$ such that both triangles commute, i.e. $H|_{Y \times \{0\}} = H_0$ and $p \circ H = h$.

In other words, the map $p : E \rightarrow X$ has the HLP if for any homotopy $h : Y \times I \rightarrow X$ of maps $h(-, 0) := f_0$ and $h(-, 1) := f_1$ of maps $Y \rightarrow X$ and a choice of a lift H_0 of f_0 , then the homotopy h lifts to a homotopy $H : Y \times I \rightarrow E$. In particular, if $f_0 \simeq f_1 : Y \rightarrow X$ and H_0 is a lift of f_0 , we find $H_0 \simeq H_1$ where H_1 lifts f_1 .

Example 10: Example of the HLP

Trivially the identity map $\text{id}_X : X \rightarrow X$ has the HLP with respect to any space Y .

Definition 19: Covering Space

A **covering space** of X is a topological space \tilde{X} along with a continuous map $p : \tilde{X} \rightarrow X$ such that for any point $x \in X$ there exists an open neighbourhood $U \subset X$ whose preimage $p^{-1}(U) = \bigcup_{j \in J} V_j$ and the opens $V_j \subset \tilde{X}$ map homeomorphically to U under p . In other words, a covering space of X looks locally like a product of X with a discrete space.

Example 11: Example of a Covering Space

1. The projection map $p : X \times Z \rightarrow X$ is a covering map if Z is a discrete topological space. If Z is not discrete, then this is not a covering map in general.
2. The identity map $\text{id}_X : X \rightarrow X$ is always trivially a covering map.
3. While the projection of $p : X \times I \rightarrow X$ from the cylinder is not a covering map, its restriction to the boundary $\partial(X \times I) = X \times \{0, 1\} =: \tilde{X}$ gives a trivial (2-fold) covering of X .
4. Recall that the Möbius band M deformation retracts onto its core \mathbb{S}^1 . Restricting to the boundary $\partial M = \mathbb{S}^1$, one obtains a (non-trivial) covering map $\mathbb{S}^1 \rightarrow \mathbb{S}^1$. This map coincides with $z \mapsto z^2$ if we identify \mathbb{S}^1 as the unit circle in \mathbb{C} .

Theorem 3: Unique HLPs from Covering Maps

Let $p : \tilde{X} \rightarrow X$ be a covering map and Y any topological space. Then p satisfies the HLP uniquely:

$$\begin{array}{ccc} Y \times \{0\} & \xrightarrow{H_0} & \tilde{X} \\ \downarrow & \nearrow \exists! & \downarrow p \\ Y \times I & \xrightarrow{h} & X \end{array}$$

i.e. the lift H not only exists, but it is also unique.

Corollary 2:

1. Let $\gamma : I \rightarrow X$ be a path and fix a point $\tilde{x}_0 \in \tilde{X}$ such that $p(\tilde{x}_0) = \gamma(0)$. Then, there exists a unique path $\tilde{\gamma} : I \rightarrow \tilde{X}$ which starts at \tilde{x}_0 and lifts γ i.e. $p \circ \tilde{\gamma} = \gamma$.
2. Let $h : I \times I \rightarrow X$ be a (relative) homotopy of paths $h(-, 0) := \gamma_0$ and $h(-, 1) := \gamma_1$, and fix a point \tilde{x}_0 such that $p(\tilde{x}_0) = h(0, t) = \gamma_0(0) = \gamma_1(0)$. Suppose $\tilde{\gamma}_0 : I \rightarrow \tilde{X}$ is a lift of γ_0 starting at $\tilde{\gamma}_0(0) = \tilde{x}_0$. Then, there exists a unique homotopy of paths $\tilde{h} : I \times I \rightarrow \tilde{X}$ which lifts h and $\tilde{h}(-, 0) = \tilde{\gamma}_0$.

Theorem 4: Fundamental Group of the Circle

The fundamental group of the circle is $\pi_1(\mathbb{S}^1) \cong \mathbb{Z}$. It is generated by the class of

$$\alpha : I \rightarrow \mathbb{S}^1, \quad t \mapsto e^{2\pi i t}.$$

Theorem 5: Brouwer's Fixed Point Theorem

Any continuous map $f : \mathbb{D}^2 \rightarrow \mathbb{D}^2$ has a fixed point, i.e. there exists $x \in \mathbb{D}^2$ such that $f(x) = x$.

Theorem 6: Fundamental Theorem of Algebra

Every non-constant complex polynomial $p \in \mathbb{C}[z]$ has at least one root, i.e. $p(z_0) = 0$ for some z_0 .

Theorem 7: Fundamental Group of \mathbb{S}^n

The fundamental group of \mathbb{S}^n is trivial for $n \geq 2$, i.e. $\pi_1(\mathbb{S}^2) \cong 1$ for $n \geq 2$

Lemma 7: Closed Paths Homotopic to Loops

Let (X, x_0) be a topological space with an open cover $\{U_j\}_{j \in J}$ such that U_j are path-connected neighbourhoods of x_0 and $U_j \cap U_{j'}$ is path-connected for any $j, j' \in J$. Then, any closed path γ based at x_0 is homotopic to a concatenation $\gamma_1 * \gamma_2 * \dots * \gamma_n$ of loops at x_0 each of them contained in a single U_j

Corollary 3: Homeomorphisms between \mathbb{R}^2 and \mathbb{R}^n

There is no homeomorphism between \mathbb{R}^2 and \mathbb{R}^n for $n \neq 2$

Recall 4: Defining the Real Projective Space

1. The space \mathbb{RP}^2 is the quotient space:

$$\mathbb{RP}^n := (\mathbb{R}^{n+1} \setminus \{0\}) / \sim$$

where $x \sim y$ if there exists $\lambda \in \mathbb{R}$ such that $x = \lambda y$. In other words, the real projective n -space represents the lines in \mathbb{R}^{n+1} through the origin.

2. Picking representatives that lie in the unit n -sphere $\mathbb{S}^n \subset \mathbb{R}^{n+1} \setminus \{0\}$, we obtain $\mathbb{RP}^n \cong \mathbb{S}^n / \sim$ where $x \sim -x$ for all $x \in \mathbb{S}^n$, i.e. identifying antipodal points on the n -sphere.
3. Further restricting to the upper half $\mathbb{D}^n \subset \mathbb{S}^n$ we obtain:

$$\mathbb{RP}^n \cong \mathbb{D}^n / \sim$$

where $x \sim -x$ for all boundary points $x \in \partial \mathbb{D}^n \cong \mathbb{S}^{n-1}$

For example, \mathbb{RP}^0 is a one point space, $\mathbb{RP}^1 \cong \mathbb{S}^1$, while \mathbb{RP}^n are different than spheres for larger n .

Definition 20: Lift of a Path

- A lift of a path $\alpha : I \rightarrow \mathbb{RP}^n$ is a path $\tilde{\alpha} : I \rightarrow \mathbb{S}^n$ such that $p \circ \tilde{\alpha} = \alpha$
- If α is a closed path, then $[\tilde{\alpha}(0)] = [\tilde{\alpha}(1)]$ which implies

$\tilde{\alpha}(0) = \pm \tilde{\alpha}(1)$. The **sign** of α is defined by

$$\text{sgn}(\alpha) \mapsto \begin{cases} +1 & \tilde{\alpha}(0) = \tilde{\alpha}(1) \\ -1 & \tilde{\alpha}(0) = -\tilde{\alpha}(1) \end{cases}$$

Theorem 8: Group Homomorphism of the Sign

The sign induces a surjective group homomorphism

$$\text{sgn} : \pi_1(\mathbb{RP}^n) \rightarrow \mathbb{Z}_2, \quad [\alpha] \mapsto \text{sgn}(\alpha)$$

which is an isomorphism for $n \geq 2$.

Definition 21: Right Lifting Property

A map $p : X \rightarrow Y$ satisfies the **right lifting property** (RLP) with respect to a map $i : A \rightarrow B$ if any commutative square

$$\begin{array}{ccc} A & \xrightarrow{\quad} & X \\ i \downarrow & \nearrow \exists! & \downarrow p \\ B & \xrightarrow{\quad} & Y \end{array}$$

has a solution to the lifting problem making both triangles commute. Explicitly, if $f : B \rightarrow Y$ and $g : A \rightarrow X$ such that $f \circ i = p \circ g$, then there exists a map $l : B \rightarrow X$ satisfying $l \circ i = g$ and $p \circ l = f$. Dually, the map $i : A \rightarrow B$ is said to satisfy the **left lifting property** (LLP) with respect to $p : X \rightarrow Y$.

Example 13: Homotopy Lifting Property WRT Spaces

1. A map $p : X \rightarrow Y$ satisfies the homotopy lifting property with respect to a space Z iff it has the RLP with respect to the inclusion map $i : Z \times \{0\} \hookrightarrow Z \times I$, i.e. solves the following lifting problem:

$$\begin{array}{ccc} Z \times \{0\} & \xrightarrow{\quad} & X \\ \downarrow & \nearrow \exists & \downarrow p \\ Z \times I & \xrightarrow{\quad} & Y \end{array}$$

In other words, given a homotopy $h : Z \times I \rightarrow Y$ and a lift $\tilde{f} : Z \rightarrow X$ of $h(-, 0) =: f$, there is a homotopy lift $\tilde{h} : Z \times I \rightarrow X$ with $\tilde{h}(-, 0) = \tilde{f}$.

2. Dually, a map $i : A \rightarrow B$ satisfies the **homotopy extension property** (HEP) with respect to a space Z iff it has the LLP with respect to the map

$$p : Z^I \rightarrow Z, \quad \gamma \mapsto \gamma(0)$$

where $Z^I := \text{Map}(I, Z)$ is the space of paths in Z . In other words, one can solve the following lifting problem:

$$\begin{array}{ccc} A & \xrightarrow{\quad} & Z^I \\ i \downarrow & \nearrow \exists & \downarrow p \\ B & \xrightarrow{\quad} & Z \end{array}$$

Note that a map $A \rightarrow Z^I$ is the same datum as a homotopy $h : A \times I \rightarrow Z$. Given an extension $\tilde{f} : B \rightarrow Z$ of $h(-, 0)$ along i , the existence of a map $B \rightarrow Z^I$ which makes both triangles commute provides an extension of the homotopy h

to a homotopy $\tilde{h} : B \times I \rightarrow Z$ along i .

Definition 22: Fibration

1. A map $p : X \rightarrow Y$ is a **fibration** if it satisfies the HLP with respect to all spaces Z . In other words, it has the RLP with respect to the set of maps $\{i : Z \times \{0\} \hookrightarrow Z \times I\}_Z$ where Z runs over all topological spaces
2. Dually, a map $i : A \rightarrow B$ is a **cofibration** if it satisfies the HEP with respect to all spaces Z .

Theorem 9: Covering Maps are Fibrations

A covering map $p : \tilde{X} \rightarrow X$ is a fibration. Additionally, the homotopy lifts are unique:

$$\begin{array}{ccc} Z \times \{0\} & \xrightarrow{\quad} & X \\ \downarrow i & \nearrow \exists! & \downarrow p \\ Z \times I & \xrightarrow{\quad} & Y \end{array}$$

Example 14: Examples of Fibrations

1. By Theorem 9, fibrations include all covering maps
2. The projection map $p : X \times F \rightarrow X$ is always a fibration. However this map is a covering map iff F is a discrete space. Hence, this includes examples of fibrations that are not coming from covering maps.
3. An important example of a cofibration is the inclusion $i : X \rightarrow M_f$ where M_f is the mapping cylinder of $f : X \rightarrow Y$. We have seen that any continuous map $f : X \times Y$ factors through the mapping cylinder:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow i & \nearrow r \\ & M_f & \end{array}$$

In particular, every map factors through a cofibration and a homotopy equivalence.

Example 15: Covering Spaces

1. The projection map $p : X \times D \rightarrow X$ where D is a discrete space. Note that $X \times D$ cannot be path-connected unless D is a one-point set.
2. The covering map $\mathbb{R} \rightarrow \mathbb{S}^1$, $t \mapsto e^{2\pi i t}$ which we can use to compute the fundamental group of \mathbb{S}^1 .
3. The degree- n map $F_n : \mathbb{S}^1 \rightarrow \mathbb{S}^1$, $z \mapsto z^n$ provides an n -fold covering of \mathbb{S}^1 by itself.
4. The product of two covering maps $p_i : \tilde{X}_i \rightarrow X_i$ is also a covering map $p_1 \times p_2 : \tilde{X}_1 \times \tilde{X}_2 \rightarrow X_1 \times X_2$
5. The product of F_n and F_m in the third example also provides a self covering of the torus:
$$T^2 = \mathbb{S}^1 \times \mathbb{S}^1 \rightarrow T^2, \quad (z, w) \mapsto (z^n, w^m)$$
6. Similarly, there is a covering $\mathbb{R}^2 \rightarrow T^2$.
7. The 2-fold covering $\mathbb{S}^n \rightarrow \mathbb{RP}^n$ which was used to compute the fundamental group of \mathbb{RP}^n

Theorem 10: Homomorphism of Covering Maps

Let $p : \tilde{X} \rightarrow X$ be a covering map. The induced group homomorphism

$$p_* : \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$$

is injective for any $\tilde{x}_0 \in p^{-1}(x_0)$. Its image consists of (classes of) loops in X based at x_0 that lift to loops in \tilde{X} based at \tilde{x}_0

Remark 5: Notation for Covers

Fix a covering map $p : \tilde{X} \rightarrow X$ and $x_0 \in X$ a fixed point. We will write

$$G := \pi_1(X, x_0)$$

for the fundamental group of X at x_0 and

$$H := p_* \pi_1(\tilde{X}, \tilde{x}_0) \subset G$$

for the subgroup determined by the covering map. The subgroup H depends on the choice of fiber point $\tilde{x}_0 \in p^{-1}(x_0)$ and we shall see that its subgroups for different fiber points are conjugate to each other. Finally, the fiber over x_0 will be denoted by

$$F_{x_0} := p^{-1}(x_0)$$

Lemma 8: Transitive Actions

If \tilde{X} is path-connected, then the G -action on F_{x_0} is transitive, i.e. for any $\tilde{x}, \tilde{x}' \in F_{x_0}$, there exists a $\alpha \in G$ such that $\tilde{x} \cdot \alpha = \tilde{x}'$.

Theorem 11: Path-Connected Correspondence

If \tilde{X} is path-connected, then there is a one-to-one correspondence between right cosets and fiber points, i.e. a bijection

$$G\tilde{x} \backslash G \rightarrow F_{x_0}, \quad G\tilde{x} \cdot g \mapsto \tilde{x} \cdot g$$

Thus the index of $G_{\tilde{x}}$ in G coincides with the cardinality of the fiber F_{x_0} :

$$[G : G_{\tilde{x}}] = |F_{x_0}| \quad (20)$$

Corollary 4: If \tilde{X} is simply-connected, then there is a bijection

$$G \rightarrow F_{x_0}$$

Equation (20) becomes

$$|G| = |F_{x_0}| \quad (21)$$

Definition 23: Deck Transformation

A **deck transformation** of a covering map $p : \tilde{X} \rightarrow X$ is a self-homeomorphism $D : \tilde{X} \xrightarrow{\cong} \tilde{X}$ such that $p \circ D = p$.

Deck transformations form a group $\text{Deck}(p)$. Indeed, for any two deck transformations D, D' their composite is also a deck transformation since $p \circ D \circ D' = p \circ D' = p$. Moreover, if D is a deck transformation then so is its inverse D^{-1} as $p \circ D^{-1} = p \circ D \circ D' = p$.

For example, deck transformations of the covering map $\mathbb{R} \rightarrow \mathbb{S}^1$ are precisely translations by integers:

$$D_n : \mathbb{R} \rightarrow \mathbb{R}, \quad t \mapsto t + n$$

In particular, the group of deck transformations is \mathbb{Z} . More generally, the group of deck transformations $\text{Deck}(p)$ of a universal covering p is isomorphic to the fundamental group G . From now on $p : \tilde{X} \rightarrow X$ will be a covering with \tilde{X} path-connected and X

path-connected and locally path-connected.

Example 16: Topologist's Sine Curve

Recall that the topologist's sine curve

$$X = \{0\} \times [-1, 1] \cup \left\{ \left(x, \sin\left(\frac{1}{x}\right) \right) \mid 0 < x \leq \frac{1}{2\pi} \right\} \subset \mathbb{R}^2$$

is an example of a connected, but not path-connected space. Let Z be the quotient of X by identifying the points $(0, 0) \sim (\frac{1}{2\pi}, 0)$. Z is a path-connected space but not locally path-connected.

Theorem 12: Solutions to the Lifting Problem

Let $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ be a covering with \tilde{X} path-connected and locally path-connected, and let $g : (Z, z_0) \rightarrow (X, x_0)$ be a pointed map. Then, there exists a solution to the lifting problem

$$\begin{array}{ccc} & (\tilde{X}, \tilde{x}_0) & \\ \exists \tilde{g} \nearrow & \downarrow p & \searrow \\ (Z, z_0) & \xrightarrow{g} & (X, x_0) \end{array} \quad \text{if and only if } g_* \pi_1(Z, z_0) \subset p_* \pi_1(\tilde{X}, \tilde{x}_0)$$

Moreover, if a solution to the lifting problem exists, then it is also unique.

Remark 10: If in Theorem 12, g is a covering map, then so is its lift \tilde{g} . In particular, homomorphisms of covering maps are also covering maps.

Corollary 5: Commuting Covering Maps

Let $p : \tilde{X} \rightarrow X$ be a covering with \tilde{X} simply connected. Then, for any covering $p' : \tilde{X}' \rightarrow X$ there exists a covering $\tilde{p} : \tilde{X} \rightarrow \tilde{X}'$ such that

$$\begin{array}{ccc} & \tilde{X}' & \\ \tilde{p} \nearrow & \downarrow p' & \\ \tilde{X} & \xrightarrow{p} & X \end{array}$$

commutes. In other words, if a simply connected covering \tilde{X} of X exists, then it covers all other possible coverings. This is why such a covering is called the **universal covering** of X . For example, we have seen \mathbb{R} as the universal covering of \mathbb{S}^1 and \mathbb{RP}^n as the universal covering of \mathbb{RP}^n .

Theorem 24: Covering Isomorphism

Let $p : \tilde{X} \rightarrow X$ be a covering with X path-connected and locally path-connected. Let $H \subset \pi_1(X, x_0)$ denote the subgroup determined by the covering map. Then, there exists a group isomorphism:

$$\text{Deck}(p) \cong N(H)/H$$

where $N(H)$ denotes the normalizer.

Definition 24: Normal Coverings

A covering $p : \tilde{X} \rightarrow X$ is called **normal** if the subgroup H is normal

Trivially, universal coverings are always normal. All the examples so far were normal since all the fundamental groups we have seen so far were abelian.

Corollary 6: Let \tilde{X} be simply-connected. Then

$$\text{Deck}(p) \cong \pi_1(X, x_0).$$

Recall 6: The Figure Eight Space

Let X be the figure eight space

$$X = \mathbb{S}^1 \vee \mathbb{S}^1.$$

Consider an oriented bicolored graph \tilde{X} whose vertices are all 4-valent with one incoming edge of each color and one outgoing edge of each color. Bicolored means that each edge is labelled by a or b . Such a graph determines a covering map

$$p : \tilde{X} \rightarrow X$$

by sending all vertices to the unique vertex of the figure-eight graph and the edges are sent to one of the loops. A universal covering is obtained by the following graph: where vertical edges are oriented upwards and labelled by b and horizontal edges are oriented to the right and labelled by a . Deck transformations are freely generated by either D_a or D_b , where D_a (resp. D_b) acts on the graph by shifting all edges once to the right, rescaling them appropriately. In other words..

Theorem 14: Fundamental Group of $\mathbb{S}^1 \vee \mathbb{S}^1$

The fundamental group of $\mathbb{S}^1 \vee \mathbb{S}^1$ is the free group generated by two elements, i.e.

$$\pi_1(\mathbb{S}^1 \vee \mathbb{S}^1) \cong \langle a, b \rangle$$

Example 20.1: Covering The Möbius Band

Recall the Möbius band $M : \mathbb{R} \times I / \sim$ is defined by the quotient $(x, y) \sim (x + 1, 1 - y)$. We obtain the homotopy equivalence using covering theory. The quotient map

$$q : \mathbb{R} \times I \rightarrow M$$

is the universal covering, since $\mathbb{R} \times I$ is simply-connected. For some $n \in \mathbb{Z}$, let D_n be the deck transformation:

$$D_n : \mathbb{R} \times I \rightarrow \mathbb{R} \times I, \quad (x, y) \mapsto (x + n, y_n)$$

where $y_n = y$ if n is even and $y_n = 1 - y$ for n odd. One can easily check that these are all deck transformations and $\text{Deck}(p)$ is generated by D_1 since

$$D_n = (D_1)^n$$

For odd n , there are n -fold self-coverings $M \rightarrow M$. For even n , there are n -fold coverings by the torus $T^2 \rightarrow M$.

Example 20.2: Covering the Klein Bottle

Consider the Klein bottle

$$K = \mathbb{R}^2 / \sim$$

where $(x, y) \sim (x + 1, 1 - y) \sim (x, y + 1)$ for all $(x, y) \in \mathbb{R}^2$. The quotient map

$$q : \mathbb{R}^2 \rightarrow K$$

is the universal covering map. Consider the deck transformation

$$D_a : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad (x, y) \mapsto (x, y + 1)$$

and

$$D_b : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad (x, y) \mapsto (x + 1, 1 - y)$$

These two deck transformations generate the deck transformation group $\text{Deck}(q)$ and satisfy the relation:

$$D_b \circ D_a \circ D_b^{-1} \circ D_a = \text{id}.$$

Proposition 10: Fundamental Group of the Klein Bottle

The fundamental group of the Klein Bottle is:

$$\pi_1(K) = \langle a, b \rangle / \langle aba^{-1}b \rangle$$

Theorem 15: Seifert-Vam Kampen Theorem

Let X be a topological space with a fixed point x_0 . Let $\{U_\alpha\}_\alpha$ be an open cover of X consisting of path-connected open sets U_α containing the fixed point x_0 . The inclusions $U_\alpha \subset X$ induce a group homomorphism:

$$\Phi : *_\alpha \pi_1(U_\alpha) \rightarrow \pi_1(X).$$

1. If $U_\alpha \cap U_\beta$ is path-connected for any α, β , then Φ is surjective.
2. If $U_\alpha \cap U_\beta \cap U_\gamma$ is path-connected for any α, β, γ , then the kernel of Φ is generated by elements $i_{\alpha\beta}(w)i_{\beta\alpha}(w)^{-1}$ where $w \in \pi_1(U_\alpha \cap U_\beta)$ and $i_{\alpha\beta} : \pi_1(U_\alpha \cap U_\beta) \rightarrow \pi_1(U_\alpha)$ is the induced homomorphism from the inclusion $U_\alpha \cap U_\beta \subset U_\alpha$.

The assumption $U_\alpha \cap U_\beta$ are path-connected ensures that words $\pi_1(U_\alpha)$ generate $\pi_1(X)$. The assumption $U_\alpha \cap U_\beta \cap U_\gamma$ is path-connected gives a presentation for the group $\pi_1(X)$.

Example 17: Sifert-Vam Kampen on $\mathbb{S}^1 \vee \mathbb{S}^1$

Consider the figure eight $\mathbb{S}^1 \vee \mathbb{S}^1 = \mathbb{S}^1 \times \{1\} \cup \{1\} \times \mathbb{S}^1$ and let

$$U_i := \mathbb{S}^1 \vee \mathbb{S}^1 \setminus \{x_i\}$$

be the complements of the points $x_1 = (-1, 1)$ and $x_2 = (1, -1)$. The sets U_1 and U_2 are open path-connected and cover $\mathbb{S}^1 \vee \mathbb{S}^1$. In fact, they are both homotopy equivalent to the circle $U_i \simeq \mathbb{S}^1$. Their intersection $U_1 \cap U_2$ is contractible, and applying SVK we find,

$$\pi_1(\mathbb{S}^1 \vee \mathbb{S}^1) \cong \mathbb{Z} * \mathbb{Z}.$$

Example 18: Fundamental Group of Wedged Circles

Let (X_α, x_α) be a fIYL of path-connected pointed spaces and consider their wedge sum

$$X := \bigvee_{\alpha} X_{\alpha}.$$

suppose that each $x_\alpha := X_\alpha \cap \bigvee_{\beta \neq \alpha} U_\beta \subset X$. By contractibility of the U_α 's we have homotopy equivalences $A_\alpha \simeq X_\alpha$. Moreover, the intersection $A_\alpha \cap A_\beta$ is contractible for any $\alpha \neq \beta$. Applying SVK we obtain

$$\pi_1(X) \cong *_\alpha \pi_1(X_\alpha).$$

In particular, the fundamental group of the n -th wedge sum of circles is the free group on n -generators:

$$\pi_1\left(\bigwedge^n \mathbb{S}^1\right) \cong \mathbb{Z} * \dots * \mathbb{Z} \cong \langle \alpha_1, \dots, \alpha_n \rangle. \quad (23)$$

Definition 25: CW Complexes

A special class of topological spaces which are constructed inductively attaching n -dimensional disks or n -cells are called **CW complexes**. They are described as follows:

1. A set X^0 of **vertices** or 0-cells
2. Inductively construct the n -skeleton X^n from X^{n-1} by attaching n -dimensional disks \mathbb{D}_α^n by attaching maps $\phi_\alpha : \partial \mathbb{D}_\alpha^n = \mathbb{S}_\alpha^{n-1} \rightarrow X^{n-1}$. In other words,

$$X^n = X^{n-1} \amalg_{\phi_\alpha} \coprod_{\alpha} \mathbb{D}_\alpha^n.$$

Equivalently, a **CW Complex** is a space X along with a filtration of subspaces

$$X^0 \subset \dots \subset X^n \subset X^{n+1} \subset \dots \subset X$$

such that $X^n \setminus X^{n-1}$ is homeomorphic to a disjoint union of n -dimensional open disks, and X^0 is discrete.

Example 19: Examples of CW Complexes

1. The Torus $T^2 = I^2 / \sim$ can be made into a CW complex with: $X^0 = \{(0, 0)\}$, $X^1 = \{(a, 0) \mid a \in I\} \cup \{(0, b) \mid b \in I\}$ and $X^2 = T^2$. In particular, it has one 0-cell, two 1-cells, and one

- 2-cell.
2. The real projective plane \mathbb{RP}^2 can be made into a CW complex with $X^0 = *$, $X^1 = \mathbb{RP}^1 = \mathbb{S}^1$ and X^2 obtained by attaching a 2-disk to \mathbb{S}^1 along the quotient map $\mathbb{S}^1 \rightarrow \mathbb{RP}^1$

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum. Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris. Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula. Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur. Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio. Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent feugiat sapien aliquet odio. Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at, consectetur sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget

lectus. Proin eu metus. Sed porttitor. In hac habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui. Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis, ultrices a, dui. Morbi luctus, wisi viverra faucibus pretium, nibh est placerat odio, nec commodo wisi enim eget quam. Quisque libero justo, consectetur a, feugiat vitae, porttitor eu, libero. Suspendisse sed mauris vitae elit sollicitudin malesuada. Maecenas ultricies eros sit amet ante. Ut venenatis velit. Maecenas sed mi eget dui varius euismod. Phasellus aliquet volutpat odio. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Pellentesque sit amet pede ac sem eleifend consectetur. Nullam elementum, urna vel imperdiet sodales, elit ipsum pharetra ligula, ac pretium ante justo a nulla. Curabitur tristique arcu eu metus. Vestibulum lectus. Proin mauris. Proin eu nunc eu urna hendrerit faucibus. Aliquam auctor, pede consequat laoreet varius, eros tellus scelerisque quam, pellentesque hendrerit ipsum dolor sed augue. Nulla nec lacus. Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui. Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem dictum tortor, vel consectetur odio sem sed wisi. Sed feugiat. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Ut pellentesque augue sed urna. Vestibulum diam eros, fringilla et, consectetur eu, nonummy id, sapien. Nullam at lectus. In sagittis ultrices mauris. Curabitur malesuada erat sit amet massa. Fusce blandit. Aliquam erat volutpat. Aliquam euismod. Aenean vel lectus. Nunc imperdiet justo nec dolor. Etiam euismod. Fusce facilisis lacinia dui. Suspendisse potenti. In mi erat, cursus id, nonummy sed, ullamcorper eget, sapien. Praesent pretium, magna in eleifend egestas, pede pede pretium lorem, quis consectetur tortor sapien facilisis magna. Mauris quis magna varius nulla scelerisque imperdiet. Aliquam non quam. Aliquam porttitor quam a lacus. Praesent vel arcu ut tortor cursus volutpat. In vitae pede quis diam bibendum placerat. Fusce elementum convallis neque. Sed dolor orci, scelerisque ac, dapibus nec, ultricies ut, mi. Duis nec dui quis leo sagittis commodo.