

# Differential Cohomology and Virasoro Central Extensions

Yu Leon Liu

Harvard University

April 3rd 2022

Based on [[arXiv:2112.10837](https://arxiv.org/abs/2112.10837)]

Joint work with Arun Debray and Christopher Weis

# Table of Contents

- 1 Motivation
- 2 Virasoro groups and central extensions
- 3 Differential cohomology
- 4 Main theorem

# Table of Contents

- 1 Motivation
- 2 Virasoro groups and central extensions
- 3 Differential cohomology
- 4 Main theorem

**Virasoro groups** is a  $\mathbf{R}$  family of central extension of  $\text{Diff}^+(S^1)$ , the group of orientation preserving smooth automorphism of  $S^1$ . The central extension is describe by the Bott-Thurston cocycle. The goal of this talk is to give a novel geometric description these central extensions, using differential cohomology. thus affirmatively answering a conjecture of Freed-Hopkins.

# Table of Contents

- 1 Motivation
- 2 Virasoro groups and central extensions
- 3 Differential cohomology
- 4 Main theorem

# Bott-Thurston cocycles

Recall that  $\text{Diff}^+(S^1)$  is the group of orientation preserving smooth automorphism of  $S^1$ . It is an infinite-dimensional Frechet Lie group.

# Bott-Thurston cocycles

Recall that  $\text{Diff}^+(S^1)$  is the group of orientation preserving smooth automorphism of  $S^1$ . It is an infinite-dimensional Frechet Lie group.

## Definition

The Virasoro group  $\tilde{\Gamma}_\lambda$ , for  $\lambda \in \mathbf{R}$ , is a  $U(1)$  central extension of  $\text{Diff}^+(S^1)$ , described by the **Bott-Thurston cocycle**

$B_\lambda : \text{Diff}^+(S^1) \times \text{Diff}^+(S^1) \rightarrow U(1)$ :

$$B_\lambda(\gamma_1, \gamma_2) = \exp \left( -\frac{i\lambda}{48\pi} \int_{S^1} \log(\gamma'_1 \circ \gamma_2) d(\log(\gamma_2))' \right) \quad (1)$$

for  $\gamma_1, \gamma_2 \in \text{Diff}^+(S^1)$ , viewed as morphisms  $S^1 \rightarrow S^1$ .

# Central Extensions

Let's briefly review what is a central extension:



# Central Extensions

Let's briefly review what is a central extension:

## Definition

Let  $G$  be a group and  $A$  be an abelian group, a central extension of  $G$  by  $A$  is a group  $\tilde{G}$  with short exact sequence:

$$0 \rightarrow A \rightarrow \tilde{G} \rightarrow G \rightarrow 1 \quad (2)$$

such that subgroup  $A \subset \tilde{G}$  is in the center, that is, it commutes with every element of  $\tilde{G}$ .

# Central extension as group cohomology: I

As many other things, central extensions can be classified by cohomology groups:

# Central extension as group cohomology: I

As many other things, central extensions can be classified by cohomology groups:

## Proposition

*Let  $G$  be a discrete group, then the isomorphism class of central extensions of  $G$  by  $A$  is classified by group cohomology class  $H^2(G; A) \simeq H^2(BG; A)$ , where  $BG$  is the classifying space of  $G$ .*

Given a cocycle class  $b \in C^2(G; A)$ , viewed as a map  $b : G \times G \rightarrow A$  satisfying some cocycle conditions. Then  $\tilde{G} = G \times A$  as a set, with multiplication  $(g, a) \cdot (g', a') := (g \cdot g', a + a' + b(g, g'))$ .

# Central extension as group cohomology: II

In our case,  $\text{Diff}^+(S^1)$  is an infinite dimension Frechet Lie group, and we care about its topology. Smooth  $U(1)$  central extensions of Lie groups are **NOT** classified by ordinary cohomology classes.

# Central extension as group cohomology: II

In our case,  $\text{Diff}^+(S^1)$  is an infinite dimension Frechet Lie group, and we care about its topology. Smooth  $U(1)$  central extensions of Lie groups are **NOT** classified by ordinary cohomology classes.

We need a cohomology theory that remembers the smooth structure.

# Central extension as group cohomology: II

In our case,  $\text{Diff}^+(S^1)$  is an infinite dimension Frechet Lie group, and we care about its topology. Smooth  $U(1)$  central extensions of Lie groups are **NOT** classified by ordinary cohomology classes.

We need a cohomology theory that remembers the smooth structure.

The answer is **differential cohomology**.

# Table of Contents

- 1 Motivation
- 2 Virasoro groups and central extensions
- 3 Differential cohomology
- 4 Main theorem

# Sheaves on smooth manifolds

Let  $M$  be a manifold, then the ordinary cohomology groups  $H^*(M; A)$  depends only on the homotopy classes of  $M$ . It is the cohomology of the constant sheave  $\underline{A}$  on  $M$ . On the other hand, the  $i$ -th cohomology form on  $M$ ,  $\Omega^i(M)$  is sensitive to the smooth structure of  $M$ .



# Sheaves on smooth manifolds

Let  $M$  be a manifold, then the ordinary cohomology groups  $H^*(M; A)$  depends only on the homotopy classes of  $M$ . It is the cohomology of the constant sheave  $\underline{A}$  on  $M$ . On the other hand, the  $i$ -th cohomology form on  $M$ ,  $\Omega^i(M)$  is sensitive to the smooth structure of  $M$ .

We can view both constant sheaves and differential forms as sheaves on  $Mfld$  the site of smooth manifolds.

# Sheaves on smooth manifolds

Let  $M$  be a manifold, then the ordinary cohomology groups  $H^*(M; A)$  depends only on the homotopy classes of  $M$ . It is the cohomology of the constant sheave  $\underline{A}$  on  $M$ . On the other hand, the  $i$ -th cohomology form on  $M$ ,  $\Omega^i(M)$  is sensitive to the smooth structure of  $M$ .

We can view both constant sheaves and differential forms as sheaves on  $Mfld$  the site of smooth manifolds.

Even though  $\Omega^i$  are not homotopy invariant, the chain complex of sheaves  $\Omega^* = 0 \rightarrow \Omega^0 \xrightarrow{d} \Omega^1 \xrightarrow{d} \dots$  is a homotopy invariant, in fact,

## Theorem (de Rham)

*The chain complex  $\Omega^*$  is the constant sheave  $\mathbb{R}$ , as a chain complex concentrated in degree 0.*

With this in mind, we define the (chain complex of) sheave  $\mathbb{Z}(n)$  as

$$\mathbb{Z}(n) = \mathbb{Z} \rightarrow \Omega^0 \rightarrow \Omega^1 \rightarrow \cdots \rightarrow \Omega^{n-1} \rightarrow 0. \quad (4)$$

With this in mind, we define the (chain complex of) sheave  $\mathbb{Z}(n)$  as

$$\mathbb{Z}(n) = \mathbb{Z} \rightarrow \Omega^0 \rightarrow \Omega^1 \rightarrow \cdots \rightarrow \Omega^{n-1} \rightarrow 0. \quad (4)$$

These sheaves  $\mathbb{Z}(n)$  are both sensitive to topology (from  $\mathbb{Z}$ ) and the smooth structure (from  $\Omega^i$ ).

With this in mind, we define the (chain complex of) sheave  $\mathbb{Z}(n)$  as

$$\mathbb{Z}(n) = \mathbb{Z} \rightarrow \Omega^0 \rightarrow \Omega^1 \rightarrow \cdots \rightarrow \Omega^{n-1} \rightarrow 0. \quad (4)$$

These sheaves  $\mathbb{Z}(n)$  are both sensitive to topology (from  $\mathbb{Z}$ ) and the smooth structure (from  $\Omega^i$ ).

There is also a form of integration: let  $M$  be a closed oriented  $d$ -dimensional manifold, then there is an integration map:

$$\int_M : H^*(M; \mathbb{Z}(n)) \rightarrow H^{*-d}(M; \mathbb{Z}(n-d)). \quad (5)$$

There is also a relative version of this.

# $\mathbb{Z}(1)$ and line bundles

For example, a degree cocycle in  $C^2(M; \mathbb{Z}(1))$  can be describe as follows: fix an open covering  $\{U_i\}$  of  $M$ , we have 1-form  $\alpha_i$  on the open subsets  $U_i$ , and 0-form  $f_{ij}$  on intersections  $U_i \cap U_j$ , such that  $\alpha_i - \alpha_j = f_{ij}$

# $\mathbb{Z}(1)$ and line bundles

For example, a degree cocycle in  $C^2(M; \mathbb{Z}(1))$  can be describe as follows: fix an open covering  $\{U_i\}$  of  $M$ , we have 1-form  $\alpha_i$  on the open subsets  $U_i$ , and 0-form  $f_{ij}$  on intersections  $U_i \cap U_j$ , such that  $\alpha_i - \alpha_j = f_{ij}$

This precisely describe the data of a  $U(1)$  principal bundle on  $M$ !

## Proposition

*$H^2(M; \mathbb{Z}(1))$  is the group of isomorphism classes of  $U(1)$  principal bundles on  $M$ .*

Furthermore,  $H^2(M; \mathbb{Z}(2))$  is the group of isomorphism classes of  $U(1)$  principal bundles with connections on  $M$ .

# $\mathbb{Z}(1)$ and central extensions

While  $H^2(-; \mathbb{Z}(1))$  classifies  $U(1)$  principal bundle,  $H^3(-; \mathbb{Z}(1))$  classifies  $U(1)$  central extensions:

## Theorem

*Let  $G$  be a smooth (possibly infinite dimensional) Lie group,  $BG$  its classifying space. Then  $H^3(BG; \mathbb{Z}(1))$  classifies smooth central extensions of  $G$  by  $U(1)$ .*



# Differential characteristic classes

Let  $G$  be a Lie group, and  $B_\bullet G$  the classifying space. Then  $H^*(B_\bullet G; \mathbb{Z})$  are the characteristic classes of  $G$ . Similarly,  $H^*(B_\bullet G; \mathbb{Z}(n))$  are differential characteristic classes of  $G$ .

# Differential characteristic classes

Let  $G$  be a Lie group, and  $B_\bullet G$  the classifying space. Then  $H^*(B_\bullet G; \mathbb{Z})$  are the characteristic classes of  $G$ . Similarly,  $H^*(B_\bullet G; \mathbb{Z}(n))$  are differential characteristic classes of  $G$ .

We will need the following key fact:

## Theorem (Bott)

$$H^4(B_\bullet \mathrm{GL}_1^+(\mathbb{R}); \mathbb{Z}(2)) \simeq \mathbb{R}.$$

# Table of Contents

- 1 Motivation
- 2 Virasoro groups and central extensions
- 3 Differential cohomology
- 4 Main theorem

# Key Idea I

We want to get a  $\mathbb{R}$  family of central extension of  $\text{Diff}^+(S^1)$  by  $U(1)$ , therefore we want a  $\mathbb{R}$  family in  $H^3(B_\bullet \text{Diff}^+(S^1), \mathbb{Z}(1))$ . We get this by pullback and integration:

# Key Idea I

We want to get a  $\mathbb{R}$  family of central extension of  $\text{Diff}^+(S^1)$  by  $U(1)$ , therefore we want a  $\mathbb{R}$  family in  $H^3(B_\bullet \text{Diff}^+(S^1), \mathbb{Z}(1))$ . We get this by pullback and integration:

Consider the canonical  $\text{Diff}^+(S^1)$  action on  $S^1$ , note that

- The quotient  $S^1/\text{Diff}^+(S^1)$  has a map to  $B_\bullet \text{Diff}^+(S^1) = */\text{Diff}^+(S^1)$ . Since the action of  $\text{Diff}^+(S^1)$  on  $S^1$  is orientation preserving, this is a oriented  $S^1$  fiber bundle.
- The tangent bundle of  $S^1$  gives a map  $TS^1 : S^1 \rightarrow B_\bullet \text{GL}_1^+(\mathbb{R})$ . Since the action of  $\text{Diff}^+(S^1)$  on  $S^1$  is smooth, the tangent bundle is  $\text{Diff}^+(S^1)$ -equivariant. Equivalently, the tangent bundle factors through the quotient as a map  $TS^1 : S^1/\text{Diff}^+(S^1) \rightarrow B\text{GL}_1^+(\mathbb{R})$ .

# Key Idea II

To summarize, we have a span of maps:

$$\begin{array}{ccc} S^1/\mathrm{Diff}^+(S^1) & \xrightarrow{TS^1} & B_\bullet \mathrm{GL}_1^+(\mathbb{R}) \\ \downarrow & & \\ B_\bullet \Gamma. & & \end{array} \quad (7)$$

Note the vertical map is a  $S^1$  fibration, something we can integrate against. Therefore we get a map:

$$\begin{array}{ccc} H^4(S^1/\mathrm{Diff}^+(S^1); \mathbb{Z}(2)) & \longleftarrow & H^4(B_\bullet \mathrm{GL}_1^+(\mathbb{R}); \mathbb{Z}(2)) \\ \downarrow \int_{S^1} & & \\ H^3(B_\bullet \Gamma; \mathbb{Z}(1)). & & \end{array} \quad (8)$$

# Main theorem

Finally, we can state the conjecture of Freed and Hopkins that we proved:

**Theorem (Y.L., Arun Debray, Christoph Weis)**

*The image of map  $\mathbb{R} \simeq H^4(B_\bullet \mathrm{GL}_1^+(\mathbb{R}); \mathbb{Z}(2)) \rightarrow H^3(B_\bullet \mathrm{Diff}^+(S^1); \mathbb{Z}(1))$  are the Virasoro central extensions  $\tilde{\Gamma}_\lambda$ .*

Furthermore, we explicitly recovers the Bott-Thurston cocycles when considering the map on cocycles.

# Proof sketch I

We construct explicit cocycles and compute the map on the level of cocycles.

- We find 1-form cocycles for  $H^4(B_\bullet \mathrm{GL}_1^+(\mathbb{R}); \mathbb{Z}(2))$ , using the canonical simplicial resolution of  $B_\bullet \mathrm{GL}_1^+(\mathbb{R})$ .



# Proof sketch I

We construct explicit cocycles and compute the map on the level of cocycles.

- We find 1-form cocycles for  $H^4(B_\bullet \mathrm{GL}_1^+(\mathbb{R}); \mathbb{Z}(2))$ , using the canonical simplicial resolution of  $B_\bullet \mathrm{GL}_1^+(\mathbb{R})$ .
- We pullback to 1-form cocycles on  $S^1/\mathrm{Diff}^+(S^1)$ , using the simplicial realization of  $S^1 = Fr_+(S^1)/\mathrm{GL}_1^+(\mathbb{R})$ , where  $Fr_+(S^1)$  is the oriented frame bundle.

# Proof sketch II

Here's the key point of the proof:

- Now we move the cocycles across the double complex associated to the bisimplicial object  $S^1/\text{Diff}^+(S^1) = \text{GL}_1^+(\mathbb{R}) \backslash Fr_+(S^1) / \text{GL}_1^+(\mathbb{R})$ , to get cocycles on the simplicial resolution for  $S^1/\text{Diff}^+(S^1)$ .

$$\begin{array}{ccccc}
 & & \Omega^1(F \times \mathbb{R}^{\times 2}) & & \\
 & & \uparrow -\log(\gamma'_1 \circ \gamma_2) d\log(\gamma_2) & & \\
 & \Omega^1(\Gamma \times F \times \mathbb{R}) & \longleftarrow & \Omega^1(F \times \mathbb{R}) & \\
 & \uparrow \begin{array}{l} \text{blue } x \log \gamma' - \\ \text{red } x \log \gamma' = 0 \end{array} & & \uparrow \log(v) \quad d\log(\gamma') & \\
 \Omega^1(\Gamma^{\times 2} \times F) & \longleftarrow & \Omega^1(\Gamma \times F) & & \\
 & \xleftarrow{x_2 dx_1} & \xleftarrow{x \quad d\log(v)} & & 
 \end{array}$$

# Proof sketch III

- Lastly, we integrate over  $S^1$  and immediately see that we recover the Bott-Thurston cocycles! Q.E.D.

Thank you for listening!