Low Energy High Precision Test

Aryan K. Somkuwar, Writipriya Paul
Indian Institute of Science, Bangalore

Abstract

The Standard Model of particle physics is remarkably successful, but it does not have a proper theory for the matter-antimatter asymmetry of the universe and the unnatural absence of Charge-Parity (CP) violation in the strong interaction. The neutcover , its connection to Parity (P) and Time reversal violation. Electric Dipole Moment (nEDM) can help us to link these phenomena.in this term paper we try to cover , its connection to Parity (P) and Time Reversal (T) violation and Strong CP problem.And of course current experimental detection methodologies for both the nEDM and the axion. In addition, we consider low-energy precision tests of the Standard Model through pion decay processes. Pion beta decay provides a theoretically clean determination of the CKM matrix element V_{ud}, while purely leptonic pion decays offer a sensitive probe of lepton universality through the ratio R_π. These observables, together with precise measurements from experiments such as PIBETA, PEN, and PIENU, serve as complementary probes of possible physics beyond the Standard Model.

Contents

1 Introduction **3**

2 Maths Pre-req **3**
2.1 The Wigner-Eckart Theorem . . . 3
2.2 P and T Violation . . . 3

3 The Strong CP Problem and Naturalness **3**
3.1 The θ-Vacuum in QCD . . . 3
3.2 The Conflict and Naturalness . . . 4

4 Calculating relation between d_n and θ **4**

5 The Peccei-Quinn Mechanism and the Axion **6**

6 Baryon Asymmetry of the Universe (BAU) **6**

7 Experimental Detection Methodologies **7**
7.1 intro . . . 7
7.2 Measuring the nEDM and nMDM (Ramsey Resonance) . . . 7
7.3 Ramsey's method . . . 7
7.4 summary of experiment . . . 8
7.5 Set Up . . . 9
7.6 Cooling process . . . 11
7.7 Check check . . . 12

8 Axion Detection **12**

9 Question asked during class: **13**

10 Conclusion **13**

11 Pion Beta Decay and Testing CKM Unitarity **15**
11.1 $\pi\beta$ Decay Rate . . . 15
11.2 Radiative Corrections in Pion Beta Decay . . . 16
11.2.1 Short-distance electroweak correction . . . 17
11.2.2 Long-distance electromagnetic correction: general structure . . . 17
11.2.3 Virtual-photon loop corrections . . . 18
11.2.4 Real-photon emission and bremsstrahlung . . . 18
11.2.5 Need for chiral perturbation theory . . . 19
11.2.6 Differential rate with radiative corrections . . . 20
11.2.7 Infrared cancellation in more detail . . . 20
11.2.8 Interpretation for the extraction of V_{ud} . . . 20
11.2.9 Corrected Decay Rate . . . 21
11.3 Experimental Realization . . . 21
11.4 The PIBETA Experiment . . . 22
11.4.1 Data Acquisition and Event Selection . . . 22
11.4.2 Event Reconstruction . . . 22
11.4.3 Background Suppression . . . 22
11.4.4 Experimental Results . . . 23
11.5 Next-Generation Experiments: PIONEER . . . 24

12 Lepton Universality and Precision Tests in Pion Decay **24**
12.1 Effective weak interaction and the pion decay constant . . . 25
12.2 Why helicity suppression occurs . . . 26
12.3 Decay rate and the ratio R_π . . . 26
12.4 Lepton universality and what R_π tests . . . 27
12.5 Chiral perturbation theory and the low-energy description . . . 27
12.5.1 Structure of radiative corrections . . . 28
12.5.2 Explicit one-loop analysis in χPT . . . 28
12.5.3 Real-photon emission and infrared cancellation . . . 30
12.5.4 Sirlin-type representation of $\delta_{\rm rad}$ . . . 30
12.6 Prediction for R_π . . . 31
12.7 The PEN Experiment . . . 31
12.7.1 Overview and Physics Goal . . . 31
12.7.2 Beam and Target System . . . 31
12.7.3 Detector Design . . . 31
12.7.4 Waveform Digitization and Signal Formation . . . 32
12.7.5 Data Acquisition and Triggering . . . 32
12.7.6 Event Reconstruction and Selection . . . 32
12.7.7 Systematic Uncertainties . . . 33
12.8 The PIENU Experiment . . . 33
12.9 Results and Implications . . . 33

13 Conclusion **34**

14 Question from the Presentation **34**

15 References **35**

1 Introduction

Despite being electrically neutral, its quark substructure gives it a magnetic dipole moment (nMDM). The core question of the paper is whether it also possesses a permanent electric dipole moment (nEDM),if yes then how??. While its magnetic dipole moment (nMDM) is well-established, the existence of a permanent electric dipole moment (nEDM), denoted as d_n, remains one of the Questionable qantity in modern physics. The current experimental upper bound is $|d_n| < 1.8 \times 10^{-26}\, e \cdot \text{cm}$ at a 90% confidence level. Finding a non-zero value would provide crucial evidence for large CP violation.

2 Maths Pre-req

2.1 The Wigner-Eckart Theorem

The Wigner–Eckart theorem is a fundamental result in quantum mechanics that simplifies the calculation of matrix elements for spherical tensor operators. It allows you to separate the physical part of a transition (the dynamics) from the geometrical part (the angular momentum alignment).Mathematically, the theorem is expressed as:

$$\langle n'j'm'|T_q^{(k)}|njm\rangle = \frac{\langle n'j'\|T^{(k)}\|nj\rangle}{\sqrt{2j'+1}}\langle jm; kq|j'm'\rangle$$

For a fundamental particle in a state of definite angular momentum, any vector operator must be proportional to the total angular momentum (spin) operator $\mathbf{S}$. According to the Wigner-Eckart theorem, the expectation value of the EDM operator $\mathbf{d}$ is rigidly aligned with the spin axis:

$$\langle \mathbf{d}\rangle = d_n \frac{\langle \mathbf{S}\rangle}{S} \tag{1}$$

where d_n is the magnitude of the EDM.

2.2 P and T Violation

The interaction Hamiltonian for a neutron in external electric ($\mathbf{E}$) and magnetic ($\mathbf{B}$) fields is:

$$\mathcal{H} = -d_n \frac{\mathbf{S}}{S}\cdot\mathbf{E} - \mu_n \frac{\mathbf{S}}{S}\cdot\mathbf{B} \tag{2}$$

Under the discrete symmetries of Parity (P) and Time Reversal (T), these vectors transform differently. The electric field is a polar vector ($\mathbf{E} \xrightarrow{P} -\mathbf{E}$, $\mathbf{E} \xrightarrow{T} \mathbf{E}$), while spin is an axial pseudovector ($\mathbf{S} \xrightarrow{P} \mathbf{S}$, $\mathbf{S} \xrightarrow{T} -\mathbf{S}$).

Consequently, the EDM interaction term $-d_n\mathbf{S}\cdot\mathbf{E}$ changes sign under both P and T transformations. If a non-zero d_n exists, the system fundamentally violates both Parity and Time Reversal symmetries. Assuming the CPT theorem holds, T violation mathematically guarantees CP violation.

3 The Strong CP Problem and Naturalness

3.1 The θ-Vacuum in QCD

we can add the following term in lagrangian as it is lorentz invariant. This manifests in the QCD Lagrangian as a term that violates CP symmetry.this term is very small in nature.

$$\mathcal{L}_\theta = \bar{\theta}\frac{g_s^2}{32\pi^2}G^a_{\mu\nu}\tilde{G}^{\mu\nu,a} \tag{3}$$

where $G^a_{\mu\nu}$ is the gluon field strength tensor, $\tilde{G}^{\mu\nu,a}$ is its dual, and $\bar{\theta}$ is the effective CP-violating phase .We can add such term but we have to tell why is this term so small.

3.2 The Conflict and Naturalness

Theoretical models utilizing QCD sum rules relate this phase directly to the neutron EDM:

$$d_n \approx 3.2 \times 10^{-16}\, \bar{\theta}\, e \cdot \text{cm} \tag{4}$$

The experimental limit of $d_n < 1.8 \times 10^{-26}\, e \cdot \text{cm}$ restricts $\bar{\theta} \lesssim 10^{-10}$. In the Standard Model, $\bar{\theta}$ is a free parameter capable of taking any value from 0 to 2π. The fact that it is infinitesimally close to zero presents a severe "naturalness" problem,suggesting a missing fundamental symmetry. A theory is considered "natural" if its parameters (like masses or coupling constants) are roughly the same order of magnitude as the physical scales of the theory, or if their smallness is protected by a symmetry.
The 't Hooft Criterion.
The most formal definition comes from Gerard 't Hooft (1979). He stated:
A parameter is technically natural if setting it to zero increases the symmetry of the system.

If setting a parameter to zero restores a symmetry, then quantum corrections to that parameter will only be proportional to the parameter itself (logarithmic divergence), rather than the high-energy scale of the universe (quadratic divergence).

4 Calculating relation between d_n and θ

we do something similar to how we find the g-factor of the muon.we find it through comparing effective lagrangian.below is the operator that couples the neutron spin to the electromagnetic field in a CP-violating way

$$\mathcal{L}_{\text{EDM}} = -\frac{i}{2}\, d_n\, \bar{n}\, \sigma^{\mu\nu} \gamma_5\, n\, F_{\mu\nu}$$

while the scattering amplitude we will be calculating will be of form:

$$T = -2iD(q^2)\, \epsilon^*_\mu(q)\, \bar{u}_{s'}(p')\, S^{\mu\nu} q_\nu i\gamma_5\, u_s(p), \tag{94.25}$$

where $q = p' - p$. In the $q \to 0$ limit, this corresponds to a term in the effective Lagrangian of

$$\mathcal{L} = D(0)\, F_{\mu\nu}\, \bar{n}\, S^{\mu\nu} i\gamma_5\, n, \tag{94.26}$$

where n is the neutron field;If the factor of $i\gamma_5$ were absent, this would represent a contribution of $D(0)$ to the magnetic dipole moment of the neutron. To account for the factor of $i\gamma_5$, we use

$$S^{\mu\nu} i\gamma_5 = -\frac{1}{2}\, \epsilon^{\mu\nu\rho\sigma} S_{\rho\sigma}, \tag{94.27}$$

to see that eq. (94.26) is equivalent to

$$\mathcal{L} = -D(0)\, \tilde{F}_{\mu\nu}\, \bar{n}\, S^{\mu\nu}\, n. \tag{94.28}$$

We write the charged-pion interaction terms as, this can be found by using chiral perturbation theory.

$$\mathcal{L}_{\pi NN} = -i\sqrt{2}\left(\frac{g_A m_N}{f_\pi}\right)\left(\pi^+ \bar{p}\gamma_5 n + \pi^- \bar{n}\gamma_5 p\right), \tag{5}$$

$$\mathcal{L}_{\theta\pi NN} = -\sqrt{2}\left(\frac{\tilde{m} c_+}{f_\pi}\right)\left(\pi^+ \bar{p} n + \pi^- \bar{n} p\right). \tag{6}$$

From these we read off the pion-nucleon vertex factors as,

$$V_{\pi^+ pn} = -i\sqrt{2}\, \frac{g_A m_N}{f_\pi}\, \gamma_5$$

$$V_{\pi^+ pn} = -\sqrt{2}\, \frac{\theta\, c_+\, \tilde{m}}{f_\pi}$$

We label the internal momenta as shown in figure below; for small q, ℓ is a pion momentum that should be cut off at $\Lambda \sim 4\pi f_\pi$. Because this diagram have a infrared divergence at small ℓ, we can treat ℓ as

much less than p and p'.
You might wonder why above term which is not associated to weak interaction consist g_A ,this comes from Goldberger-Treiman relation.(this is derived in **Peskin and schroder** ,Chapter 19),At zero momentum transfer ($q^2 = 0$):

$$g_A = \frac{f_\pi \, g_{\pi NN}}{m_N} \, .$$

Thus the internal proton is nearly on-shell, which justifies the use of eq. (94.24) for the CP conserving interaction.



Figure 1: required feynman diagram

The diagram yield an amplitude of

$$\begin{aligned} iT = \left(\frac{i}{\hbar}\right)^3 (ie) \left(\sqrt{2} g_A m_N / f_\pi\right) \left(-i\sqrt{2}\delta c_+ \tilde{m}/f_\pi\right) \epsilon^*_\mu \int^\Lambda \frac{d^4\ell}{(2\pi)^4} \\ \times \frac{(2\ell^\mu)\, \bar{u}' \left[(-\not{\ell} - \not{\bar{p}} + m_N)\gamma_5 + \gamma_5(-\not{\ell} - \not{\bar{p}} + m_N)\right] u}{\left((\ell+\bar{p})^2 + m_N^2\right)\left((\ell + \frac{1}{2}q)^2 + m_\pi^2\right)\left((\ell - \frac{1}{2}q)^2 + m_\pi^2\right)}, \end{aligned} \tag{7}$$

where $\bar{p} = \frac{1}{2}(p' + p)$. Using $\{\gamma^\mu, \gamma_5\} = 0$, the spinor factors in the numerator simplify to

$$\bar{u}'[\ldots]u = 2m_N \, \bar{u}'\gamma_5 u. \tag{8}$$

From the spinor properties, $\bar{u}'\gamma_5 u$ vanishes when $p' = p$; thus $\bar{u}'\gamma_5 u$ must be $O(q)$, and so we can set $q = 0$ everywhere else. taking $\ell \ll p$, we can set $(\ell + \bar{p})^2 + m_N^2 = 2p \cdot \ell$ in the denominator of eq. (7). We now have

$$T = 4\left(\frac{e\theta g_A c_+ \tilde{m} m_N^2}{f_\pi^2}\right) \epsilon^*_\mu \int_0^\Lambda \frac{d^4\ell}{(2\pi)^4} \frac{(2\ell^\mu)\, \bar{u}'\gamma_5 u}{(2p\cdot\ell)\,(\ell^2 + m_\pi^2)^2}. \tag{9}$$

Integrating over the directions of ℓ results in

$$\frac{\ell^\mu}{p\cdot\ell} \;\to\; \frac{p^\mu}{p^2} = -\frac{p^\mu}{m_N^2}. \tag{10}$$

Next we use the Gordon identity,

$$p^\mu \, \bar{u}'\gamma_5 u = \bar{u}' S^{\mu\nu} q_\nu \gamma_5 u + O(q^2), \tag{11}$$

which verifies that $\bar{u}'\gamma_5 u$ is linear in q. We now have

$$T = -4\left(\frac{e\theta g_A c_+ \tilde{m}}{f_\pi^2}\right) \epsilon^*_\mu \, \bar{u}' S^{\mu\nu} q_\nu \gamma_5 u \int_0^\Lambda \frac{d^4\ell}{(2\pi)^4} \frac{1}{(\ell^2 + m_\pi^2)^2}. \tag{12}$$

In the limit $m_\pi \to 0$, the integral diverges at small ℓ, generating a chiral log. This infrared divergence can only arise from diagrams with two pion propagators, which is why it appears only if the photon is attached to the pion.

After a Wick rotation, the integral evaluates to $(i/16\pi^2)\ln\left(\Lambda^2/m_\pi^2\right)$.Comparing with eq. (94.25), we see that the electric dipole moment of the neutron is

$$d_n = \frac{e\theta g_A c_+ \tilde{m}}{8\pi^2 f_\pi^2}\left[\ln\left(\Lambda^2/m_\pi^2\right) + O(1)\right]. \tag{13}$$

Putting in numbers ($g_A = 1.27$, $c_+ = 1.7$, $\tilde{m} = 1.2$ MeV), we find

$$d_n = 3.2 \times 10^{-16}\ e \cdot \text{cm}. \tag{14}$$

5 The Peccei-Quinn Mechanism and the Axion

The most elegant solution to the Strong CP problem is the Peccei-Quinn (PQ) mechanism. It postulates a new global, chiral $U(1)_{PQ}$ symmetry that is spontaneously broken at a high energy scale f_a.

According to the Goldstone theorem, this spontaneous symmetry breaking yields a pseudo-scalar particle: the axion (a). The static parameter $\bar{\theta}$ is promoted to a dynamical field:

$$\theta_{\text{eff}}(x) = \bar{\theta} + \frac{a(x)}{f_a} \tag{5}$$

Non-perturbative QCD instanton effects generate a periodic potential for the axion field:

$$V(a) \approx \Lambda_{\text{QCD}}^4 \left[1 - \cos\left(\bar{\theta} + \frac{a(x)}{f_a}\right)\right] \tag{6}$$

The field dynamically relaxes to the minimum of this potential, where $\bar{\theta} + a(x)/f_a = 0$. This perfectly cancels the CP-violating term, naturally restoring CP symmetry in the strong sector and leaving behind a **light axion**, Tadaaa XD.

6 Baryon Asymmetry of the Universe (BAU)

The observable universe consists almost entirely of matter. To generate this Baryon Asymmetry dynamically from an initially symmetric state, the three Sakharov conditions must be met:

1. Baryon number (B) violation.
2. C and CP violation.
3. Interactions outside of thermal equilibrium.

It is usually quantified by the baryon–to–photon ratio:

$$\eta \equiv \frac{n_B - n_{\bar{B}}}{n_\gamma} \sim 6 \times 10^{-10}$$

B is baryon and $\bar{B}$ is anti baryon,as measured independently by primordial nucleosynthesis and the cosmic microwave background.
Most accepted mechanism suggest that in early universe there exist heavy right-handed Majorana neutrinos.which further decay into a lepton and a Higgs doublet.in this process CP violation occured resulting in Baryon Asymmetry

The CP violation parameterized by the CKM matrix in the Standard Model is insufficient by several orders of magnitude to account for the observed BAU. A non-zero nEDM would indicate undiscovered sources of CP violation, offering a mechanism for baryogenesis (e.g., via leptogenesis frameworks).

7 Experimental Detection Methodologies

7.1 intro

To measure Larmor frequency we use Ramsey's method of separated oscillatory fields.this method gives more acurate and precise measurements compared to Rabi's resonance method.Below we try to find electric diople moment ,But as you will see this method can be used to measure magnetic dipole moemnt as well.As fact of matter, In 1949, Norman Ramsey (student of I.I. Rabi) realized that the standard "Rabi resonance" was limited by the uniformity of the magnetic field over the entire interaction length.he proposed the separated oscillatory fields method, and in 1951, he published the results of its application to the neutron.although it the first accurate, direct measurement of the magnetic moment of a free neutron was performed by Luis Alvarez and Felix Bloch in 1940, Finding magnetic dipole moment was amazing ,it tells us that neutron is a composite particle and its magnetic moment is the result of its complex structure of quarks.

7.2 Measuring the nEDM and nMDM (Ramsey Resonance)

experiments utilize Ultracold Neutrons (UCNs) stored in highly shielded vacuum chambers. The measurement relies on the Ramsey method of separated oscillatory fields to determine the Larmor precession frequency of the neutron spin.

Neutrons are subjected to perfectly aligned parallel and anti-parallel magnetic (**B**) and electric (**E**) fields. The resonant frequencies for both configurations are:

$$h\nu_{\pm} = 2\mu_n B \pm 2d_n E \tag{7}$$

By calculating the difference between these two frequencies, the magnetic moment term cancels out, isolating the EDM:

$$\Delta\nu = \nu_+ - \nu_- = \frac{4d_n E}{h} \tag{8}$$

7.3 Ramsey's method

Consider a spin-$\frac{1}{2}$ particle in a uniform magnetic field

$$\mathbf{B}_0 = B_0\hat{z}.$$

The Hamiltonian is

$$\hat{H}_0 = -\boldsymbol{\mu}\cdot\mathbf{B}_0 = -\gamma\hbar B_0\hat{S}_z = -\frac{\hbar\omega_0}{2}\sigma_z,$$

where γ is the gyromagnetic ratio and

$$\omega_0 = \gamma B_0$$

is the Larmor frequency.

The energy eigenstates satisfy

$$E_{\uparrow,\downarrow} = \mp\frac{\hbar\omega_0}{2}.$$

A transverse oscillating magnetic field is applied:

$$\mathbf{B}_1(t) = B_1\big(\hat{x}\cos\omega t - \hat{y}\sin\omega t\big).$$

The interaction Hamiltonian is

$$\hat{H}_{\mathrm{RF}}(t) = -\gamma\hbar B_1\big(\hat{S}_x\cos\omega t - \hat{S}_y\sin\omega t\big).$$

In the frame rotating at angular frequency ω, and within the rotating-wave approximation, the effective Hamiltonian becomes

$$\hat{H}_{\mathrm{rot}} = -\frac{\hbar}{2}\left[(\omega_0-\omega)\sigma_z + \omega_1\sigma_x\right],$$

where

$$\omega_1 = \gamma B_1$$

is the Rabi frequency.

At resonance ($\omega = \omega_0$),

$$\hat{H}_{\text{rot}} = -\frac{\hbar\omega_1}{2}\sigma_x,$$

which generates a rotation about the x-axis.

The time-evolution operator is

$$\hat{U}(t) = \exp\left(-\frac{i}{\hbar}\hat{H}_{\text{rot}}t\right) = \exp\left(i\frac{\omega_1 t}{2}\sigma_x\right).$$

This corresponds to a rotation by angle

$$\theta = \omega_1 t.$$

A $\pi/2$ pulse is defined by

$$\theta = \frac{\pi}{2} \quad \Rightarrow \quad t_{\pi/2} = \frac{\pi}{2\omega_1} = \frac{\pi}{2\gamma B_1}.$$

State After the First $\pi/2$ Pulse

Assuming the initial state

$$|\psi(0)\rangle = |\uparrow\rangle,$$

the $\pi/2$ pulse produces

$$|\psi\rangle = e^{i\frac{\pi}{4}\sigma_x}|\uparrow\rangle = \frac{1}{\sqrt{2}}\left(|\uparrow\rangle + i|\downarrow\rangle\right).$$

The spin now lies in the transverse plane.

After the RF is turned off, the system evolves under $\hat{H}_0$:

$$\hat{H}_0 = -\frac{\hbar\omega_0}{2}\sigma_z.$$

After a free-precession time T,

$$|\psi(T)\rangle = e^{i\frac{\omega_0 T}{2}\sigma_z}|\psi\rangle = \frac{1}{\sqrt{2}}\left(|\uparrow\rangle + ie^{i\omega_0 T}|\downarrow\rangle\right).$$

The accumulated relative phase is therefore

$$\phi = \omega_0 T.$$

A small frequency shift $\delta\omega$ leads to

$$\delta\phi = \delta\omega\, T,$$

Applying a second $\pi/2$ pulse converts the phase ϕ into a population difference:

$$P_\uparrow = \frac{1}{2}\big(1 + \cos\phi\big), \qquad P_\downarrow = \frac{1}{2}\big(1 - \cos\phi\big),$$

producing the Ramsey interference fringes.

7.4 summary of experiment

1. Polarize UCN: pass the ultracold neutrons through a 5 T superconducting magnet to strongly polarize the beam by reflecting one spin state back toward the source, leaving an almost single-spin population generally the one aligned with Magnetic field B.use adiabatic fast-passage spin flipper for initial polarization.

2. Fill chamber:: since neutron has magnetic dipole moment it can be transported using non uniform magnetic field.by this deflect polarized UCN into the Ramsey precession chamber for an optimized filling time ($\sim$ 28 s) to maximize usable counts and polarization, then close the chamber .

3. Admit co-magnetometer: inject polarized ^{199}Hg into chamber ($\sim$ 2 s).for providing a continuous reference for magnetic-field tracking.

4. Start Ramsey precession: apply $\pi/2$ pulses to Hg ($\sim$ 8 Hz) and neutrons ($\sim$ 30 Hz).

5. Free precession: wait $\sim$ 180 s; continuously read out Hg frequency optically (UV beam) to track B_0 drifts.

6. Second Ramsey pulse: apply the final, phase-coherent neutron $\pi/2$ pulse to convert the accumulated phase (frequency detuning) into a measurable spin-population asymmetry that is maximally sensitive near resonance.

7. Spin analysis: send neutrons through polarized iron foil; use SF2A/SF2B to swap analyzed spin states between detector arms.

8. Field reversals: item Field reversals: reverse the high voltage to flip E (minutes timescale) and periodically reverse B_0 (hours timescale) to isolate genuine E-correlated shifts; form the ratio $R = f_n/f_{\mathrm{Hg}}$ to suppress common-mode magnetic-field variations and extract an EDM-sensitive signal.

9. Run structure: group cycles into stable field-configuration sets; apply blinding (if used) at analysis stage.

by Theory this experiment is quite simple but to actualy measure this quantity their goes lot of equipment and calculations to do it.

7.5 Set Up

- **Neutron source and transport:** prepare a well-defined initial neutron spin state and allow controlled spin flips for systematic checks.CryoEDM: the incoming cold-neutron beam is polarized with a long polariser that also filters wavelengths (CryoEDM). The produced UCN must then maintain polarization through guides and valves, requiring low stray transverse fields and smooth magnetic holding fields along transport.in PSI, a 5 T superconducting magnet polarizes UCN by reflecting one spin state. SF1 selects the initial spin state. SF2A/SF2B in the detector arms allow alternating which spin state is counted by which detector, reducing detector-efficiency asymmetry systematics.

- **UCN production and storage:** CryoEDM stores UCN in two adjacent BeO cylindrical cells (about 2.2 L each) capped with Ti electrodes coated with 1 m diamond-like carbon (DLC) (CryoEDM). The electrode arrangement sets one cell with an electric field while the neighboring cell is held at neutral electric field. Analyzing both cells separately provides an internal control against systematic effects that could mimic an EDM.

- **Precession (Ramsey) chamber:** UCN stored for 100–200 s in a highly uniform B_0, with an applied high-voltage electric field.: CryoEDM generates a holding field of about 50 mG using a superconducting NbTi solenoid, with additional trim coils to improve homogeneity (CryoEDM). The storage cells sit in a superfluid containment vessel made of non-magnetic titanium alloy at the center of the magnetic shields.

- **Magnetic environment control:** CryoEDM uses multiple shielding layers: three layers of mu-metal plus a superconducting lead screen held at 4 K (CryoEDM). The lead screen provides strong attenuation of time-varying external magnetic fields once superconducting. The holding-field solenoid is placed just inside the superconducting screen, and trim coils around the storage chambers are used to tune gradients and improve uniformity.

- **Magnetometry:** cohabiting magnetometers (e.g., ^{199}Hg at PSI) and external sensor arrays (e.g., Cs magnetometers), and SQUID systems to read and subtract out any dipole due to magnetic field fluctuations.In neutron EDM experiments, polarized Hg atoms are stored in the same precession



Figure 1. Sketch of the spectrometer used to search for an electric-field induced shift in the magnetic resonance frequency of polarized UCN exposed to an electric field of strength E = 11 kV/cm and a magnetic field of B_0 = 1.035 μT.

chamber as neutrons and undergo Ramsey precession in the same magnetic field. Mercury acts as a co-magnetometer, continuously measuring magnetic-field fluctuations via optical readout. By forming the frequency ratio $R = \frac{\omega_n}{\omega_{Hg}}$ first-order magnetic-field drifts cancel, allowing the tiny electric-field-dependent shift in neutron precession frequency to be isolated.

- **Spin manipulation and readout:** RF pulses for Ramsey sequence; spin analyzers and UCN detectors to count each spin state at the end of storage.the incoming cold-neutron beam is polarized with a long polariser that also filters wavelengths (CryoEDM). The produced UCN must then maintain polarization through guides and valves, requiring low stray transverse fields and smooth magnetic holding fields along transport.in PSI, a 5 T superconducting magnet polarizes UCN by reflecting one spin state. SF1 selects the initial spin state. SF2A/SF2B in the detector arms allow alternating which spin state is counted by which detector, reducing detector-efficiency asymmetry systematics.

- **Detector:** UCN detectors and polarisation measurement.The detector system consists of a set of solid state surface barrier detectors covered with ^{6}LiF. The signal in the detector is generated by the alpha and triton particles from the $n(^6Li, ^3H)^4He$ reaction. This is a system that has been shown to work efficiently within superfluid helium. The detectors are placed in a magnetic field that the UCN will have followed adiabatically. Two of the detectors have an additional Fe layer on top of the ^6Li coating that will reflect one spin state of the UCN. The ratio in neutron counts in the detectors with Fe layer to the counts in the detectors without Fe layer is a measure of the UCN polarisation.

- **Optical Readout of the ^{199}Hg Co-Magnetometer**

  In neutron electric dipole moment (nEDM) experiments, polarized ^{199}Hg atoms are stored in the same precession chamber as the neutrons and act as a co-magnetometer for measuring the magnetic field B_0.

  Magnetic Precession of Mercury

  The nuclear spin of ^{199}Hg is $I = \frac{1}{2}$. In a uniform magnetic field

$$\mathbf{B}_0 = B_0\hat{z},$$

the mercury spins precess at the Larmor frequency

$$\omega_{\mathrm{Hg}} = \gamma_{\mathrm{Hg}} B_0,$$

where γ_{Hg} is the gyromagnetic ratio of ^{199}Hg.

After polarization, the transverse spin component evolves as

$$S_{\perp}(t) \propto \cos(\omega_{\mathrm{Hg}} t + \phi_0).$$

Mercury atoms possess an allowed optical transition

$$^1S_0 \;\rightarrow\; ^3P_1$$

at wavelength $\lambda \approx 254$ nm. A linearly polarized ultraviolet (UV) probe beam is sent through the precession chamber.

The atomic absorption probability depends on the projection of the nuclear spin along the light polarization axis. As the mercury spin precesses, this projection oscillates, producing a time-dependent absorption of the probe light.

Let

- I_0 be the incident light intensity,
- n the mercury atom density,
- L the optical path length,
- σ the absorption cross section,
- $P(t)$ the normalized spin projection.

The transmitted intensity is

$$I(t) = I_0 \exp\big[-n\sigma(1 + \alpha P(t))L\big],$$

where α characterizes the spin-dependent absorption.

For small absorption, this reduces to

$$I(t) \approx I_0 \left[1 - A\cos(\omega_{\mathrm{Hg}} t + \phi)\right],$$

with modulation amplitude $A \propto \alpha$.

Thus the detected optical signal oscillates directly at the mercury Larmor frequency.

- **Controls and DAQ:** high-voltage supply and reversal logic; timing sequencer; detector readout; blinding and data-quality monitoring (PSI strategy).

7.6 Cooling process

Bring apparatus to superfluid-helium temperatures (~ 0.7 K in measurement region). Purify helium: prepare isotopically pure ^{4}He using superleaks to suppress UCN loss via He absorption (target $^3\mathrm{He} < 10^{-10}$). Configure UCN source: send ~ 9 Å cold neutrons through superthermal source volume to produce UCN by downscattering (single-phonon process).Guide & store UCN open cryogenic valves to fill storage cells through Be-coated guides; close valves to define storage time.
Apply electric-field configuration: energize electrodes so one cell is $E \neq 0$ while neighboring cell is $E = 0$ (systematic monitor).
Ramsey sequence: apply first $\pi/2$ RF pulse, wait free-precession time T, then apply second $\pi/2$ pulse.record SQUID/fluxgate signals to track drifts/gradients near cells. Detect UCN: open valves to send UCN to in-superfluid LiF-coated silicon detectors; use Fe-coated channels (if enabled) for polarization analysis. Reverse and repeat: repeat cycles with electric-field polarity reversed; extract EDM from E-correlated frequency shift, using $E = 0$ cell as control.

7.7 Check check

1. Verify cryogenic readiness: confirm liquid-helium inventories for shield cryostats and readiness of ^{3}He circulation system (CryoEDM cryostat reaches $\sim$ 420 mK).

2. Magnetic shields and coils: ensure mu-metal layers and superconducting lead shield are at operating temperature ($\sim$ 4 K for lead screen); solenoid and trim coils powered and stable.

3. High-voltage readiness: verify electrode connections, insulation distances, interlocks, and ramp-rate limits for liquid/superfluid helium operation.

4. Detector readiness: confirm silicon surface-barrier detectors with ^{6}LiF conversion layers are biased and reading properly; check Fe foils if used.

5. UCN-valve checks: cycle superfluid-operable UCN valves to confirm sealing and timing reproducibility.

6. Mercury atoms form a thermal gas and sample the full chamber volume, while ultracold neutrons preferentially sample the lower part of the chamber due to gravity. In the presence of magnetic-field gradients,
$$\langle B\rangle_{\mathrm{Hg}} \neq \langle B\rangle_n,$$
which can lead to false EDM signals and must be carefully controlled.

7. we have approx atandard deviation given by:this is statistical error only because this experiment are designed so beautifully that $\sigma_{exp} << \sigma_{stat}$

$$\sigma_{stat}(d_n^{\mathrm{meas}}) = \frac{\hbar}{2\alpha T E\sqrt{N}\,\sqrt{N_{\mathrm{cycles}}}}$$

What is measured; The transmitted light intensity is recorded continuously by a UV photodetector during the neutron free-precession time. A sinusoidal fit yields ω_{Hg}, from which the magnetic field is obtained as
$$B_0(t) = \frac{\omega_{\mathrm{Hg}}(t)}{\gamma_{\mathrm{Hg}}}.$$

This provides a non-destructive, real-time measurement of magnetic-field fluctuations.
In above experiment ΔR is measured while flipping E and B. The neutron Larmor frequency is

$$\omega_n = \gamma_n B_0 \pm \frac{2d_n E}{\hbar}.$$

Forming the frequency ratio
$$R = \frac{\omega_n}{\omega_{\mathrm{Hg}}} = \frac{\gamma_n}{\gamma_{\mathrm{Hg}}} \pm \frac{2d_n E}{\hbar\gamma_{\mathrm{Hg}} B_0},$$
cancels first-order magnetic-field fluctuations.

8 Axion Detection

Let say if we by chance get to find an EDM,By PQ mech,we predicted a goldstone particle **Axion**.So now,we give ourselves more work.Detecting Axion use Primakoff effect, whereby an axion couples to two photons. In the presence of a strong, static magnetic field, an axion can convert into a detectable microwave photon:
$$\mathcal{L}_{a\gamma\gamma} = -g_{a\gamma\gamma}\, a\, \mathbf{E}\cdot\mathbf{B} \tag{9}$$

In a strong, static magnetic field $\mathbf{B}_0$, one of the photons at the $a\gamma\gamma$ vertex is effectively supplied by the external field. As a result, an axion propagating through the magnetic field can convert into a real photon with the same energy and momentum, up to small corrections determined by the axion mass. *Haloscopes* are designed to detect axions constituting the local dark matter halo. In these experiments,

a tunable microwave cavity is placed inside a strong superconducting magnetic field. Galactic axions passing through the apparatus convert into microwave photons via the Primakoff effect.

If the cavity resonance frequency ν_c matches the axion mass,

$$\nu_c = \frac{m_a c^2}{h}, \tag{10}$$

the conversion is resonantly enhanced. The expected signal power in a haloscope is

$$P \propto g_{a\gamma\gamma}^2 \, B_0^2 \, V \, C \, Q \, \rho_{\mathrm{DM}}, \tag{11}$$

where V is the cavity volume, Q its quality factor, C a mode-dependent form factor describing the overlap with the magnetic field, and ρ_{DM} the local dark matter density.

Since the axion mass is unknown, haloscopes must scan over frequency by tuning the cavity. **Haloscopes** (such as ADMX) use tunable microwave cavities immersed in superconducting magnets to search for resonant photon conversions from the ambient dark matter halo.

$$\gamma + Z \to a + Z. \tag{12}$$

In these experiments, a long, strong dipole magnet is pointed toward the Sun. Solar axions entering the magnet convert back into photons, typically in the X-ray energy range, which are detected at the ends of the magnet.

The axion–photon conversion probability over a magnet of length L is given by

$$P_{a\to\gamma} \propto (g_{a\gamma\gamma} B_0 L)^2 \, \frac{\sin^2(qL/2)}{(qL/2)^2}, \qquad q = \frac{m_a^2}{2E}, \tag{13}$$

where E is the axion energy. For sufficiently small axion masses, $qL \ll 1$, the conversion is coherent and maximal. At higher masses, coherence can be restored by introducing a buffer gas, which gives the photon an effective plasma mass.

9 Question asked during class:

here are some question which i remembered where asked me during class.:)
Q;How does spin flipper in image work??
A: It is more like a filter, what it does is allow specifically spin oriented neutron to pass.since,The polarizer gives mostly one spin state. The spin flipper allows you to switch which spin state you inject into the chamber because we need spin aligned to B as discussed above.
With flipper OFF $\to$ spin-up neutrons
With flipper ON $\to$ spin-down neutrons

Q:Why is their so many neutron??is it macroscopic experiment and what is size of equipment? A:The effect is obviously on level of particle itself,but ramsey's method uses population difference arised from phase of spins due to EDM.Well about the size when i saw the image i thought it was as large as a room, but effectively the size of cylindrical chamber where measurements are taken is:
Diameter: $\approx$ 45–50 cm
Height: $\approx$ 12 cm

10 Conclusion

Neutron EDM which is yet to find is a experiment that can explain big problems that standard model is currently facing.We by brute calculation have found the relation between the nEDM and θ by feynman diagram,of course we can reduce uncertainty by calculating higher order diagrams as done for g-2.while people at PSI are trying to find the observed value of d_n Using Ramsey's method.With the same method physicist have not only found Magnetic dipole moment of neutron and other particle but also to very very precise number(i.e. Least count of 10^{-18}).As prof. said you might look on surface and say we know standard model but their goes a lot of Calculation to find even a Simple relation and hardwork to Engineer this beautiful experiment.

References

[1] C. A. Baker et al., "CryoEDM: A cryogenic experiment to measure the neutron electric dipole moment," *Journal of Physics: Conference Series* **251**, 012055 (2010). doi:10.1088/1742-6596/251/1/012055

[2] C. Abel et al., "nEDM experiment at PSI: Data-taking strategy and sensitivity of the dataset," arXiv:1811.04012 [physics.ins-det] (2018).

[3] J. Engel, M. J. Ramsey-Musolf, and U. van Kolck, "Electric dipole moments of nucleons, nuclei, and atoms: The Standard Model and beyond," *Progress in Particle and Nuclear Physics* **71**, 21–74 (2013). arXiv:1303.2371 [nucl-th].

11 Pion Beta Decay and Testing CKM Unitarity

Pion beta decay, $\pi^+ \to \pi^0 e^+ \nu_e$, is a rare semileptonic weak decay that provides one of the cleanest probes of the Standard Model (SM). Unlike nuclear beta decay, it is free from nuclear structure uncertainties and thus allows a theoretically clean extraction of the CKM matrix element V_{ud}. The decay is a pure vector ($0^- \to 0^-$) transition governed by the conserved vector current (CVC) hypothesis, making it an ideal laboratory for testing CKM unitarity, studying radiative corrections, probing physics beyond the Standard Model.



11.1 $\pi\beta$ Decay Rate

At low energies, the charged-current weak interaction is described by the Fermi effective Hamiltonian:

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{ud} \left(\bar{u}\gamma^\mu(1-\gamma^5)d\right) \left(\bar{e}\gamma_\mu(1-\gamma^5)\nu_e\right) \tag{14}$$

This interaction mediates the transition between quark and lepton sectors. The hadronic matrix Element is

$$\mathcal{M} = \frac{G_F}{\sqrt{2}} V_{ud} \langle \pi^0(p')|J^\mu|\pi^+(p)\rangle \bar{u}_\nu \gamma_\mu (1-\gamma^5) v_e \tag{15}$$

The current is:

$$J^\mu = V^\mu - A^\mu \tag{16}$$

Since both initial and final states are pseudoscalars (0^-), parity conservation implies:

$$\langle \pi^0|A^\mu|\pi^+\rangle = 0 \tag{17}$$

Thus only the vector current contributes. Lorentz invariance implies:

$$\langle \pi^0|V^\mu|\pi^+\rangle = f_+(q^2)(p+p')^\mu + f_-(q^2)q^\mu \tag{18}$$

where:

$$q^\mu = (p-p')^\mu \tag{19}$$

Using the Dirac equation:

$$q^\mu \bar{u}_\nu \gamma_\mu (1-\gamma^5) v_e = -m_e \bar{u}_\nu (1+\gamma^5) v_e \tag{20}$$

Thus:

$$f_- \text{ term is suppressed by } m_e \Rightarrow f_- \approx 0 \tag{21}$$

The conserved vector current hypothesis implies:

$$f_+(0) = \sqrt{2} \tag{22}$$

This follows from isospin symmetry relating weak and electromagnetic currents. The squared amplitude:

$$|\mathcal{M}|^2 = \frac{G_F^2 |V_{ud}|^2}{2} |f_+(q^2)|^2 L_{\mu\nu} H^{\mu\nu} \tag{23}$$

After phase space integration, we obtain:

$$\Gamma_{\pi\beta} = \frac{G_F^2 |V_{ud}|^2 M_\pi^5}{64\pi^3} |f_+(0)|^2 I_{\pi\beta} \tag{24}$$

where $I_{\pi\beta}$ is the phase space integral.

11.2 Radiative Corrections in Pion Beta Decay

For a precision extraction of V_{ud} from pion beta decay, the tree-level decay amplitude is not sufficient. The decay

$$\pi^+(p) \to \pi^0(p')\, e^+(p_e)\, \nu_e(p_\nu) \tag{25}$$

is a semileptonic weak process, and once one aims at sub-percent or per-mille precision, electroweak radiative corrections become indispensable. These corrections arise from several physically distinct sources. At high momentum scales, one has virtual electroweak corrections associated with the running from the weak scale down to a hadronic matching scale. At low momentum scales, one has electromagnetic loop effects, real-photon emission, and hadronic-structure-dependent contributions associated with the fact that the pion is not an elementary point particle. The physically measurable quantity is therefore not the naive tree-level width, but the inclusive decay width

$$\Gamma_{\pi\beta}^{\rm incl} = \Gamma(\pi^+ \to \pi^0 e^+ \nu_e) + \Gamma(\pi^+ \to \pi^0 e^+ \nu_e \gamma) + \cdots\,, \tag{26}$$

where the real-photon channel must be included to obtain an infrared-finite result.

At tree level, the low-energy charged-current interaction is described by the Fermi Hamiltonian

$$\mathcal{H}_{\rm eff} = \frac{G_F}{\sqrt{2}}\, V_{ud} \left[\bar{u}\gamma^\mu(1-\gamma_5)d\right] \left[\bar{e}\gamma_\mu(1-\gamma_5)\nu_e\right] + \text{h.c.} \tag{27}$$

Because both the initial and final hadrons are pseudoscalars, the axial current does not contribute to the hadronic matrix element, and one may write

$$\langle \pi^0(p')|\bar{u}\gamma^\mu d|\pi^+(p)\rangle = f_+(q^2)(p+p')^\mu + f_-(q^2)q^\mu, \qquad q^\mu = (p-p')^\mu. \tag{28}$$

The term proportional to $f_-(q^2)$ is suppressed by the electron mass, since contraction with the leptonic current gives

$$q_\mu\, \bar{u}_\nu \gamma^\mu (1-\gamma_5) v_e = -\, m_e\, \bar{u}_\nu (1+\gamma_5) v_e, \tag{29}$$

and is therefore negligible for the precision level considered here. The dominant tree-level amplitude is thus

$$\mathcal{M}_0 = \frac{G_F}{\sqrt{2}}\, V_{ud}\, f_+(q^2)\, (p+p')_\mu\, \bar{u}_\nu \gamma^\mu (1-\gamma_5) v_e. \tag{30}$$

In the exact isospin limit, the conserved vector current hypothesis implies

$$f_+(0) = \sqrt{2}, \tag{31}$$

and the tree-level width may then be expressed in the standard form

$$\Gamma_{\pi\beta}^{(0)} = \frac{G_F^2 |V_{ud}|^2 M_{\pi^+}^5}{64\pi^3}\, |f_+(0)|^2\, I_{\pi\beta}, \tag{32}$$

where $I_{\pi\beta}$ denotes the three-body phase-space integral. The goal of the radiative-correction program is to replace this by

$$\Gamma_{\pi\beta} = \Gamma_{\pi\beta}^{(0)}\, (1 + \Delta_{\rm RC}), \tag{33}$$

with $\Delta_{\rm RC}$ computed in a systematic and physically transparent way.

A standard and very useful decomposition is

$$1 + \Delta_{\rm RC} = S_{\rm EW}\, (1 + \delta_{\rm EM}), \tag{34}$$

where $S_{\rm EW}$ is the universal short-distance electroweak factor, while $\delta_{\rm EM}$ contains the long-distance electromagnetic and hadronic-structure corrections. This decomposition is conceptually important because it separates perturbatively calculable high-scale physics from genuinely low-energy strong-interaction effects.

11.2.1 Short-distance electroweak correction

The short-distance factor $S_{\rm EW}$ arises when one matches the full Standard Model onto the low-energy four-fermion theory. In the full theory, the semileptonic charged-current amplitude receives one-loop corrections from virtual W, Z, and photon exchange, as well as from γW box diagrams. Once the heavy electroweak degrees of freedom are integrated out, these effects are encoded in the Wilson coefficient of the effective semileptonic operator. The leading universal contribution has the familiar logarithmic structure identified by Sirlin:

$$S_{\rm EW} = 1 + \frac{2\alpha}{\pi} \ln \frac{M_Z}{M_\rho} + \cdots , \tag{35}$$

where M_Z is the weak scale and M_ρ is a hadronic matching scale of order the rho-meson mass. The logarithm originates from the integration over virtual momenta between the weak scale and the hadronic scale, and is universal in the sense that it depends only on the semileptonic charged current, not on the detailed internal structure of the hadron.

To understand this more explicitly, one may think of the renormalized semileptonic amplitude as

$$\mathcal{M}_{\rm ren} = \frac{G_F}{\sqrt{2}} V_{ud} \left[1 + \delta_{\rm EW}^{\rm SD}(\mu)\right] \langle \pi^0 | \bar{u}\gamma^\mu d | \pi^+ \rangle \, \bar{u}_\nu \gamma_\mu (1-\gamma_5) v_e, \tag{36}$$

with

$$\delta_{\rm EW}^{\rm SD}(\mu) = \frac{\alpha}{\pi}\Big[c_0 + c_1 \ln(M_Z/\mu) + \cdots\Big]. \tag{37}$$

The matching scale μ is chosen in the hadronic regime so that all remaining low-energy physics is absorbed into the long-distance part. The scale dependence of the short-distance coefficient is compensated by that of the low-energy matrix element and counterterms, leaving the full physical amplitude scale independent. In practice, $S_{\rm EW}$ provides a multiplicative enhancement of order a few percent, and it is crucial because the extraction of V_{ud} is linearly sensitive to it.

11.2.2 Long-distance electromagnetic correction: general structure

The long-distance part $\delta_{\rm EM}$ is more subtle. It includes the effects of virtual photons, real bremsstrahlung, and local structure-dependent terms required by renormalization. Schematically, the corrected amplitude can be written as

$$\mathcal{M} = \mathcal{M}_0 + \mathcal{M}_{\rm virt}^\gamma + \mathcal{M}_{\rm ct} + \mathcal{M}_{\rm real}, \tag{38}$$

where $\mathcal{M}_{\rm virt}^\gamma$ denotes virtual-photon loop corrections, $\mathcal{M}_{\rm ct}$ denotes local counterterms, and $\mathcal{M}_{\rm real}$ corresponds to the radiative channel $\pi^+ \to \pi^0 e^+ \nu_e \gamma$. The decay rate is then obtained from

$$d\Gamma = \frac{1}{2M_{\pi^+}} \sum_{\rm spins} \left(|\mathcal{M}_0 + \mathcal{M}_{\rm virt}^\gamma + \mathcal{M}_{\rm ct}|^2 \, d\Phi_3 + |\mathcal{M}_{\rm real}|^2 \, d\Phi_4 \right), \tag{39}$$

where $d\Phi_3$ and $d\Phi_4$ are the three-body and four-body phase-space elements, respectively. Expanding to first order in α, one finds

$$d\Gamma = d\Gamma_0 \left[1 + 2\,{\rm Re}\left(\frac{\mathcal{M}_{\rm virt}^\gamma + \mathcal{M}_{\rm ct}}{\mathcal{M}_0}\right)\right] + d\Gamma_{\rm real}. \tag{40}$$

The long-distance radiative correction may therefore be split into three pieces,

$$\delta_{\rm EM} = \delta_{\rm virt} + \delta_{\rm real} + \delta_{\rm local}, \tag{41}$$

but only the sum is physically meaningful, because the individual virtual and real parts are infrared divergent.

11.2.3 Virtual-photon loop corrections

The virtual-photon corrections come from several classes of one-loop diagrams. First, there are wave-function renormalization diagrams for the charged pion and the positron. Second, there are electromagnetic vertex corrections in which a virtual photon connects the charged pion leg to the positron leg. Third, there are hadronic electromagnetic corrections to the semileptonic form factor itself. At the point-particle level, the generic structure of the virtual contribution is

$$\mathcal{M}^{\gamma}_{\rm virt} = \mathcal{M}_0 \, \frac{\alpha}{4\pi} \, F_{\rm virt}(E_e, q^2, m_e, M_{\pi^+}, \lambda), \tag{42}$$

where λ is an infrared regulator, often introduced as a fictitious photon mass. The function $F_{\rm virt}$ contains both ultraviolet and infrared singularities.

To illustrate the structure, consider the photon exchange between the external positron and the charged pion. The corresponding loop integral is of the form

$$I_\mu = e^2 \int \frac{d^d k}{(2\pi)^d} \frac{N_\mu(k)}{\left[k^2 - \lambda^2\right]\left[(p_e - k)^2 - m_e^2\right]\left[(p-k)^2 - M_{\pi^+}^2\right]}, \tag{43}$$

where $N_\mu(k)$ arises from the Dirac and scalar numerators. Using Feynman parameters,

$$\frac{1}{abc} = 2 \int_0^1 dx \int_0^{1-x} dy \, \frac{1}{\left[ax + by + c(1-x-y)\right]^3}, \tag{44}$$

one combines denominators, shifts the loop momentum, and performs the d-dimensional integral. The scalar part of the resulting expression has the characteristic form

$$I_{\rm scalar} \sim \frac{i}{16\pi^2} \left[\frac{1}{\epsilon_{\rm UV}} - \ln \frac{\Delta}{\mu^2} + \ln \frac{\lambda^2}{\mu^2} + \text{finite terms} \right]. \tag{45}$$

The ultraviolet divergence is absorbed by counterterms, while the $\ln \lambda$ term signals the infrared divergence that must be canceled by real-photon emission.

After adding all virtual one-loop contributions, the differential rate takes the schematic form

$$d\Gamma_{\rm virt} = d\Gamma_0 \left[1 + \frac{\alpha}{\pi} \, g_{\rm virt}(E_e, q^2) + \frac{\alpha}{\pi} \, L_{\rm IR}(\lambda) \right], \tag{46}$$

where

$$L_{\rm IR}(\lambda) \propto \ln \frac{\lambda}{E_{\rm ref}} \tag{47}$$

contains the infrared-divergent logarithm and $g_{\rm virt}$ is a finite kinematic function.

11.2.4 Real-photon emission and bremsstrahlung

The radiative process

$$\pi^+(p) \to \pi^0(p') \, e^+(p_e) \, \nu_e(p_\nu) \, \gamma(k, \varepsilon) \tag{48}$$

must be included because the infrared divergence in Eq. (46) is unphysical and disappears only after combining virtual and real emission. In the soft-photon approximation, the amplitude factorizes:

$$\mathcal{M}^{\rm soft}_{\rm real} = e \, \mathcal{M}_0 \left(\frac{p \cdot \varepsilon^*}{p \cdot k} - \frac{p_e \cdot \varepsilon^*}{p_e \cdot k} \right). \tag{49}$$

Squaring the amplitude and summing over photon polarizations yields

$$|\mathcal{M}^{\rm soft}_{\rm real}|^2 = -e^2 |\mathcal{M}_0|^2 \left[\frac{M_{\pi^+}^2}{(p \cdot k)^2} + \frac{m_e^2}{(p_e \cdot k)^2} - 2 \frac{p \cdot p_e}{(p \cdot k)(p_e \cdot k)} \right]. \tag{50}$$

Integrating over photon momenta with $E_\gamma < \Delta E$, one finds

$$d\Gamma_{\rm soft} = d\Gamma_0 \, \frac{\alpha}{\pi} \left[- L_{\rm IR}(\lambda) + g_{\rm soft}(E_e, \Delta E) \right]. \tag{51}$$

The crucial point is that the infrared logarithm appears with the opposite sign to the one in the virtual correction, so that

$$d\Gamma_{\rm virt} + d\Gamma_{\rm soft} = d\Gamma_0 \left[1 + \frac{\alpha}{\pi}\Big(g_{\rm virt} + g_{\rm soft}\Big)\right] \tag{52}$$

is infrared finite.

Beyond the soft approximation, one must also include hard-photon emission. The full radiative amplitude contains, in addition to inner bremsstrahlung, structure-dependent terms sensitive to the hadronic electromagnetic current. However, for pion beta decay these structure-dependent effects are well controlled and comparatively small. The inclusive rate after integration over the full allowed radiative phase space may therefore be expressed in the form

$$d\Gamma_{\rm incl} = d\Gamma_0 \left[1 + \frac{\alpha}{\pi} g_{\rm EM}(E_e, q^2)\right], \tag{53}$$

where $g_{\rm EM}$ is finite and free of infrared singularities.

11.2.5 Need for chiral perturbation theory

At this stage one might ask whether low-energy QED alone is sufficient. The answer is no, because the pion is not an elementary point particle. Long-distance photons probe hadronic structure, and therefore the low-energy description must incorporate the symmetry structure of QCD. The appropriate framework is chiral perturbation theory (χPT), in which the effective Lagrangian is expanded in powers of momenta, quark masses, and the electric charge:

$$\mathcal{L}_{\rm eff} = \mathcal{L}_{p^2} + \mathcal{L}_{e^2p^0} + \mathcal{L}_{p^4} + \mathcal{L}_{e^2p^2} + \cdots . \tag{54}$$

The leading term $\mathcal{L}_{p^2}$ governs the strong low-energy interactions of the Goldstone bosons, while $\mathcal{L}_{e^2p^0}$ and $\mathcal{L}_{e^2p^2}$ contain the electromagnetic contributions. The weak semileptonic current is introduced through external sources.

In χPT, the pion beta amplitude is expanded around the conserved vector current limit. The loop diagrams generate logarithms and finite terms, while local operators from $\mathcal{L}_{e^2p^2}$ absorb divergences and encode short-distance hadronic information not explicitly resolved within the effective theory. Thus the renormalized form factor may be written as

$$f_+(q^2) = f_+^{(0)}(q^2) + \delta f_+^{\rm loop}(q^2, \mu) + \delta f_+^{\rm ct}(q^2, \mu), \tag{55}$$

with the scale dependence canceling between loop and counterterm pieces. The electromagnetic correction to the decay rate is then not just the point-particle QED correction, but the sum of universal and structure-dependent pieces:

$$\delta_{\rm EM} = \delta_{\rm pt} + \delta_{\rm str}. \tag{56}$$

A useful way to organize this is

$$\mathcal{M} = \frac{G_F}{\sqrt{2}} V_{ud} \left[f_+^{\rm eff}(q^2)\right] (p + p')_\mu \, \bar{u}_\nu \gamma^\mu (1 - \gamma_5) v_e, \tag{57}$$

where

$$f_+^{\rm eff}(q^2) = f_+(q^2) \left[1 + \frac{\alpha}{4\pi} \Gamma_c(v, \lambda)\right] + \delta f_{+,\rm local}^{\rm EM}(q^2). \tag{58}$$

Here $\Gamma_c(v, \lambda)$ represents the universal long-distance electromagnetic function, depending on the usual Dalitz variables, while $\delta f_{+,\rm local}^{\rm EM}$ contains the local structure-dependent electromagnetic terms. This decomposition makes clear that the inclusive decay rate is controlled both by universal bremsstrahlung physics and by the low-energy structure of the pion current.

11.2.6 Differential rate with radiative corrections

The tree-level differential rate for pion beta decay may be written, schematically, as

$$d\Gamma_0 = \frac{1}{2M_{\pi^+}} \frac{G_F^2 |V_{ud}|^2}{2} |f_+(q^2)|^2 L_{\mu\nu} H^{\mu\nu} \, d\Phi_3. \tag{59}$$

Including the radiative corrections, one obtains

$$d\Gamma = d\Gamma_0 \, S_{\rm EW} \left[1 + \frac{\alpha}{\pi} g_{\rm EM}(E_e, q^2) \right] + d\Gamma^{\gamma}_{\rm SD}, \tag{60}$$

where $d\Gamma^{\gamma}_{\rm SD}$ denotes the explicitly structure-dependent hard-photon contribution beyond inner bremsstrahlung. After integration over the allowed kinematic region, this leads to the standard master formula

$$\Gamma_{\pi\beta} = \frac{G_F^2 |V_{ud}|^2 M_{\pi^+}^5}{64\pi^3} |f_+(0)|^2 I_{\pi\beta} \, (1 + \Delta_{\rm RC}), \tag{61}$$

with

$$\Delta_{\rm RC} = \Delta^{\rm EW}_{\rm SD} + \delta^{\rm LD}_{\rm EM} + \delta^{\chi\rm PT}_{\rm str}. \tag{62}$$

The first term is the short-distance Sirlin correction, the second is the universal long-distance QED correction obtained after infrared cancellation, and the third is the residual hadronic structure contribution computed in χPT.

11.2.7 Infrared cancellation in more detail

Since the infrared cancellation is conceptually central, it is useful to make it explicit. Let the virtual correction contain an infrared-singular contribution

$$\delta^{\rm IR}_{\rm virt} = \frac{\alpha}{\pi} A(E_e, q^2) \, \ln \lambda, \tag{63}$$

where λ is the fictitious photon mass. The soft-photon emission contributes

$$\delta^{\rm IR}_{\rm soft} = -\frac{\alpha}{\pi} A(E_e, q^2) \, \ln \lambda, \tag{64}$$

with the same coefficient but opposite sign. Therefore

$$\delta^{\rm IR}_{\rm virt} + \delta^{\rm IR}_{\rm soft} = 0. \tag{65}$$

This cancellation is guaranteed by the Bloch–Nordsieck mechanism and is one reason why the experimentally relevant quantity must be inclusive. The resulting finite correction depends on the experimental definition of the inclusive rate only through the treatment of hard photons, which can be implemented consistently in the Monte Carlo simulation and acceptance calculation.

11.2.8 Interpretation for the extraction of V_{ud}

The importance of the radiative correction can be seen by solving for V_{ud}:

$$|V_{ud}| = \left[\frac{64\pi^3 \, \Gamma_{\pi\beta}}{G_F^2 M_{\pi^+}^5 |f_+(0)|^2 I_{\pi\beta} (1 + \Delta_{\rm RC})} \right]^{1/2}. \tag{66}$$

A relative error in $\Delta_{\rm RC}$ propagates directly into the extracted value of V_{ud}. Since pion beta decay is theoretically attractive precisely because it is free from nuclear-structure corrections, the radiative correction becomes the main theoretical bottleneck. That is why modern analyses devote so much effort to computing $S_{\rm EW}$, the γW box contribution, the long-distance QED functions, and the χPT counterterms with high precision.

The conceptual advantage of pion beta decay is that the hadronic current is a pure vector current between pseudoscalar states, so the underlying structure is exceptionally simple. The same simplicity that makes the decay theoretically clean also makes the radiative correction especially meaningful: one is not correcting a poorly controlled hadronic process, but refining an already clean weak transition. This is precisely why pion beta decay remains an important independent avenue for determining V_{ud} and testing CKM unitarity.

11.2.9 Corrected Decay Rate

Including radiative corrections, the master formula for the pion beta decay width becomes

$$\Gamma_{\pi\beta} = \frac{G_F^2 |V_{ud}|^2 M_\pi^5}{64\pi^3} |f_+(0)|^2 I_{\pi\beta} \, (1 + \Delta_{\mathrm{RC}}), \tag{67}$$

or equivalently,

$$\Gamma_{\pi\beta} = \frac{G_F^2 |V_{ud}|^2 M_\pi^5}{64\pi^3} |f_+(0)|^2 I_{\pi\beta} \, S_{\mathrm{EW}} (1 + \delta_{\mathrm{EM}}). \tag{68}$$

Here $I_{\pi\beta}$ is the phase-space integral and $f_+(0)$ is the vector form factor at zero momentum transfer. This formula shows clearly that the extraction of V_{ud} depends not only on the measured decay rate but also on a precise theoretical understanding of both the universal short-distance correction and the process-dependent long-distance electromagnetic effects.

In summary, radiative corrections are indispensable in pion beta decay. The short-distance part is under good perturbative control, while the long-distance part is computed using chiral perturbation theory. Because pion beta decay is a pure vector transition with very small hadronic uncertainty, the reliability of these radiative corrections is what makes it such a clean and important test of the Standard Model.

11.3 Experimental Realization

One of the primary motivations for studying pion beta decay lies in its direct connection to the unitarity of the Cabibbo–Kobayashi–Maskawa (CKM) matrix. Within the Standard Model, the CKM matrix describes the mixing between quark flavor eigenstates and mass eigenstates, and its unitarity is a fundamental requirement of the theory. In particular, the first-row unitarity condition is given by

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1. \tag{69}$$

Since the element $|V_{ub}|$ is extremely small, this relation is dominated by the contributions from V_{ud} and V_{us}. Any experimentally observed deviation from unity would signal new physics beyond the Standard Model, such as non-standard charged currents or additional fermion generations.

Traditionally, the most precise determinations of V_{ud} have been obtained from superallowed nuclear beta decays. However, such extractions require detailed theoretical corrections accounting for nuclear structure, isospin breaking, and radiative effects. In contrast, pion beta decay, $\pi^+ \to \pi^0 e^+ \nu_e$, is a pure vector transition between pseudoscalar mesons. As a result, it is free from nuclear structure uncertainties and is governed by the conserved vector current (CVC) hypothesis. This makes pion beta decay one of the theoretically cleanest processes for extracting V_{ud}, providing an important independent test of CKM unitarity.

The identification of pion beta decay relies on reconstructing the decay chain

$$\pi^+ \to \pi^0 e^+ \nu_e, \qquad \pi^0 \to \gamma\gamma. \tag{70}$$

Thus, the experimentally accessible final state consists of a positron and two photons, along with missing energy carried by the neutrino. The neutrino escapes detection, and therefore the experimental analysis must rely entirely on the visible particles.

A central observable is the invariant mass of the two photons, defined as

$$M_{\gamma\gamma}^2 = (p_{\gamma_1} + p_{\gamma_2})^2. \tag{71}$$

Signal events are characterized by a peak in this distribution at the neutral pion mass, M_{π^0}. In addition to this invariant-mass reconstruction, the detected positron and photons must be in temporal coincidence and originate from a common spatial vertex corresponding to the pion stopping point. Energy and momentum conservation further constrain the event, allowing one to infer the presence of the undetected neutrino.

Despite its theoretical simplicity, pion beta decay is extremely challenging to measure experimentally. The most significant difficulty arises from its very small branching ratio, of order 10^{-8}. This implies

that only one in approximately 10^8 pion decays produces the desired signal, necessitating extremely high statistics and long data-taking periods.

A second major challenge comes from background processes. The dominant decay mode of the charged pion is $\pi^+ \to \mu^+\nu_\mu$, which occurs nearly 100% of the time. The subsequent decay $\mu^+ \to e^+\nu_e\bar{\nu}_\mu$ produces a large flux of positrons that can mimic the signal if not carefully distinguished. Additionally, the decay $\pi^+ \to e^+\nu_e$, although suppressed, contributes further background.

A third complication is the small Q-value of the pion beta decay, approximately 4 MeV. This limited phase space reduces the energy of the final-state particles and makes it more difficult to separate signal from background using kinematic criteria alone. Consequently, the experiment must rely on a combination of precise timing, energy measurement, and event topology to isolate the signal.

11.4 The PIBETA Experiment

A major advance in the study of pion beta decay was achieved by the PIBETA experiment at the Paul Scherrer Institute (PSI). This experiment was specifically designed to measure rare pion decays with high precision by combining an intense pion beam with a detector optimized for simultaneous detection of positrons and photons.

- The experimental setup begins with a high-intensity beam of positive pions, with a rate on the order of 10^6 particles per second. These pions are slowed down and brought to rest in an active scintillator target. The use of a stopping target is crucial, as it ensures that the decays occur at a well-defined location and simplifies the kinematic reconstruction. The target itself is instrumented to detect the incoming pion and record the time at which it stops, thereby providing a reference for the subsequent decay.

- Surrounding the target is a tracking system composed of cylindrical multi-wire proportional chambers. These detectors measure the trajectories of charged particles emerging from the decay, in particular the positron. By reconstructing the positron track, the experiment can determine its direction, momentum, and point of origin. This information is essential for establishing that the positron originates from the same decay vertex as the photons.

- The most important component of the PIBETA detector is the electromagnetic calorimeter, which consists of a nearly spherical arrangement of pure cesium iodide (CsI) crystals covering almost the full solid angle. This calorimeter is designed to detect photons with high efficiency and excellent energy resolution. When the neutral pion decays into two photons, these photons are absorbed in the calorimeter, producing electromagnetic showers. By measuring the energy and position of these showers, the experiment reconstructs the photon momenta and computes their invariant mass.

11.4.1 Data Acquisition and Event Selection

11.4.2 Event Reconstruction

The reconstruction of pion beta decay events proceeds by combining information from the tracking system and the calorimeter. The photon energies and positions are used to determine their four-momenta, from which the invariant mass is calculated. Events with $M_{\gamma\gamma}$ close to M_{π^0} are identified as candidates for containing a neutral pion.

The positron track is reconstructed independently, and its kinematic properties are measured. Using energy and momentum conservation, the experiment checks whether the observed particles are consistent with a three-body decay involving a missing neutrino. This overconstrained system provides a powerful tool for rejecting background events.

11.4.3 Background Suppression

Given the rarity of the signal, careful suppression of background is essential. One of the most effective methods is the use of timing information. Pion beta decay occurs promptly after the pion stops, whereas



Figure 2: Schematic cross section of the PIBETA detector showing the active target, tracking chambers, veto barrel, and CsI calorimeter.

The data acquisition system is designed to efficiently select events consistent with the pion beta decay signature. A typical trigger requires the detection of a positron and two photon-like energy deposits in the calorimeter within a narrow time window. This ensures that only events with the correct topology are recorded for further analysis. Once the data are collected, a series of selection criteria is applied to isolate the signal. Events are required to have exactly two photon candidates with energies consistent with the decay $\pi^0 \to \gamma\gamma$. The invariant mass of the photon pair must lie within a narrow window around the neutral pion mass. In addition, the positron must be reconstructed with a trajectory pointing back to the pion stopping target, and its timing must be consistent with a prompt pion decay.



Figure 3: π^0 energy spectrum, New results from the PIBETA experiment, D. Pocanic (Virginia U.)

muon decays occur on a much longer timescale. By selecting only events within a narrow time window after the pion stop, the experiment significantly reduces contamination from muon decay.

Energy cuts also play an important role. The energy distribution of positrons from muon decay differs from that of the signal, allowing further discrimination. Finally, the spatial configuration of the detected particles is used to enforce consistency with the expected decay topology. These combined techniques result in a clean sample of pion beta decay events.

11.4.4 Experimental Results

The PIBETA experiment collected an unprecedented dataset of pion decays, corresponding to approximately 6×10^{12} pion stops. From this dataset, a high-purity sample of pion beta decay events was extracted. The measured branching ratio is

$$B_{\pi\beta} = (1.036 \pm 0.006) \times 10^{-8}. \tag{72}$$



Figure 4: Top panel: A typical histogram of time differences between the beam pion stop, t(deg), and 1-arm ht event time, t(e), (dots), compared with a sum of the Monte Carlo simulated responses for π_{e2} decay (π), muon decay (μ), and muon pile-up events (μp). The π_{e2} pile-up background, being much lower, is off scale in the plot. Prompt events are suppressed. Bottom panel: CsI calorimeter energy spectrum for the π_{e2} decay events, after background subtraction.

Using the theoretical expression for the decay rate, this measurement yields

$$|V_{ud}| = 0.9728 \pm 0.0030. \tag{73}$$

Although the uncertainty is larger than that obtained from nuclear beta decay, the result is fully consistent and provides an important independent verification of the Standard Model prediction.

11.5 Next-Generation Experiments: PIONEER

Future experiments aim to improve the precision of pion beta decay measurements by at least an order of magnitude. The PIONEER experiment represents a significant step forward in this direction. It employs a novel detector design based on silicon active targets using low-gain avalanche diode (LGAD) technology, which offers superior spatial and timing resolution compared to traditional scintillator targets.

In addition to improved tracking capabilities, PIONEER incorporates advanced calorimetry optimized for photon detection. These enhancements enable more precise reconstruction of decay events and more effective suppression of background. With an expected dataset of order 10^7 pion beta decay events, the experiment aims to reach a precision at the level of 10^{-4}.

Such a level of precision would allow pion beta decay to become competitive with nuclear beta decay as a probe of V_{ud}. More importantly, it would provide a stringent test of CKM unitarity and could reveal subtle deviations from the Standard Model, potentially pointing toward new physics.

12 Lepton Universality and Precision Tests in Pion Decay

Among low-energy weak processes, pion decay occupies a special place because it is both theoretically simple and experimentally precise. The charged pion is the lightest hadron carrying isospin and electric charge, and its dominant weak decay channels are

$$\pi^+ \to \mu^+\nu_\mu, \qquad \pi^+ \to e^+\nu_e. \tag{74}$$

The muonic channel dominates overwhelmingly, while the electronic one is suppressed by about four orders of magnitude. At first sight this seems surprising. Since the electron is much lighter than the muon, one might naively expect the electronic channel to be favored by phase space. The opposite happens, and the reason is deeply dynamical: the weak interaction is chiral.

The Standard Model predicts that the W-boson couples universally to leptons. In other words, apart from trivial mass effects and radiative corrections, the coupling strengths to e, μ, τ are identical. This statement is called *lepton universality.* Pion decay is an excellent place to test it because in the ratio

$$R_\pi \equiv \frac{\Gamma(\pi^+ \to e^+\nu_e(\gamma))}{\Gamma(\pi^+ \to \mu^+\nu_\mu(\gamma))}, \tag{75}$$

many common factors cancel, including the Fermi constant G_F, the CKM matrix element V_{ud}, and the pion decay constant f_π at leading order. What remains is highly sensitive to the mass dependence generated by helicity suppression and to small radiative corrections that can be computed systematically.

The theoretical cleanliness of this observable is precisely why it has long been considered a sharp probe of the Standard Model. Any deviation of R_π from its predicted value would indicate either a violation of lepton universality or the presence of new pseudoscalar, scalar, or right-handed interactions. To understand this precision test properly, one must first derive the basic pion decay amplitude.

12.1 Effective weak interaction and the pion decay constant

At energies much below the W-boson mass, the charged-current weak interaction is described by the four-fermion effective Hamiltonian

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{ud} \left[\bar{u}\gamma^\mu(1-\gamma_5)d\right] \left[\bar{\nu}_\ell\gamma_\mu(1-\gamma_5)\ell\right] + \text{h.c.} \tag{76}$$

The hadronic part of the matrix element is therefore controlled by the current $\bar{u}\gamma^\mu(1-\gamma_5)d$. Since the pion is a pseudoscalar meson with quantum numbers $J^P = 0^-$, parity implies that the vector current cannot connect the vacuum to a one-pion state, whereas the axial current can. Thus one defines the pion decay constant f_π through

$$\langle 0|\bar{d}\gamma^\mu\gamma_5 u|\pi^+(p)\rangle = if_\pi p^\mu. \tag{77}$$

Equivalently,

$$\langle 0|\bar{u}\gamma^\mu(1-\gamma_5)d|\pi^+(p)\rangle = -if_\pi p^\mu, \tag{78}$$

up to convention-dependent signs that do not affect the rate.

The amplitude for $\pi^+(p) \to \ell^+(p_\ell)\nu_\ell(p_\nu)$ is then

$$i\mathcal{M} = \frac{G_F}{\sqrt{2}} V_{ud} \left(if_\pi p_\mu\right) \left[\bar{u}(p_\nu)\gamma^\mu(1-\gamma_5)v(p_\ell)\right]. \tag{79}$$

Using momentum conservation, $p^\mu = p_\ell^\mu + p_\nu^\mu$, one obtains

$$\mathcal{M} = \frac{G_F}{\sqrt{2}} V_{ud} f_\pi\, \bar{u}(p_\nu)\not{p}(1-\gamma_5)v(p_\ell). \tag{80}$$

Now split $\not{p}$ into the lepton and neutrino momenta:

$$\not{p} = \not{p}_\ell + \not{p}_\nu. \tag{81}$$

Therefore,

$$\mathcal{M} = \frac{G_F}{\sqrt{2}} V_{ud} f_\pi\, \bar{u}(p_\nu) \left(\not{p}_\ell + \not{p}_\nu\right) (1-\gamma_5)v(p_\ell). \tag{82}$$

For the neutrino, which is treated as massless to excellent approximation,

$$\bar{u}(p_\nu)\not{p}_\nu = 0. \tag{83}$$

For the outgoing antilepton spinor,

$$\not{p}_\ell v(p_\ell) = -m_\ell v(p_\ell), \tag{84}$$

so the amplitude becomes

$$\mathcal{M} = -\frac{G_F}{\sqrt{2}} V_{ud} f_\pi\, m_\ell\, \bar{u}(p_\nu)(1-\gamma_5)v(p_\ell). \tag{85}$$

This is the crucial result. The amplitude is proportional to the charged-lepton mass m_ℓ. Hence

$$\mathcal{M} \propto m_\ell, \qquad \Gamma(\pi \to \ell\nu) \propto m_\ell^2. \tag{86}$$

This is the mathematical expression of helicity suppression.

12.2 Why helicity suppression occurs

The mass factor above reflects a conflict between angular momentum conservation and the chiral structure of the weak interaction. The pion has spin zero. In its rest frame, the final-state lepton and neutrino are emitted back-to-back. Because the total angular momentum must remain zero, the two final spins must combine to a spin singlet. In helicity language, this means the two outgoing fermions must have the same helicity when viewed along their own directions of motion.

Now consider the weak interaction. The charged current has the form $V - A$, so it couples only to left-chiral fermions and right-chiral antifermions. For a massless neutrino, chirality and helicity coincide, so the neutrino is left-handed, that is, it has helicity $-1/2$. For the antilepton produced in the decay, the weak interaction selects the corresponding chiral state. In the massless limit, that state has the "wrong" helicity from the point of view of angular momentum conservation in pion decay. As a result, the decay would be impossible if the charged lepton were exactly massless.

The finite lepton mass resolves this conflict. For a massive fermion, chirality and helicity are not identical; a chiral state contains a small admixture of the opposite helicity component. That admixture is proportional to m_ℓ/E_ℓ. Therefore the decay amplitude can proceed only through this helicity-flip component, and that is why the amplitude is proportional to m_ℓ. The electron channel is much more suppressed than the muon channel simply because $m_e \ll m_\mu$.

This makes pion decay a textbook example of a process that is not merely phase-space controlled, but instead dynamically controlled by the chiral structure of the interaction.

12.3 Decay rate and the ratio R_π

Let us now derive the decay rate. For a two-body decay of a spinless particle of mass m_π, the width is

$$\Gamma = \frac{1}{8\pi m_\pi^2}\,|\mathbf{p}_\ell|\,\overline{|\mathcal{M}|^2}, \tag{87}$$

where

$$|\mathbf{p}_\ell| = \frac{m_\pi^2 - m_\ell^2}{2m_\pi} \tag{88}$$

in the limit of vanishing neutrino mass. Starting from

$$\mathcal{M} = -\frac{G_F}{\sqrt{2}} V_{ud} f_\pi\, m_\ell\, \bar{u}(p_\nu)(1-\gamma_5)v(p_\ell), \tag{89}$$

one finds after summing over spins

$$\overline{|\mathcal{M}|^2} = G_F^2 |V_{ud}|^2 f_\pi^2\, m_\ell^2\,(m_\pi^2 - m_\ell^2). \tag{90}$$

Substituting into the phase-space formula gives

$$\Gamma(\pi^+ \to \ell^+\nu_\ell) = \frac{G_F^2 |V_{ud}|^2 f_\pi^2}{8\pi}\, m_\pi\, m_\ell^2 \left(1 - \frac{m_\ell^2}{m_\pi^2}\right)^2. \tag{91}$$

This is the standard tree-level result. Now taking the ratio for $\ell = e$ and $\ell = \mu$:

$$R_\pi^{(0)} = \frac{\Gamma(\pi \to e\nu)}{\Gamma(\pi \to \mu\nu)} = \frac{m_e^2}{m_\mu^2} \frac{\left(1 - \dfrac{m_e^2}{m_\pi^2}\right)^2}{\left(1 - \dfrac{m_\mu^2}{m_\pi^2}\right)^2}. \tag{92}$$

This can be written more compactly as

$$R_\pi^{(0)} = \left(\frac{m_e}{m_\mu}\right)^2 \left(\frac{m_\pi^2 - m_e^2}{m_\pi^2 - m_\mu^2}\right)^2. \tag{93}$$

The cancellation of G_F, V_{ud}, and f_π is now explicit. Thus the ratio is theoretically very clean even before radiative corrections are included.

Numerically, the dominant suppression comes from the factor $(m_e/m_\mu)^2$, which is tiny:

$$\left(\frac{m_e}{m_\mu}\right)^2 \sim 2.3 \times 10^{-5}. \tag{94}$$

The phase-space factor partially compensates this suppression because the electron channel has a larger available phase space, but not nearly enough. Consequently the final ratio is of order 10^{-4}, not order one.

12.4 Lepton universality and what R_π tests

The Standard Model charged current can be written schematically as

$$\mathcal{L}_{cc} = -\frac{g}{\sqrt{2}} \left(\bar{\nu}_e \gamma^\mu P_L e + \bar{\nu}_\mu \gamma^\mu P_L \mu + \bar{\nu}_\tau \gamma^\mu P_L \tau\right) W_\mu^+ + \text{h.c.}, \tag{95}$$

where $P_L = (1 - \gamma_5)/2$. The same gauge coupling g multiplies all three lepton flavors. This is the statement of lepton universality at the level of the Standard Model Lagrangian.

Because pion decay depends on the weak charged current, its branching ratios provide a direct test of whether the electron and muon really couple identically to the W. If one allows, more generally, different effective couplings g_e and g_μ, then the ratio becomes

$$R_\pi = \left(\frac{g_e}{g_\mu}\right)^2 \left(\frac{m_e}{m_\mu}\right)^2 \left(\frac{m_\pi^2 - m_e^2}{m_\pi^2 - m_\mu^2}\right)^2 (1 + \delta R_\pi), \tag{96}$$

where δR_π denotes radiative and structure-dependent corrections. Therefore a precision measurement of R_π directly constrains the ratio g_e/g_μ.

This is why pion decay is so useful. In many hadronic processes, strong-interaction uncertainties obscure the weak-physics interpretation. Here, however, the leading hadronic quantity f_π cancels in the ratio, leaving behind a very sharp test of the Standard Model. Agreement between theory and experiment confirms both helicity suppression and lepton universality. A discrepancy would point immediately toward new physics.

12.5 Chiral perturbation theory and the low-energy description

Although the tree-level decay amplitude depends only on f_π, radiative corrections probe the composite structure of the pion. One cannot treat the pion as an elementary point particle all the way down to arbitrarily low energies and still maintain controlled precision. The correct framework is chiral perturbation theory (χPT), the effective field theory of low-energy QCD based on the spontaneous breaking

$$SU(3)_L \times SU(3)_R \to SU(3)_V. \tag{97}$$

The pseudo-Goldstone bosons are collected in the matrix field

$$U(x) = \exp\left(\frac{i\phi(x)}{F}\right), \tag{98}$$

with F the pion decay constant in the chiral limit. In the presence of external sources and electromagnetism, the lowest-order chiral Lagrangian is

$$\mathcal{L}_2 = \frac{F^2}{4} \operatorname{Tr}\left(D_\mu U D^\mu U^\dagger\right) + \frac{F^2}{4} \operatorname{Tr}\left(\chi U^\dagger + U \chi^\dagger\right), \tag{99}$$

where $\chi = 2B\mathcal{M}$ contains the quark mass matrix and the covariant derivative includes the electromagnetic field and external weak currents. The axial current is obtained by differentiating the generating functional with respect to the appropriate external source, and at leading order one finds

$$A_\mu^a = F\, \partial_\mu \pi^a + \cdots, \tag{100}$$

so that $f_\pi = F$ at leading order. Beyond leading order, loops and higher-order local terms shift this relation, and those same loop effects enter the radiative corrections to pion decay.

When photons and leptons are added, the effective Lagrangian is organized in a simultaneous expansion in small momenta, quark masses, and the electric charge:

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{p^2} + \mathcal{L}_{e^2p^0} + \mathcal{L}_{p^4} + \mathcal{L}_{e^2p^2} + \cdots . \tag{101}$$

The term $\mathcal{L}_{p^2}$ gives the leading strong chiral dynamics, $\mathcal{L}_{e^2p^0}$ gives the leading electromagnetic mass and interaction effects, $\mathcal{L}_{p^4}$ contains the standard Gasser–Leutwyler low-energy constants, and $\mathcal{L}_{e^2p^2}$ contains the electromagnetic counterterms needed to renormalize one-loop radiative corrections. This structure is essential: it is what turns the phrase "hadronic corrections" into a systematic field-theoretic computation.

12.5.1 Structure of radiative corrections

The full decay amplitude may be written schematically as

$$\mathcal{M}_\ell = \mathcal{M}_{0,\ell}\left[1 + \delta_\ell^{\text{virt}} + \delta_\ell^{\text{real}} + \delta_\ell^{\text{SD}}\right], \tag{102}$$

where $\delta_\ell^{\text{virt}}$ denotes virtual-loop corrections, $\delta_\ell^{\text{real}}$ denotes the real-photon bremsstrahlung contribution, and δ_ℓ^{SD} denotes structure-dependent terms. For the ratio R_π, one needs the difference between the electron and muon channels:

$$\delta R_\pi = \delta_e - \delta_\mu. \tag{103}$$

It is useful to separate the correction into short-distance electroweak and long-distance electromagnetic pieces,

$$\delta_\ell = \delta_\ell^{\text{SD,EW}} + \delta_\ell^{\text{LD,EM}}. \tag{104}$$

The short-distance electroweak part arises from momenta between the hadronic scale and M_Z, and is encoded in the matching coefficient of the low-energy four-fermion operator. The long-distance part arises from photons with momenta of order m_π and below, and it is here that χPT is indispensable.

The classic Sirlin viewpoint is that one should first integrate out the heavy electroweak bosons and write a renormalized low-energy weak Hamiltonian. The resulting Wilson coefficient contains universal logarithms of the form

$$\delta^{\text{SD,EW}} \sim \frac{\alpha}{\pi}\ln\frac{M_Z}{\mu}, \tag{105}$$

with μ a hadronic matching scale. These logarithms are process-independent in their universal part and are common to many semileptonic weak processes. What remains is the low-energy photon physics and hadronic structure, which must be treated with an infrared-safe and gauge-invariant low-energy effective theory.

12.5.2 Explicit one-loop analysis in χPT

To compute the long-distance electromagnetic correction, one considers the one-loop diagrams built from the lowest-order chiral and QED interactions. The relevant classes of diagrams are as follows. First, there is the wave-function renormalization of the charged pion from a photon loop attached to the external pion line. Second, there is the wave-function renormalization of the outgoing charged lepton. Third, there is the electromagnetic vertex correction in which a virtual photon is exchanged between the pion and the charged lepton. Fourth, there are local electromagnetic counterterm insertions from $\mathcal{L}_{e^2p^2}$, required to absorb ultraviolet divergences and encode short-distance hadronic information not explicitly resolved in the effective theory. Finally, there is real-photon emission, which must be added to cancel the infrared singularities of the virtual graphs.

Let the tree-level amplitude be denoted $\mathcal{M}_{0,\ell}$. The one-loop corrected amplitude can then be written as

$$\mathcal{M}_\ell^{1-\text{loop}} = \mathcal{M}_{0,\ell}\left[1 + \frac{1}{2}\delta Z_\pi + \frac{1}{2}\delta Z_\ell + \delta V_\ell + \delta C_\ell\right], \tag{106}$$

where δZ_π and δZ_ℓ are the pion and lepton wave-function renormalizations, δV_ℓ is the virtual vertex correction, and δC_ℓ is the local counterterm contribution.

To illustrate the structure explicitly, consider dimensional regularization in $d = 4 - 2\epsilon$ dimensions. The photon-loop correction to an external charged particle generically produces ultraviolet and infrared singular terms of the form

$$\Delta_{\rm UV} = \frac{1}{\epsilon} - \gamma_E + \ln 4\pi + \ln \frac{\mu_\chi^2}{\mu_{\rm UV}^2}, \qquad \Delta_{\rm IR} = \ln \frac{m_\gamma^2}{\mu_\chi^2}, \tag{107}$$

where m_γ is an infrared regulator and μ_χ is the chiral renormalization scale. In a gauge-invariant inclusive calculation, the dependence on m_γ cancels between virtual and real emission.

For the pion wave-function renormalization one finds schematically

$$\delta Z_\pi = -\frac{\alpha}{4\pi}\left[\Delta_{\rm UV} - \Delta_{\rm IR} + F_\pi^{(Z)}\left(\frac{m_\pi^2}{\mu_\chi^2}\right)\right], \tag{108}$$

while for the outgoing charged lepton

$$\delta Z_\ell = -\frac{\alpha}{4\pi}\left[\Delta_{\rm UV} - 2\Delta_{\rm IR} + F_\ell^{(Z)}\left(\frac{m_\ell^2}{\mu_\chi^2}\right)\right]. \tag{109}$$

The specific finite functions $F_\pi^{(Z)}$ and $F_\ell^{(Z)}$ depend on the renormalization scheme, but the key point is that ultraviolet singularities are removed by local counterterms, while infrared singularities survive until the real-photon process is included.

The virtual photon exchange between the charged pion and the lepton gives the vertex integral

$$I_V^\mu = e^2 \int \frac{d^dk}{(2\pi)^d} \frac{(2p-k)^\nu\, \bar{u}_\nu \gamma_\nu (\not{p}_\ell - \not{k} + m_\ell)\gamma^\mu (1-\gamma_5) v_\ell}{\left[k^2 - m_\gamma^2\right]\left[(p-k)^2 - m_\pi^2\right]\left[(p_\ell - k)^2 - m_\ell^2\right]}. \tag{110}$$

After standard Feynman parametrization,

$$\frac{1}{abc} = 2\int_0^1 dx \int_0^{1-x} dy\, \frac{1}{\left[ax + by + c(1-x-y)\right]^3}, \tag{111}$$

one shifts the loop momentum and performs the d-dimensional integral,

$$\int \frac{d^dq}{(2\pi)^d} \frac{1}{(q^2-\Delta)^3} = -\frac{i}{(4\pi)^{d/2}} \frac{\Gamma(3-d/2)}{2\,\Gamma(3)}\, \Delta^{d/2-3}. \tag{112}$$

The result takes the form

$$\delta V_\ell = \frac{\alpha}{4\pi}\left[-\Delta_{\rm UV} + \Delta_{\rm IR}\, \mathcal{C}_{\rm IR} + F_\ell^{(V)}(r_\ell)\right], \qquad r_\ell \equiv \frac{m_\ell^2}{m_\pi^2}. \tag{113}$$

The finite function $F_\ell^{(V)}(r_\ell)$ contains logarithms and dilogarithms arising from the two-scale kinematics. In practice, one-loop calculations are organized so that the sum

$$\frac{1}{2}\delta Z_\pi + \frac{1}{2}\delta Z_\ell + \delta V_\ell + \delta C_\ell \tag{114}$$

is ultraviolet finite, but still infrared divergent. The remaining infrared singularity is cancelled by soft bremsstrahlung.

The local counterterms in $\mathcal{L}_{e^2p^2}$ are crucial. They represent the most general chirally and gauge-invariant local operators involving the pion, photon, weak spurions, and lepton fields at that order. Denoting the relevant renormalized low-energy constants collectively by K_i^r and X_i^r, one may write the local contribution in the decay amplitude as

$$\delta C_\ell = \frac{\alpha}{4\pi}\left[\sum_i c_i K_i^r(\mu_\chi) + \sum_j d_j X_j^r(\mu_\chi)\right]. \tag{115}$$

The coefficients c_i and d_j are fixed by the operator basis and the kinematics of the process. Their scale dependence cancels the μ_χ-dependence of the loop functions, so the final physical amplitude is scale independent.

12.5.3 Real-photon emission and infrared cancellation

The radiative process

$$\pi^+(p) \to \ell^+(p_\ell)\,\nu_\ell(p_\nu)\,\gamma(k,\varepsilon) \tag{116}$$

contains two conceptually distinct contributions. The first is inner bremsstrahlung, in which the photon is emitted from the external charged pion or lepton legs. This part is fixed by QED gauge invariance and by the non-radiative amplitude. The second is structure-dependent radiation, which probes the internal hadronic structure of the pion and is parameterized in terms of radiative form factors.

For the purpose of the inclusive width, the infrared-divergent part comes entirely from inner bremsstrahlung. In the soft-photon limit $k \to 0$, the amplitude factorizes as

$$\mathcal{M}_\ell^{\rm IB} = e\,\mathcal{M}_{0,\ell}\left(\frac{p\cdot\varepsilon^*}{p\cdot k} - \frac{p_\ell\cdot\varepsilon^*}{p_\ell\cdot k}\right). \tag{117}$$

Squaring and integrating over photon phase space below a soft cutoff $E_\gamma < \Delta E$ gives

$$\delta_\ell^{\rm soft} = \frac{\alpha}{\pi}\left[B_\ell(r_\ell)\,\ln\frac{2\Delta E}{m_\gamma} + F_\ell^{\rm soft}(r_\ell,\Delta E)\right]. \tag{118}$$

The coefficient $B_\ell(r_\ell)$ is exactly such that the $\ln m_\gamma$ dependence cancels the infrared logarithm present in the virtual correction. Therefore

$$\delta_\ell^{\rm virt} + \delta_\ell^{\rm soft} \tag{119}$$

is infrared finite. Adding the hard-photon region and the structure-dependent radiative contribution gives the fully inclusive radiative correction.

A particularly important point for R_π is that the correction depends nontrivially on the final-state lepton mass. This is why the electron and muon channels do not receive identical QED corrections, and why δR_π is nonzero even though many hadronic factors cancel. The lepton-mass dependence enters both through the kinematics and through the collinear logarithms that enhance the electron channel relative to the muon channel.

12.5.4 Sirlin-type representation of $\delta_{\rm rad}$

The corrected width may be represented in a Sirlin-like factorized form,

$$\Gamma(\pi\to\ell\nu(\gamma)) = \Gamma_0(\pi\to\ell\nu)\,S_{\rm EW}\left[1+\frac{\alpha}{\pi}F(r_\ell)\right]\left[1+\delta_\ell^{\rm str}\right], \tag{120}$$

where $S_{\rm EW}$ is the short-distance electroweak factor, $F(r_\ell)$ is the universal point-like QED function, and $\delta_\ell^{\rm str}$ is the structure-dependent hadronic correction calculated in χPT. Forming the ratio, one obtains

$$R_\pi = R_\pi^{(0)}\left\{1+\frac{\alpha}{\pi}\big[F(r_e)-F(r_\mu)\big] + \big(\delta_e^{\rm str}-\delta_\mu^{\rm str}\big)\right\}. \tag{121}$$

Thus

$$\delta R_\pi = \frac{\alpha}{\pi}\big[F(r_e)-F(r_\mu)\big] + \big(\delta_e^{\rm str}-\delta_\mu^{\rm str}\big). \tag{122}$$

At one loop, the point-like function contains terms of the schematic form

$$F(r_\ell) = \frac{3}{2}\ln\frac{m_\pi^2}{m_\ell^2} + \frac{13-19r_\ell}{8(1-r_\ell)} - \frac{8-5r_\ell}{4(1-r_\ell)^2}r_\ell\ln r_\ell + \Phi(r_\ell), \tag{123}$$

where $\Phi(r_\ell)$ contains finite terms built from logarithms and dilogarithms. The exact analytic expression depends on the convention used for separating inner bremsstrahlung from structure-dependent pieces and on the treatment of the inclusive phase space, but the logarithmic dependence on m_ℓ is universal and is the main source of the difference between the electron and muon channels.

The structure-dependent correction in χPT may be written schematically as

$$\delta_\ell^{\rm str} = \frac{\alpha}{\pi}\left[A_\chi\ln\frac{m_\rho}{\mu_\chi} + B_\chi(r_\ell) + 4\pi^2\,\mathcal{K}_\ell^r(\mu_\chi)\right], \tag{124}$$

where A_χ is fixed by the chiral loop divergence structure, $B_\chi(r_\ell)$ is the finite loop function, and $\mathcal{K}_\ell^r$ denotes an appropriate combination of renormalized electromagnetic low-energy constants. The resonance scale m_ρ often appears in practical matching estimates of the low-energy constants, since vector-meson saturation provides a useful phenomenological guide.

Combining the short-distance logarithm, the universal one-loop QED correction, and the χPT structure-dependent part, one arrives at the final correction

$$\delta_{\rm rad} \equiv \delta R_\pi = \delta_{\rm EW}^{\rm short} + \delta_{\rm QED}^{\rm point} + \delta_{\chi{\rm PT}}^{\rm structure}. \tag{125}$$

The practical significance of this formula is that each term has a definite field-theoretic meaning. The first comes from electroweak matching and renormalization-group running, the second from the universal long-distance QED dressing of a point-like charged pion decay, and the third from genuinely hadronic electromagnetic structure effects. This is exactly the sense in which the modern calculation is "Sirlin-type" but completed at low energies by χPT.

12.6 Prediction for R_π

Combining the phase-space factors and helicity suppression, one obtains

$$R_\pi = \left(\frac{m_e}{m_\mu}\right)^2 \left(\frac{M_\pi^2 - m_e^2}{M_\pi^2 - m_\mu^2}\right)^2 (1 + \delta_{\rm rad}). \tag{126}$$

A crucial feature of this ratio is that the decay constant f_π cancels exactly, eliminating hadronic uncertainties. This makes R_π one of the cleanest observables in particle physics.

The Standard Model prediction is known to very high precision,

$$R_\pi^{\rm SM} = (1.2352 \pm 0.0001) \times 10^{-4}. \tag{127}$$

12.7 The PEN Experiment

12.7.1 Overview and Physics Goal

The PEN experiment at the Paul Scherrer Institute (PSI) was designed to measure the branching ratio $R_\pi = \Gamma(\pi^+ \to e^+\nu_e)/\Gamma(\pi^+ \to \mu^+\nu_\mu)$ with high precision, thereby providing a stringent test of lepton universality. The experiment builds upon earlier measurements, particularly the PIBETA experiment, but introduces significant improvements in signal discrimination through waveform digitization techniques.

The primary experimental challenge is the suppression of the dominant $\pi^+ \to \mu^+ \to e^+$ decay chain, which exceeds the signal by several orders of magnitude. The PEN experiment addresses this challenge by exploiting differences in both energy and time structure between the signal and background processes.

12.7.2 Beam and Target System

A continuous high-intensity beam of positive pions, with a rate of approximately 10^6 π^+ per second, is delivered to the experimental setup. These pions are slowed and brought to rest in an active plastic scintillator target. The stopping of the pion produces a prompt scintillation signal, which defines the initial time reference for the event.

The use of a stopping target is essential for simplifying the kinematics of the decay. Since the pion decays at rest, the energy of the emitted positron in $\pi \to e\nu$ decay is fixed, providing a clean experimental signature. Furthermore, the well-defined spatial location of the decay allows for precise reconstruction of the event vertex.

12.7.3 Detector Design

The detector system surrounding the target is designed to measure both the energy and trajectory of the decay products. Charged particles are tracked using cylindrical multi-wire proportional chambers (MWPCs), which provide information about the direction and origin of the positron. This ensures that selected events originate from the pion stopping region, reducing contamination from background

processes. The energy of the positron is measured using a large-acceptance electromagnetic calorimeter composed of pure CsI crystals. The calorimeter provides high-resolution measurements of energy and timing, enabling the distinction between the monoenergetic $\pi \to e\nu$ signal and the continuous Michel spectrum from muon decay. However, finite resolution and shower leakage introduce a low-energy tail in the signal distribution, which must be carefully modeled.



Figure 5: Schematic cross section of the PEN apparatus, shown in the 2009-10 running configuration

12.7.4 Waveform Digitization and Signal Formation

A defining feature of the PEN experiment is the use of high-speed waveform digitization of signals from the active target. The detector electronics sample the signal at gigahertz frequencies, recording the full time evolution of the waveform rather than a single integrated value.

Each physical interaction within the detector produces a distinct pulse in the waveform. In the case of direct pion decay, the waveform consists of two pulses: one corresponding to the pion stopping in the target and a second corresponding to the positron emission. In contrast, the decay chain $\pi^+ \to \mu^+ \to e^+$ produces three pulses, with the intermediate pulse arising from the muon.

The time separation between these pulses reflects the difference between the pion lifetime (~ 26 ns) and the muon lifetime (~ 2.2 μs). By fitting the waveform with template pulse shapes, the experiment can determine the number and timing of pulses, providing a powerful method for distinguishing signal from background.

12.7.5 Data Acquisition and Triggering

The data acquisition system is designed to trigger on events consistent with pion decay. A typical trigger condition requires a valid pion stop signal in the target followed by a positron-like signal in the calorimeter within a predefined time window.

Once triggered, the full waveform from the target, along with signals from the calorimeter and tracking detectors, is recorded for offline analysis. This approach ensures that all relevant information is preserved, allowing for flexible reconstruction and event classification.

12.7.6 Event Reconstruction and Selection

In the offline analysis, the recorded waveforms are processed to extract the timing and amplitude of individual pulses. This is achieved by fitting the waveform to a sum of template pulse shapes, each representing a distinct physical process.

Events are classified based on the number of pulses identified in the waveform. Two-pulse events are identified as candidates for $\pi \to e\nu$ decay, while three-pulse events are attributed to the $\pi \to \mu \to e$ decay chain. Additional selection criteria are applied based on the positron energy and track reconstruction to further suppress background.



Figure 6: Data points: decay time spectra for the clean samples of $\pi \to e\nu$ (red) and $\pi \to \mu \to e$ (blue) events from the 2010 PEN data set

The extraction of R_π is performed using a maximum likelihood fit to the combined energy, timing, and waveform distributions. Signal and background templates are obtained from detailed Monte Carlo simulations, which incorporate detector effects and radiative corrections.

12.7.7 Systematic Uncertainties

The precision of the PEN measurement is limited not only by statistical uncertainties but also by systematic effects. The dominant sources of systematic uncertainty include the calibration of the calorimeter energy scale, the modeling of the detector response function, and the efficiency of waveform-based event classification.

Additional uncertainties arise from radiative processes such as $\pi \to e\nu\gamma$, which modify the observed energy spectrum. These effects are incorporated into the simulation and their impact is evaluated as part of the systematic error analysis.

Through careful calibration, detailed simulation, and robust analysis techniques, the PEN experiment achieves a high level of control over systematic uncertainties, enabling a precise determination of R_π.

12.8 The PIENU Experiment

The PIENU experiment at TRIUMF adopts a complementary strategy focused on achieving superior energy resolution. The central component of the detector is a large NaI(Tl) calorimeter, which provides excellent resolution for electromagnetic showers due to its high light yield and uniform response.

In PIENU, pions are stopped in a thin target, and the emitted positrons are detected in the calorimeter. The high resolution of the detector results in a sharply defined peak for the $\pi \to e\nu$ decay, minimizing overlap with the Michel spectrum. Surrounding detectors capture escaping energy and improve the accuracy of energy reconstruction, reducing the impact of shower leakage.

Timing information is used in conjunction with energy measurements to separate prompt and delayed events. By applying cuts in both energy and time, the experiment isolates a clean sample of signal events. The analysis involves fitting the observed energy spectrum with contributions from the signal peak and the background distribution, taking into account the detector response function.

A key strength of the PIENU experiment is its precise control of systematic uncertainties related to energy measurement. The detector is calibrated using known sources, and its stability is continuously monitored. This allows for an accurate determination of the energy scale and resolution, which are essential for extracting the signal yield.

12.9 Results and Implications

The results from the PEN and PIENU experiments yield a value of

$$R_\pi^{\text{exp}} \approx 1.234 \times 10^{-4}, \tag{128}$$

which is in excellent agreement with the Standard Model prediction. This agreement confirms lepton universality at the level of 10^{-3} and places strong constraints on new physics scenarios.

In particular, models involving scalar or tensor interactions could modify the decay amplitude and lead to deviations from the predicted value of R_π. The absence of such deviations therefore constrains the parameter space of these models and reinforces the validity of the Standard Model description.

13 Conclusion

Pion decays provide a theoretically clean and experimentally precise framework for testing the Standard Model at low energies. Pion beta decay enables a direct determination of the CKM matrix element V_{ud}, while leptonic decays test lepton universality through the ratio R_π. The simplicity of the pion system allows for controlled theoretical predictions, with radiative corrections systematically incorporated using electroweak theory and chiral perturbation theory.

On the experimental side, measurements from PIBETA, PEN, and PIENU have achieved high precision through advanced techniques in timing, calorimetry, and waveform analysis, with future experiments such as PIONEER aiming to further improve sensitivity. The overall agreement between theory and experiment places strong constraints on possible physics beyond the Standard Model.

14 Question from the Presentation

In high-intensity pion decay experiments such as PIBETA and PEN, a very large number of π^+ particles are delivered to the detector to achieve high statistics. Does this lead to pile-up of multiple pion stops and decay events within short time intervals, and if so, how does this affect the measurement of branching ratios?

Answer

Pile-up is indeed a relevant concern in both PIBETA and PEN due to the high beam intensities. The probability of overlapping events is roughly given by

$$P_{\text{pile-up}} \sim \Phi_\pi \, \Delta t, \tag{129}$$

where Φ_π is the pion beam rate and Δt is the detector time resolution or analysis window. Since pion and muon decays occur on nanosecond to microsecond timescales, overlapping signals can occur if the beam intensity is sufficiently high.

However, both experiments are specifically designed to control and suppress pile-up effects. In PIBETA, strong kinematic and timing constraints are imposed. Events must reconstruct a consistent decay topology, including a positron and two photons satisfying

$$M_{\gamma\gamma} \approx m_{\pi^0}, \tag{130}$$

and must originate from the same decay vertex within a narrow time window. Random overlaps from different pion decays typically fail these conditions and are rejected.

In PEN, pile-up rejection is even more powerful due to waveform digitization in the active target. The detector records the full time structure of the signal, allowing identification of multiple overlapping pulses. Since genuine $\pi \to e\nu$ events produce a two-pulse structure, while background and pile-up events produce additional pulses, fitting the waveform enables event-by-event discrimination.

Any residual pile-up that survives selection cuts is treated as a background component and modeled using Monte Carlo simulations. Its contribution is included in the statistical analysis and does not bias the final result.

Thus, although high beam intensities increase the likelihood of pile-up, its effects are well understood and controlled, ensuring reliable extraction of the branching ratios.

15 References

A. Sirlin, "Current Algebra Formulation of Radiative Corrections in Gauge Theories and the Universality of the Weak Interactions," *Rev. Mod. Phys.* **50**, 573 (1978).

W. J. Marciano and A. Sirlin, "Improved Calculation of Electroweak Radiative Corrections and the Value of V_{ud}," *Phys. Rev. Lett.* **96**, 032002 (2006).

J. Gasser and H. Leutwyler, "Chiral Perturbation Theory to One Loop," *Annals Phys.* **158**, 142 (1984).

J. Gasser and H. Leutwyler, "Chiral Perturbation Theory: Expansions in the Mass of the Strange Quark," *Nucl. Phys. B* **250**, 465 (1985).

D. Počanić *et al.* (PIBETA Collaboration), "Precise Measurement of the $\pi^+ \to \pi^0 e^+ \nu$ Branching Ratio," *Phys. Rev. Lett.* **93**, 181803 (2004).

D. Počanić *et al.*, "Pion Beta Decay: Theory, Experiment, and Applications," *Phys. Rev. C* **79**, 055502 (2009).

Particle Data Group, "Review of Particle Physics," *Prog. Theor. Exp. Phys.* **2024**, 083C01 (2024).

A. Aguilar-Arevalo *et al.* (PEN Collaboration), "Improved Measurement of the $\pi \to e\nu$ Branching Ratio," *Phys. Rev. Lett.* **115**, 071801 (2015).

A. Aguilar-Arevalo *et al.* (PIENU Collaboration), "Improved Measurement of the $\pi^+ \to e^+\nu$ Branching Ratio," *Phys. Rev. Lett.* **115**, 071801 (2015).

PIONEER Collaboration, "PIONEER: A Next-Generation Experiment for Rare Pion Decays," available at `https://inspirehep.net/experiments/1511062`.

N. Cabibbo, "Unitary Symmetry and Leptonic Decays," *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, "CP Violation in the Renormalizable Theory of Weak Interaction," *Prog. Theor. Phys.* **49**, 652 (1973).

M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Addison-Wesley (1995).