Pseudo-Rational Choice with Multiple Tastes and Beliefs

Gil Riella* Leonardo Sampaio†

November 6, 2026

Abstract

This paper aims to provide an axiomatic characterization of pseudo-rational multi-taste, multi-prior choice over menus of Anscombe–Aumann acts. Taking a choice correspondence as primitive, the framework allows for violations of the Weak Axiom of Revealed Preference. The main objective is to characterize choice correspondences admitting a representation of the form

$$c(A) = \bigcup_{(\pi,u)\in\mathcal{M}} \arg\max_{f\in A} \sum_{s\in S} \pi(s)\mathbb{E}_{f(s)}[u],$$

where $\mathcal{M}$ is a set of prior–utility pairs, with each prior representing beliefs over states and each utility index representing a possible evaluation of prizes.

The model follows the notion of justifiable choice: an act is chosen whenever it maximizes subjective expected utility under at least one admissible belief–taste pair. This allows for uncertainty regarding both the states of the world and the decision maker's tastes.

The analysis focuses initially on finite prize spaces and explores an extension to compact metric prize spaces. The proposed approach draws on maximal-menu arguments and Heller's (2012) minimax theorem method to investigate the connection between revealed choice and multi-utility representations. Finally, the paper seeks to explore how indecisiveness and indifference may be identified and distinguished from observed choice behavior.

Keywords: Menu effects, Multiple priors, Multiple utilities, Incomplete preferences, Revealed preference, Indecisiveness.

JEL Codes: D81, D83

*Department of Economics, University of Brasília
†Department of Economics, University of Brasília

1 Introduction

The literature studying indecisiveness in beliefs and tastes using a binary relation as the primitive is recent but includes several notable contributions (Riella, 2014; Hara and Riella, 2015; Galaabaatar and Karni, 2013). By contrast, comparatively little work addresses these questions using a choice correspondence.

Eliaz and Ok (2006) provide a pioneering contribution by introducing the WARNI axiom and characterizing when observed behavior can distinguish indecisiveness from indifference. More recently, Heller (2012) and Safonov (2023) study this distinction in environments involving risk and uncertainty, offering insights into indecisiveness in either tastes or beliefs. Their models implicitly treat the decision maker's behavior as pseudo-rational choice in the sense of Aizerman and Malishevski (1981).

To the best of our knowledge, however, no model characterizes conditions on a choice correspondence under which a decision maker behaves as if she were simultaneously indecisive about both her tastes and her beliefs, rather than about either dimension separately.

2 Model

We consider an Anscombe–Aumann setup with a finite state space S and a compact metric prize space X. An act is a function that maps each state of the world $s \in S$ to a lottery $f(s) \in \Delta(X)$, and we denote the set of all acts by $\mathcal{F}$.

Here, $\Delta(X)$ denotes the set of Borel probability measures on X, endowed with any metric that metrizes the topology of weak convergence, and $\mathcal{F} := \Delta(X)^S$ is endowed with the product metric. Thus, $f_n \to f$ in $\mathcal{F}$ if and only if $f_n(s)$ converges weakly to $f(s)$ for every $s \in S$. Since X is compact and metrizable and S is finite, both $\Delta(X)$ and $\mathcal{F}$ are compact metrizable spaces. Whenever continuous utility indices are considered, $C(Y)$ denotes the space of continuous real-valued functions on a compact metric space Y, endowed with the supremum norm. In particular, state-dependent utility indices belong to $C(X \times S)$, while state-independent utility indices belong to $C(X)$.

Let $\mathcal{A}$ be the collection of nonempty compact menus of acts, endowed with the Hausdorff metric induced by the metric on $\mathcal{F}$, so that $\mathcal{A} \subseteq 2^{\mathcal{F}} \setminus \{\emptyset\}$. Our primitive is a choice correspondence $c\colon \mathcal{A} \to 2^{\mathcal{F}} \setminus \{\emptyset\}$ such that, for every $A \in \mathcal{A}$, $\emptyset \neq c(A) \subseteq A$. The choice correspondence is an object that the analyst can usually observe and is therefore more primitive than the notion of preference (Ok and Eliaz, 2006). Its interpretation is as follows: when the decision maker is presented with a menu A, she may choose any act in $c(A)$.

We define mixtures of acts state by state. That is, for every $\alpha \in [0,1]$, every $g, h \in \mathcal{F}$, and every $s \in S$, $\big(\alpha g + (1-\alpha)h\big)(s) = \alpha g(s) + (1-\alpha)h(s)$. Hence, $\mathcal{F}$ can also be viewed as a mixture space. We abuse notation by using $x \in X$ to denote the degenerate constant act that yields outcome x with certainty in every state, and $p \in \Delta(X)$ to denote the constant act that yields lottery p in every state $s \in S$. For $\nu \in C(X \times S)$ and $f \in \mathcal{F}$, the associated state-dependent expected utility is

$$U_\nu(f) := \sum_{s \in S} \int_X \nu(x,s)\, df(s)(x).$$

For a prior $\pi \in \Delta(S)$ and a state-independent utility index $u \in C(X)$, define the subjective expected utility of $f \in \mathcal{F}$ by

$$V_{\pi,u}(f) := \sum_{s \in S} \pi(s) \int_X u(x)\, df(s)(x).$$

Here, $\Delta(S)$ is the simplex of probability distributions on S, endowed with its Euclidean topology. The space $\Delta(S) \times C(X)$ carries the product of this topology and the supremum-norm topology. Since u is continuous and X is compact, the integrals are well defined and $V_{\pi,u}$ is continuous and affine on $\mathcal{F}$. For a constant act $p \in \Delta(X)$, $V_{\pi,u}(p) = \int_X u(x)\, dp(x)$. When X is finite, both U_ν and $V_{\pi,u}$ reduce to the corresponding finite sums.

2.1 Axioms

We impose the following axioms on the choice correspondence c.

Axiom 1 (Pseudo-WARP). For all $A, B \in \mathcal{A}$, if $f \in c(B) \cap A$ and $c(A) \subseteq B$, then $f \in c(A)$.

Axiom 2 (Independence). For every $A \in \mathcal{A}$, $\lambda \in (0,1)$, and $h \in \mathcal{F}$,[1] $c\big(\lambda A + (1-\lambda)h\big) = \lambda c(A) + (1-\lambda)h$.

Axiom 3 (Continuity). The set $\big\{(A, f) \in \mathcal{A} \times \mathcal{F} : f \in c(A)\big\}$ is closed.

Axiom 4 (Indirect Ambiguity Aversion). For every $A \in \mathcal{A}$, $\lambda \in [0,1]$, and $f, g, h \in \mathcal{F}$, if $f \notin c\big(A \cup \{h\}\big)$ and $f \notin c\big(A \cup \{g\}\big)$, then $f \notin c\big(A \cup \{\lambda g + (1-\lambda)h\}\big)$.

Axiom 5 (No-Hedging). For every $A \in \mathcal{A}$, $\lambda \in [0,1]$, and $g, h \in \mathcal{F}$, if $\{g, h\} \subseteq A$, then $c(A) \subseteq c\big(A \cup \{\lambda g + (1-\lambda)h\}\big)$.

Axiom 6 (Best and Worst). There exist distinct constant acts $p^*, p_* \in \Delta(X)$ such that, for every $A \in \mathcal{A}$ and every $\lambda \in (0,1)$,

$$c\left(\{p^*\} \cup \big(\lambda p_* + (1-\lambda)A\big)\right) = \{p^*\}, \tag{BW1}$$

and

$$p_* \notin c\left(\{p_*\} \cup \big(\lambda p^* + (1-\lambda)A\big)\right). \tag{BW2}$$

Axiom 7 (One-Side Monotonicity). For every $A \in \mathcal{A}$ and every $f, g \in \mathcal{F}$, if $f \notin c\big(A \cup \{g(s)\}\big)$ for every $s \in S$, then $f \notin c\big(A \cup \{g\}\big)$.

Pseudo-WARP imposes consistency across menus while allowing the choice correspondence to be generated by multiple possible rationales. Independence requires choice to commute with a common mixture transformation, and Continuity rules out discontinuous changes in choice as menus and acts converge. Indirect Ambiguity Aversion imposes a form of consistency under hedging: if f is not chosen in the presence of h, and h is not chosen in the presence of g, then replacing h with any mixture of g and h cannot make f chosen.

[1] We abuse notation and write $\lambda A + (1-\lambda)h$ for the set $\{\lambda f + (1-\lambda)h : f \in A\}$.

No-Hedging requires that adding a mixture of two acts already available in a menu does not eliminate any act that was previously chosen. Best and Worst imposes two complementary requirements: (BW1) makes p^* uniquely chosen when it is added alongside mixtures of each act in the menu with p_*, while (BW2) rules out p_* from choice when it is added alongside mixtures of each act in the menu with p^*. Finally, One-Side Monotonicity requires that if f is not chosen when each statewise lottery of g is added separately to a menu as a constant act, then f is not chosen when g itself is added to that menu.

3 Representation

3.1 Additively Separable Multi-Utility Representation

For this subsection, let X be a compact metric prize space. Consider a strict relation $\succ \subseteq \mathcal{F} \times \mathcal{F}$ and the following axioms:

(S1) Independence. For all $f, g, h \in \mathcal{F}$ and $\alpha \in (0, 1]$, $f \succ g \iff \alpha f + (1-\alpha)h \succ \alpha g + (1-\alpha)h$

(S2) Open Continuity. For every $f \in \mathcal{F}$, the sets $\{g \in \mathcal{F} : f \succ g\}$ and $\{g \in \mathcal{F} : g \succ f\}$ are open in $\mathcal{F}$.

(S3) Constant Non-Triviality. There exist $\hat{p}, \hat{q} \in \Delta(X)$ such that $\hat{p} \succ \hat{q}$, where lotteries are identified with the corresponding constant acts.

Proposition 1. A strict partial order $\succ \subseteq \mathcal{F} \times \mathcal{F}$ satisfies (S1)–(S3) if and only if there exists a nonempty compact and convex set $\mathcal{V} \subset C(X \times S)$ and there exist $\hat{p}, \hat{q} \in \Delta(X)$ such that $U_\nu(\hat{p}) > U_\nu(\hat{q})$ for every $\nu \in \mathcal{V}$, and, for every $f, g \in \mathcal{F}$,

$$f \succ g \iff U_\nu(f) > U_\nu(g) \quad \text{for every } \nu \in \mathcal{V}.$$

The proof builds on the representation results of Hara and Riella (2023), particularly Theorems 1 and 8. Following the construction in the proof of their Theorem 8, we associate the primitive strict partial order with an auxiliary state-dependent preorder satisfying the conditions of their Theorem 1. This yields a compact family of continuous utilities, from which the original strict relation is recovered through strict inequalities. Finally, taking the closed convex hull of the representing family preserves both compactness and the strict representation, yielding the desired compact convex set of utilities.

Building on Proposition 1, we can follow Heller's (2012) approach and apply a minimax theorem to obtain a justifiable-choice representation. The argument begins by constructing an auxiliary relation from the maximal menus associated with each act in the Anscombe–Aumann framework.

Theorem 1 *The following statements are equivalent:*

(1) The choice correspondence c satisfies Axioms 1–6.

(2) There exist distinct lotteries $p^*, p_* \in \Delta(X)$ and a nonempty compact set $\mathcal{W} \subseteq C(X \times S)$ such that, for every $A \in \mathcal{A}$,

$$c(A) = \bigcup_{\nu \in \mathcal{W}} \arg\max_{g \in A} U_\nu(g),$$

and, for every $\nu \in \mathcal{W}$,

$$U_\nu(p^*) = 1, \qquad U_\nu(p_*) = 0, \qquad 0 \leq U_\nu(g) \leq 1 \quad \textit{for every } g \in \mathcal{F}.$$

3.2 Multi-Prior Multi-Utility Representation

Theorem 2 *The following statements are equivalent:*

(1) The choice correspondence c satisfies Axioms 1–7.

(2) There exists a nonempty closed subset $\mathcal{M} \subseteq \Delta(S) \times C(X)$ of probability–utility pairs such that, for every menu $A \in \mathcal{A}$,

$$c(A) = \bigcup_{(\pi,u) \in \mathcal{M}} \arg\max_{g \in A} V_{\pi,u}(g).$$

Moreover, there exist $\hat{p}, \hat{q} \in \Delta(X)$ such that

$$\mathbb{E}_{\hat{p}}(u) > \mathbb{E}_{\hat{q}}(u) \quad \textit{for every } (\pi, u) \in \mathcal{M}.$$

4 Indecisiveness and Indifference

Write later

5 Applications

Write later

6 Appendix

6.1 Proof of Proposition 1

Necessity. The necessity of the axioms is standard. Suppose that $\succ$ admits the stated representation. Open Continuity follows from the compactness of $\mathcal{V}$ and the continuity of U_ν. Independence follows from the affinity of each U_ν, and Constant Non-Triviality follows from the existence of $\hat{p}, \hat{q}$ such that $U_\nu(\hat{p}) > U_\nu(\hat{q})$ for every $\nu \in \mathcal{V}$. ■

Sufficiency. Suppose that $\succ$ satisfies (S1)–(S3). By Constant Non-Triviality, there exist $\hat{p}, \hat{q} \in \Delta(X)$ such that $\hat{p} \succ \hat{q}$. Define the binary relation $\succsim^*$ on $\mathcal{F}$ by

$$f \succsim^* g \iff \alpha\hat{p} + (1-\alpha)f \succ \alpha\hat{q} + (1-\alpha)g \quad \text{for every } \alpha \in (0,1).$$

By the argument in the proof of Theorem 8 of Hara and Riella (2023), $\succsim^*$ is a preorder satisfying Independence, Continuity, and Directional Open-Continuity with respect to the pair $(\hat{p}, \hat{q})$. Moreover, $\succ \subseteq \succsim^*$.

We may therefore apply Theorem 1 of Hara and Riella (2023). In particular, its proof yields a compact set $\mathcal{V}_0 \subset C(X \times S)$ such that

$$f \succsim^* g \iff U_\nu(f) \geq U_\nu(g) \quad \text{for every } \nu \in \mathcal{V}_0,$$

and $U_\nu(\hat{p}) - U_\nu(\hat{q}) = 1$ for every $\nu \in \mathcal{V}_0$.

We first show that $f \succ g$ implies $U_\nu(f) > U_\nu(g)$ for every $\nu \in \mathcal{V}_0$.

Suppose that $f \succ g$. By Independence, $f \succ \frac{1}{2}f + \frac{1}{2}g \succ g$. Let $h = \frac{1}{2}f + \frac{1}{2}g$. By Open Continuity, there exist open neighborhoods O_f of f and O_g of g such that $f' \succ h \succ g'$ for all $(f', g') \in O_f \times O_g$. By transitivity, $f' \succ g'$ for all $(f', g') \in O_f \times O_g$.

Since the mappings $\alpha \mapsto \alpha\hat{q} + (1-\alpha)f$ and $\alpha \mapsto \alpha\hat{p} + (1-\alpha)g$ are continuous and converge to f and g, respectively, as $\alpha \to 0$, there exists $\alpha \in (0,1)$ sufficiently small such that $\alpha\hat{q} + (1-\alpha)f \in O_f$ and $\alpha\hat{p} + (1-\alpha)g \in O_g$. Consequently, $\alpha\hat{q} + (1-\alpha)f \succ \alpha\hat{p} + (1-\alpha)g$.

Since $\succ \subseteq \succsim^*$, the representation of $\succsim^*$ implies that, for every $\nu \in \mathcal{V}_0$,

$$\alpha U_\nu(\hat{q}) + (1-\alpha)U_\nu(f) \geq \alpha U_\nu(\hat{p}) + (1-\alpha)U_\nu(g).$$

Rearranging,

$$U_\nu(f) - U_\nu(g) \geq \frac{\alpha}{1-\alpha}\left[U_\nu(\hat{p}) - U_\nu(\hat{q})\right].$$

By the normalization obtained in the proof of Theorem 1 of Hara and Riella (2023), we have $U_\nu(\hat{p}) - U_\nu(\hat{q}) = 1$ for every $\nu \in \mathcal{V}_0$. Therefore,

$$U_\nu(f) - U_\nu(g) \geq \frac{\alpha}{1-\alpha} > 0 \quad \text{for every } \nu \in \mathcal{V}_0.$$

Hence, $f \succ g$ implies $U_\nu(f) > U_\nu(g)$ for every $\nu \in \mathcal{V}_0$. We now prove the converse implication. Suppose that $f, g \in \mathcal{F}$ satisfy $U_\nu(f) > U_\nu(g)$ for every $\nu \in \mathcal{V}_0$.

Since $\mathcal{V}_0$ is compact and the map $\nu \mapsto U_\nu(f) - U_\nu(g)$ is continuous,

$$m := \min_{\nu \in \mathcal{V}_0} \big[U_\nu(f) - U_\nu(g)\big] > 0.$$

Similarly, define $M := \max_{\nu \in \mathcal{V}_0} \big[U_\nu(\hat{p}) - U_\nu(\hat{q})\big]$. By the normalization obtained from Theorem 1, $M = 1$.

Choose $\alpha \in (0,1)$ sufficiently small such that $(1-\alpha)m > \alpha M$. Then, for every $\nu \in \mathcal{V}_0$, the affinity of U_ν implies

$$\begin{aligned} &U_\nu\big(\alpha\hat{q} + (1-\alpha)f\big) - U_\nu\big(\alpha\hat{p} + (1-\alpha)g\big) \\ &\quad = (1-\alpha)\big[U_\nu(f) - U_\nu(g)\big] - \alpha\big[U_\nu(\hat{p}) - U_\nu(\hat{q})\big] \\ &\quad \geq (1-\alpha)m - \alpha M > 0. \end{aligned}$$

Hence, by the representation of $\succsim^*$, $\alpha\hat{q} + (1-\alpha)f \succsim^* \alpha\hat{p} + (1-\alpha)g$.

By the definition of $\succsim^*$, for every $\beta \in (0,1)$,

$$\begin{aligned}&\beta\hat{p} + (1-\beta)\big[\alpha\hat{q} + (1-\alpha)f\big]\\ &\quad\succ \beta\hat{q} + (1-\beta)\big[\alpha\hat{p} + (1-\alpha)g\big].\end{aligned}$$

In particular, let $\beta = \frac{\alpha}{1+\alpha}$. Since $\beta = (1-\beta)\alpha$, the preceding comparison becomes

$$\begin{aligned}&\beta\hat{p} + (1-\beta)\big[\alpha\hat{q} + (1-\alpha)f\big]\\ &\quad\succ \beta\hat{q} + (1-\beta)\big[\alpha\hat{p} + (1-\alpha)g\big].\end{aligned}$$

Simplifying,

$$\begin{aligned}&\frac{2\alpha}{1+\alpha}\left(\frac{\hat{p}+\hat{q}}{2}\right) + \frac{1-\alpha}{1+\alpha}f\\ &\quad\succ \frac{2\alpha}{1+\alpha}\left(\frac{\hat{p}+\hat{q}}{2}\right) + \frac{1-\alpha}{1+\alpha}g.\end{aligned}$$

Since $\frac{1-\alpha}{1+\alpha} \in (0,1)$, Independence implies $f \succ g$.

Therefore,

$$f \succ g \iff U_\nu(f) > U_\nu(g) \quad \text{for every } \nu \in \mathcal{V}_0.$$

It remains to show that $\mathcal{V}$ can be chosen convex. Let $\mathcal{V}_0$ be the compact representing set obtained above and define $\mathcal{V} := \overline{\text{co}}(\mathcal{V}_0)$, where the closure is taken in the supremum norm. Since $C(X \times S)$ is a Banach space, the closed convex hull of a compact set is compact. Moreover, for any $f \succ g$, continuity and compactness imply $m := \min_{\nu\in\mathcal{V}_0}[U_\nu(f) - U_\nu(g)] > 0$. By linearity in ν, this lower bound is preserved under convex combinations and, by continuity, under closure. Thus $U_\nu(f) > U_\nu(g)$ for every $\nu \in \mathcal{V}$. The converse follows from $\mathcal{V}_0 \subseteq \mathcal{V}$. Hence, convexification preserves the representation.

■

6.2 Proof of Theorem 1

For any $f, g \in \mathcal{F}$, define the relation $\succ_c$ by

$$f \succ_c g \quad \iff \quad g \notin c(\{f,g\}).$$

Claim 1 *The relation $\succ_c$ is a strict partial order that satisfies axioms (S1)–(S3).*

Proof. Nonemptiness of $c(A)$ for every $A \in \mathcal{A}$, together with Pseudo-WARP, implies that $\succ_c$ is a strict partial order. Independence of $c(\cdot)$ implies that $\succ_c$ satisfies Independence. Best and Worst implies Constant Non-Triviality: indeed, $p_* \notin c(\{p^*, p_*\})$, so $p^* \succ_c p_*$. Similarly, Continuity of $c(\cdot)$ implies Open Continuity of $\succ_c$. Therefore, $\succ_c$ satisfies all the properties required to apply Proposition 1. ■

Now fix $f \in \mathcal{F}$ and $A \in \mathcal{A}$ such that $f \in c(A)$.

Claim 2 *For every $A \in \mathcal{A}$ and $f \in c(A)$, the set $\{B \in \mathcal{A} : A \subseteq B \text{ and } f \in c(B)\}$ has a maximal element, A_f, with respect to set inclusion.*

Proof. Suppose $\mathcal{B}$ is a subset of $\{B \in \mathcal{A} : A \subseteq B \text{ and } f \in c(B)\}$ linearly ordered by $\supseteq$. Let $\bar{B} := \text{cl}(\bigcup \mathcal{B})$. This implies that there exists a sequence (B^m) of finite subsets of $\bar{B}$ such that $B^m \to \bar{B}$. By the definition of $\bar{B}$, we can find such a sequence with $B^m \subseteq \bigcup \mathcal{B}$ for every $m \in \mathbb{N}$. Finally, since each B^m is finite and $\mathcal{B}$ is linearly ordered by $\supseteq$, we can find, for each $m \in \mathbb{N}$, a set $\bar{B}^m \in \mathcal{B}$ with $B^m \subseteq \bar{B}^m$. But then, $\bar{B}^m \to \bar{B}$ and $f \in c(\bar{B}^m)$ for every $m \in \mathbb{N}$. Continuity implies that $f \in c(\bar{B})$. That is, $\bar{B}$ is an upper bound for $\mathcal{B}$. Now Zorn's lemma implies that the collection $\{B \in \mathcal{A} : A \subseteq B \text{ and } f \in c(B)\}$ has a maximal element. ■

Claim 3 *Both A_f and $\mathcal{F} \setminus A_f$ are convex.*

Proof. Since $f \in c(A_f)$, if $g, h \in A_f$ and $\alpha \in (0,1)$, No-Hedging and maximality of A_f immediately imply that $\alpha g + (1-\alpha)h \in A_f$. Thus, A_f is convex.

To prove that $\mathcal{F} \setminus A_f$ is convex, suppose that $g \notin A_f$ and $h \notin A_f$, and let $\alpha \in (0,1)$. By maximality of A_f, $f \notin c(A_f \cup \{g\})$ and $f \notin c(A_f \cup \{h\})$. Therefore,

$$f \notin c(A_f \cup \{g\}) \cup c(A_f \cup \{h\}).$$

Indirect Ambiguity Aversion then implies that

$$f \notin c\big(A_f \cup \{\alpha g + (1-\alpha)h\}\big).$$

Since $f \in c(A_f)$, it follows that $\alpha g + (1-\alpha)h \notin A_f$. Hence, $\mathcal{F} \setminus A_f$ is convex. ■

By Claim 1 and Proposition 1, let $\mathcal{V} \subseteq C(X \times S)$ be a nonempty compact and convex set representing $\succ_c$. By Claim 2, for any fixed $f \in c(A)$, there exists A_f such that $f \in c(A_f)$ and $A \subseteq A_f$. Pseudo-WARP implies that, for every $g \in A_f$, $f \in c(\{f, g\})$. Therefore,

$$\begin{aligned} f \in c(A) &\implies \forall g \in A_f, \quad g \not\succ_c f \\ &\implies \forall g \in A_f, \ \exists \nu \in \mathcal{V} : \quad U_\nu(f) \geq U_\nu(g) \\ &\iff \min_{g \in A_f} \max_{\nu \in \mathcal{V}} \big[U_\nu(f) - U_\nu(g)\big] \geq 0. \end{aligned}$$

Since A_f and $\mathcal{V}$ are compact and convex, and $U_\nu(f) - U_\nu(g)$ is continuous and affine in each variable, Sion's Minimax Theorem gives

$$\min_{g \in A_f} \max_{\nu \in \mathcal{V}} \big[U_\nu(f) - U_\nu(g)\big] = \max_{\nu \in \mathcal{V}} \min_{g \in A_f} \big[U_\nu(f) - U_\nu(g)\big].$$

It follows that

$$f \in c(A) \implies \exists \nu \in \mathcal{V}, \ \forall g \in A_f, \quad U_\nu(f) \geq U_\nu(g).$$

And therefore,

$$c(A) \subseteq \bigcup_{\nu \in \mathcal{V}} \arg\max_{g \in A} U_\nu(g).$$

References

Aizerman, M., and A. Malishevski (1981). General theory of best variants choice: Some aspects. *IEEE Transactions on Automatic Control*, 26(5), 1030–1040.

Hara, K., and G. Riella (2023). Multiple tastes and beliefs with an infinite prize space. *Economic Theory*, 76(2), 417–444.

Heller, Y. (2012). Justifiable choice. *Games and Economic Behavior*, 76(2), 375–390.

Ok, E. A., P. Ortoleva, and G. Riella (2012). Incomplete preferences under uncertainty: Indecisiveness in beliefs versus tastes. *Econometrica*, 80(4), 1791–1808.

Riella, G. (2015). On the representation of incomplete preferences under uncertainty with indecisiveness in tastes and beliefs. *Economic Theory*, 58(3), 571–600.

Safonov, E. (n.d.). Framing of ambiguity. Available at SSRN: `https://ssrn.com/abstract=5260555`.