

The μ -calculus' Alternation Hierarchy is Strict over Non-Trivial Fusion Logics

Leonardo Pacheco
TU Wien

20 February 2024

Available at: leonardopacheco.xyz/slides/fics2024.pdf

THE μ -CALCULUS

modal μ -calculus = modal logic + fixed-point operators

- ▶ μ : least fixed-point operator
- ▶ ν : greatest fixed-point operator

ALTERNATION DEPTH

The valuation of νX and μY depend on each other:

$$\nu X. \overbrace{\mu Y. (P \wedge \Diamond X) \vee (\neg P \wedge \Diamond Y)}^{\text{scope of } \mu Y}$$

scope of νX

Alternation depth of φ

Maximum number of codependent alternating μ and ν operators in φ .

Alternation hierarchy

Classifies μ -formulas with respect to their alternation depth.

EXISTING RESULTS

Theorem (Bradfield [2])

The μ -calculus alternation hierarchy is strict over all frames.

Theorem (Alberucci–Facchini [1])

The μ -calculus alternation hierarchy collapses to the alternation hierarchy over transitive frames.

Theorem (Alberucci–Facchini [1])

The μ -calculus alternation hierarchy collapses to modal logic over equivalence relations.

(For a survey, see [4].)

OUR RESULT — SIMPLIFIED

The fusion $S5 \otimes S5$ contains two independent modalities \Box_0 and \Box_1 satisfying S5.

Theorem

The μ -calculus' alternation hierarchy is strict over the fusion $S5 \otimes S5$.

This holds for the fusion of any two non-trivial logics.

SEMANTICS

Let $M = \langle W, R_0, R_1, V \rangle$ be a Kripke model. Then:

- ▶ $M, w \models \Box_i \varphi$ iff, for all v , if $wR_i v$ then $M, v \models \varphi$, $i \in \{0, 1\}$;
- ▶ $M, w \models \Diamond_i \varphi$ iff there is v such that $wR_i v$ and $M, v \models \varphi$, $i \in \{0, 1\}$.

Given a μ -formula φ , define:

$$\Gamma_{\varphi(X)}(A) \rightarrow \|\varphi(A)\|^M.$$

Then:


- ▶ $M, w \models \mu X. \varphi$ iff w is in the least fixed point of $\Gamma_{\varphi(X)}$;
- ▶ $M, w \models \nu X. \varphi$ iff w is in the greatest fixed point of $\Gamma_{\varphi(X)}$.

ALTERNATION HIERARCHY

- ▶ $\Sigma_0^\mu (= \Pi_0^\mu)$ is the set of all μ -formulas with no fixed-point operators.
- ▶ Σ_{n+1}^μ is the closure of $\Sigma_n^\mu \cup \Pi_n^\mu$ under propositional operators, modal operators, μX , and the substitution: if $\varphi(X) \in \Sigma_{n+1}^\mu$ and $\psi \in \Sigma_{n+1}^\mu$ are such that no free variable of ψ becomes bound in $\varphi(\psi)$, then $\varphi(\psi) \in \Sigma_{n+1}^\mu$.
- ▶ Π_{n+1}^μ is the closure of $\Sigma_n^\mu \cup \Pi_n^\mu$ under propositional symbols, modal operators, νX , and the analogous substitution: if $\varphi(X) \in \Pi_{n+1}^\mu$ and $\psi \in \Pi_{n+1}^\mu$ are such that no free variable of ψ becomes bound in $\varphi(\psi)$, then $\varphi(\psi) \in \Pi_{n+1}^\mu$.

GAME SEMANTICS

Verifier and Refuter discuss whether $\Box\mu X.P \vee \Diamond X$ holds at w .



V : $\Box\mu X.P \vee \Diamond X$ holds at w

R : $\mu X.P \vee \Diamond X$ fails at v_1

V : $P \vee \Diamond X$ holds at v_1

V : $\Diamond X$ holds at v_1

V : X holds at v_2

V : $P \vee \Diamond X$ holds at v_2

V : P holds at v_2

On an infinite run, if the variable with biggest scope which repeats infinitely often is ν , then Verifier wins. Kripke semantics and game semantics are equivalent.


PARITY GAMES

- ▶ $\mathcal{P} = \langle V_{\exists}, V_{\forall}, v_0, E, \Omega \rangle$
- ▶ Two players \exists and \forall move a token in the graph $\langle V_{\exists} \cup V_{\forall}, E \rangle$ starting at v_0 .
- ▶ \exists wins $\rho = v_0, v_1, v_2, \dots$ iff the greatest priority $\Omega(v_i)$ which appears infinitely often in ρ is even.

Proposition



Evaluation games are parity games.

PARITY GAMES AS UNIMODAL KRIPKE FRAMES



$$W_n := \eta X_n \dots \nu X_0. \bigvee_{0 \leq j \leq n} [(P_j \wedge P_{\exists} \wedge \Diamond X_j) \vee (P_j \wedge P_{\forall} \wedge \Box X_j)].$$

PARITY GAMES AS $S5 \otimes S5$ FRAMES


 \Rightarrow


BIMODAL WINNING REGION FORMULAS

$$W'_n := \eta X_n \dots \nu X_0. \bigvee_{0 \leq j \leq n} [(P_j \wedge P_\exists \wedge \blacklozenge X_j) \vee (P_j \wedge P_\forall \wedge \blacksquare X_j)].$$

BIMODAL WINNING REGION FORMULAS

$$W'_n := \eta X_n \dots \nu X_0. \bigvee_{0 \leq j \leq n} [(P_j \wedge P_{\exists} \wedge \blacklozenge X_j) \vee (P_j \wedge P_{\forall} \wedge \blacksquare X_j)].$$

Where

- ▶ $\blacklozenge \varphi := \mu Y. \text{pre}_0 \wedge \text{bd} \wedge \Diamond_0(\text{nxt}_0 \wedge \text{pre}_1 \wedge \text{bd} \wedge \Diamond_1(\text{nxt}_1 \wedge \text{bd} \wedge ((Y \wedge \neg \text{st}) \vee (\varphi \wedge \text{st})))));$ and
- ▶ $\blacksquare \varphi := \nu Y. \text{pre}_0 \wedge \text{bd} \rightarrow \Box_0(\text{nxt}_0 \wedge \text{pre}_1 \wedge \text{bd} \rightarrow \Box_1(\text{nxt}_1 \wedge \text{bd} \rightarrow ((Y \wedge \neg \text{st}) \wedge (\varphi \wedge \text{st})))),$

PROOF SKETCH

- ▶ Let n be even. Then $W_n \in \Pi_{n+1}^\mu$.
- ▶ Suppose that W_n is equivalent to some formula in Π_n^μ . Let $\varphi \in \Sigma_n^\mu$ be equivalent to $\neg W_n$.
- ▶ Define $f_{\varphi \wedge \varphi}(M, w) = (\mathcal{G}^K(M, w \models \varphi \wedge \varphi), \langle w, \varphi \wedge \varphi \rangle)$.
- ▶ Let (M, w) be a fixed-point of $f_{\varphi \wedge \varphi}$. Then

$$\begin{aligned} M, w \models \neg W_n &\iff M, w \models \varphi \wedge \varphi \\ &\iff f_{\varphi \wedge \varphi}(M, w) \models W_n \\ &\iff M, w \models W_n. \end{aligned}$$

- ▶ This is a contradiction.

OUR RESULT

Theorem

Let F_0 , F_1 , and F_2 be classes of unimodal Kripke frames closed under isomorphic copies and disjoint unions. If

- 1. $\circ \leftarrow \circ \rightarrow \circ$ is a subframe of F_0 and $\circ \rightarrow \circ$ a subframe of F_1 ; or*
- 2. $\circ \rightarrow \circ \rightarrow \circ$ is a subframe of F_0 and $\circ \rightarrow \circ$ a subframe of F_1 ;*

then the μ -calculus' alternation hierarchy is strict over $F_0 \otimes F_1$. If

- 3. $\circ \rightarrow \circ$ is a subframe of F_0 , F_1 , and F_2 ;*

then the μ -calculus' alternation hierarchy is strict over $F_0 \otimes F_1 \otimes F_2$.

Conjecture

Suppose $\circ \rightarrow \circ$ is a subframe of F_0 and F_1 . Then every μ -formula is equivalent to one with alternation depth 1 over $F_0 \otimes F_1$.

COLLAPSE ON MULTIMODAL LOGICS

GLP is a provability logic which contains countably many modal operators.

Theorem (Ignatiev [3])

GLP has the fixed-point property.

IS5 is an intuitionistic version of S5 which can be treated as a bimodal logic.

Theorem (P. [5])

The μ -calculus collapses to modal logic over IS5.

REFERENCES

- [1] L. Alberucci, A. Facchini, “The modal μ -calculus hierarchy over restricted classes of transition systems”, 2009.
- [2] J.C. Bradfield, “Simplifying the modal mu-calculus alternation hierarchy”, 1998.
- [3] K.N. Ignatiev, “On Strong Provability Predicates and the Associated Modal Logics”, 1993.
- [4] L. Pacheco, “Exploring the difference hierarchies on μ -calculus and arithmetic—from the point of view of Gale–Stewart games”, PhD Thesis, 2023.
- [5] L. Pacheco, “Game Semantics for the Constructive μ -Calculus”, arXiv:2308.16697, 2023.