Beyond mean field: does the phase-space approximation capture three-body neutrino correlations?

Giovanna Fátima de Britto Vieira[1], **Marco Ruas Sales Peixoto**[1]
Leonardo Villas Boas Fischel[1]

[1]Instituto de Tecnologia e Liderança – Inteli
São Paulo – SP – Brasil
Orientadores: Jose Erinaldo da Fonseca e Bryan Kano Ferreira

Abstract. *Collective neutrino oscillations motivate methods capable of describing quantum many-body effects. The phase-space approximation offers a lower-cost alternative, but its accuracy for three-body correlations requires specific assessment. This work aims to establish its validity regimes in a two-flavor model, using exact evolution as a reference. Flavor probabilities and connected pair and triple cumulants will be compared across interaction strengths and system sizes. Trotterized circuits will be assessed with and without error mitigation. The expected contribution is to establish the approximation's accuracy and the noise and measurement thresholds that allow its predictions to be distinguished from exact results.*

Resumo. *As oscilações coletivas de neutrinos motivam métodos capazes de descrever efeitos quânticos de muitos corpos. A aproximação de espaço de fase oferece uma alternativa de menor custo, mas sua precisão para correlações de três corpos requer avaliação específica. Este trabalho busca delimitar seus regimes de validade em um modelo de dois sabores, usando a evolução exata como referência. Serão comparados probabilidades de sabor e cumulantes conectados de pares e trios, variando a interação e o tamanho do sistema. Circuitos trotterizados serão avaliados com e sem mitigação de erros. Espera-se estabelecer a precisão da aproximação e os limiares de ruído e de medições que permitem distinguir suas previsões dos resultados exatos.*

1. Introduction

1.1. Astrophysical context and motivation

Stars with initial masses above approximately eight solar masses can undergo gravitational core collapse. In a typical scenario, an iron-rich core loses support, contracts, and undergoes a bounce, generating a shock. This shock can stall through energy losses from nuclear dissociation and neutrino emission. Neutrino absorption and fluid motion can revive it during accretion, meaning matter falling onto the remnant (Figure 1) [Janka 2012, Abi et al. 2020].

Neutrinos are electrically neutral elementary particles that interact weakly with matter. A typical supernova emits about 10^{58} neutrinos, with characteristic energies of $10\,\mathrm{MeV}$, over approximately ten seconds, carrying more than 99% of the gravitational




Figure 1. Stellar core collapse, not to scale. (a) Core contraction and bounce. (b) Stalled shock and neutrino heating around the proto-neutron star (PNS). (c) Explosion or black-hole formation, including after an explosion. Source: original illustration based on Janka (2012).

energy released by the collapsed core [Abi et al. 2020]. Their flavor states are superpositions of mass states, whose propagation phases change detection probabilities, producing oscillations [Duan et al. 2010].

Flavor evolution can alter heating behind the shock and the neutron-to-proton composition relevant to nucleosynthesis. Its effects depend on stellar configuration, as shown by Nagakura (2023) and Akaho et al. (2026); the relationship between neutron-rich matter and heavy-element formation is discussed by Cowan et al. (2021). Events from SN 1987A and detection prospects at DUNE, Hyper-Kamiokande, and JUNO reinforce the importance of predicting emitted flavors [Vissani et al. 2010, Baum et al. 2022, Sarmah et al. 2026]. Near the core, coherent interactions between neutrinos couple their evolution, generating collective oscillations and the many-body problem discussed next [Duan et al. 2010].

1.2. State of the art and research gap

The collective dynamics described above constitutes a quantum many-body problem in which correlations and entanglement can develop between neutrinos, with entanglement associated with the nonseparability of the joint quantum state. In the general two-flavor representation, a pure state of N neutrinos has 2^N complex amplitudes. This growth limits exact evolution of large systems on classical computers when no specific symmetries or compression schemes are exploited [Duan et al. 2010, Cervia et al. 2019].

One strategy to reduce this cost is the mean-field approximation, which replaces the effects of other particles with an effective field determined by their averages. By factorizing expectation values of operators acting on distinct particles, this description neglects connected correlations and, for initially factorized states, does not track the genera-

tion of entanglement between neutrinos. Cervia et al. (2019) related differences between exact and mean-field evolution to the development of entanglement, with a dependence on initial conditions. In a three-flavor model, Mangin-Brinet, Bauge, and Lacroix (2026) also observed short-time deviations in the configurations examined. The validity of the approximation must therefore be assessed in terms of the model, initial conditions, and dynamical regime.

The phase-space approximation (PSA) seeks to incorporate effects beyond mean field through an ensemble of independent trajectories. Initial conditions are sampled to reproduce selected quantum moments, and each trajectory evolves according to mean-field equations; observables are estimated by statistical averages over the ensemble. Lacroix et al. (2022) obtained good agreement with mean values and entropies of one- and two-neutrino reduced states, calculated from their respective density matrices, in a two-beam model, although they did not reproduce long-time oscillations of fluctuations. The extension to multiple beams also compared collective two-body correlations through the covariance-based observable C_{ZZ}. This collective validation is not equivalent to assessing every pair individually and does not establish the accuracy of third-order cumulants [Lacroix et al. 2024]. The extension to three flavors was subsequently compared with exact solutions for up to eight neutrinos [Mangin-Brinet et al. 2026].

Another approach beyond mean field truncates the Bogoliubov–Born–Green–Kirkwood–Yvon (BBGKY) hierarchy, which relates the evolution of one-particle observables to higher-order correlations. In the preprint by Riccio and Balantekin (2026), the second-order closure retains pair contributions and sets connected three-body cumulants to zero to close the equations. This condition does not necessarily set the expectation value of a product of three operators to zero, because contributions constructed from individual means and pairs remain. PSA incorporates higher moments without imposing the same closure; its accuracy for third-order cumulants must be examined directly [Lacroix et al. 2022].

Quantum simulations offer a complementary route to investigate these correlations. Hall et al. (2021) demonstrated simulations of small neutrino systems on noisy devices, using error mitigation, namely techniques to reduce the effects of noise on observable estimates, to extract entanglement measures. Illa and Savage (2023) investigated systems of up to twelve neutrinos on the Quantinuum H1-1 trapped-ion processor, including two- and three-body correlations and multipartite entanglement indicators. At a larger scale, Kiss et al. (2026) studied thermalization with random circuits on IBM processors using more than one hundred qubits. Their performance was assessed through thermalization observables; the fidelity of specific cumulants during evolution requires separate validation.

Within the literature examined, no systematic comparison of PSA three-body cumulants with exact evolution was identified that controls system size, interaction strength, and computational and measurement uncertainties. Agreement in flavor averages, collective pair correlations, or reduced entropies does not guarantee accurate third-order cumulants. Addressing this gap requires consistent observables and a separation of approximation error, finite sampling, time discretization, and noise.

The accuracy of third-order cumulants matters because, for a Hamiltonian

with pair interactions, the evolution hierarchy couples one-particle observables to two-body correlations, whose equations depend on third-order moments. Deviations in the description of triples may therefore affect lower-order observables during evolution [Lacroix et al. 2024, Riccio and Balantekin 2026]. The reported long-time limitations of PSA motivate examining this relationship, without attributing all flavor deviations to a single cumulant component in advance [Lacroix et al. 2022].

The quantum stage seeks to establish the precision with which circuits can provide a reference for testing approximations. In larger systems, where exact classical calculations become inaccessible, this possibility will depend on the accuracy of measured observables. Small systems allow this condition to be assessed against a known reference and discretization, sampling, and noise errors to be separated. The study assumes neither a computational advantage at the sizes investigated nor an automatic extrapolation of its results to larger systems.

This work therefore asks in which regimes of interaction strength, system size, and evolution time PSA reproduces connected two- and three-body correlations and, when deviations occur, under which noise conditions and measurement counts quantum circuits can distinguish them from exact results.

To this end, the study proposes to establish PSA accuracy against exact calculations, with mean field as an additional reference, in a two-flavor model with factorized pure-flavor initial states. Systems of four to approximately sixteen neutrinos will be compared, with circuits for the subset of four to eight. The ratio $\mu/\bar{\omega}$ will range from 0.1 to 10, where μ denotes the neutrino–neutrino interaction scale and $\bar{\omega}$ the mean vacuum oscillation frequency. The supernova context motivates the model; conclusions will remain restricted to the simplified configurations examined.

Increasing interaction strength increases the coefficients of the terms coupling the hierarchy's orders, motivating the expectation that triples are more sensitive to effects beyond mean-field trajectories. This is a conjecture dependent on the initial state and time window, not a consequence of PSA setting higher cumulants to zero. Hypothesis **H1** predicts that PSA reproduces flavor probabilities and pair cumulants more accurately than triple cumulants and that, when the reference permits normalization, the error for triples tends to grow with interaction strength. Hypothesis **H2** predicts that mitigation increases the noise threshold at which PSA can be distinguished from exact evolution at the same total measurement budget. Their falsification criteria will be specified in the research design.

Section 2 presents the model, observables, comparison stages, and hypothesis assessment criteria. Section 3 describes the planned presentation of results. The present text sets out the research design; experimental results will be incorporated after the protocol has been executed and validated.

2. Research design

2.1. Model and initial states

A two-flavor model with distinct energies and interactions between all pairs will be adopted. The electron and effective nonelectron flavors will be represented by the states $|0\rangle = |\nu_e\rangle$ and $|1\rangle = |\nu_x\rangle$ of a qubit, the basic unit of quantum information. Initial states

will be products of pure flavor states. The main comparison will use even-sized systems with half the neutrinos in each flavor; frequencies, couplings, and flavor assignments will be recorded for each scenario. Other initial configurations will be treated separately because approximation deviations depend on these conditions [Cervia et al. 2019].

The Hamiltonian and its normalization will be shared by all methods. A published configuration will be reproduced before parameter scans, checking flavor, frequency, and coupling conventions. Comparisons will use common windows of dimensionless time $\bar{\omega}t$, also reporting μt, which varies when interaction strength changes. The range from four to sixteen neutrinos defines the main study; an extension to twenty or twenty-two will depend on timing and memory measurements for exact calculations. Dependence on N will be investigated without presupposing a scaling law that has not yet been justified.

2.2. Observables and connected correlations

To make the meaning of correlations explicit, consider the flavor operator $Z_i = |\nu_e\rangle_i\langle\nu_e| - |\nu_x\rangle_i\langle\nu_x|$, equivalent to the Pauli matrix σ_z on qubit i, where ν_x is the effective nonelectron flavor. Its mean is $m_i(t) = \langle Z_i\rangle_t$, and the electron-flavor probability is $P_{e,i}(t) = [1 + m_i(t)]/2$. For distinct particles, these operators commute, and second- and third-order cumulants are defined as

$$C^{(2)}_{ij}(t) = \langle Z_i Z_j\rangle_t - m_i(t)m_j(t)$$

and

$$\begin{aligned} C^{(3)}_{ijk}(t) = {} & \langle Z_i Z_j Z_k\rangle_t - m_i(t)m_j(t)m_k(t) \\ & - C^{(2)}_{ij}(t)m_k(t) - C^{(2)}_{ik}(t)m_j(t) - C^{(2)}_{jk}(t)m_i(t). \end{aligned}$$

Figure 2 represents this decomposition: a three-particle moment combines products of means, pair contributions, and a connected three-body term. Neglecting $C^{(3)}_{ijk}$ in a closure is therefore not equivalent to neglecting $\langle Z_i Z_j Z_k\rangle$. This identity is a particular case of the cumulant expansion discussed by Riccio and Balantekin (2026). The main analysis will use the Z, ZZ, and ZZZ components, with particle sets specified in the protocol; other components will require separate assessment.

2.3. Method comparison and error control

Figure 3 organizes the proposed protocol. For each system size, interaction strength, and initial state, the same Hamiltonian and observable definitions will be used for exact, mean-field, PSA, and circuit evolution. Initial conditions sampled by PSA must reproduce the moments prescribed by the method for the chosen physical state. Systems of four to eight neutrinos will be shared by all four methods; classical comparisons will extend to approximately sixteen neutrinos. This organization combines the systematic exact-reference comparisons of Lacroix et al. (2024) with the distinction between exact evolution, Trotterization, and measurements used by Hall et al. (2021) and Illa and Savage (2023).

Before interpreting physical differences, convergence will be checked by refining time integration, increasing the number of PSA trajectories, and reducing the Trotter step




Figure 2. Decomposition of the moment $\langle Z_i Z_j Z_k \rangle$. (a) Product of individual means $m_i m_j m_k$. (b) Three contributions consisting of a pair cumulant times the mean of the remaining particle. (c) Connected cumulant $C_{ijk}^{(3)}$, highlighted as the part remaining after subtracting the preceding terms. Circles identify distinct particles i, j, and k; one link denotes a pair contribution, and the three-linked diamond denotes the triple cumulant without representing another particle. This statistical notation represents neither additional Hamiltonian interactions nor an entanglement certificate. All terms are evaluated at the same time. Source: original illustration based on the cumulant expansion of Riccio and Balantekin (2026).

in ideal simulation. Noisy assessments will follow this validation. Cumulant uncertainties must account for the fact that means and products may be estimated from the same measurements or trajectories, preserving their covariances during error propagation or resampling.

2.4. Errors, uncertainties, and assessment of H1

The same observables, flavor basis, and normalization will be used in every method. A nonzero cumulant alone does not certify genuine entanglement. Absolute error will be reported over time and summarized by the root mean square of deviations (RMSE). For each order r, normalized error will be $E_r = \mathrm{RMSE}_r / R_r$, where R_r is the root mean square of the exact reference over the same time grid and observable set for that order. The same definition will be applied to flavor probabilities.

Normalization will be used only when $R_r > \varepsilon_r$, with $\varepsilon_r = \max(10\delta_r, 10^{-6})$. Here, δ_r is the largest absolute change in the reference observed under numerical refinement. The dimensionless floor 10^{-6} is an operational choice to avoid near-zero denominators; sensitivity of classification to a factor-of-ten change in this floor will be reported. Cases below the threshold will retain absolute errors and will not enter normalized comparisons. This rule will not be changed after confirmatory results have been observed.

To assess H1, differences between errors for triples, pairs, and probabilities will be compared within the same scenarios and windows. The trend with interaction strength will be estimated at fixed N, initial state, frequencies, and relative couplings; its direction will be assessed using a trend measure recorded in advance. The hypothesis will be contradicted if errors for triples are smaller than or equivalent to those of the other




Figure 3. Proposed comparison protocol. (a) A Hamiltonian H and initial state $|\Psi_0\rangle$ shared by the methods, in the two-flavor model with distinct energies and $\mu/\bar{\omega}$ between 0.1 and 10. The four-node network illustrates interactions between all pairs; it represents neither processor connectivity nor a limit on system size. (b) Exact, mean-field, and PSA evolution for four to approximately sixteen neutrinos; circuits for the subset of four to eight, under ideal and noisy conditions. (c) Comparison of probabilities and cumulants against exact evolution, highlighting $C_{ijk}^{(3)}$. The bottom row groups integration, sampling, Trotter, measurement, and noise controls. Comparisons with and without mitigation will use the same total measurement budget; hardware execution will depend on availability. Source: original illustration informed by the comparisons of Lacroix et al. (2024), Hall et al. (2021), and Illa and Savage (2023).

quantities within the specified margin, or if the increasing trend with interaction strength is absent within that margin or reversed. Uncertainties preventing the comparison from being resolved will be reported as inconclusive, rather than as evidence of equivalence.

PSA trajectory counts will be refined by doubling from 1,024, using at least five independent seeds. The convergence criterion will require changes in estimates and interval widths to remain below the established observable tolerances. The protocol will fix the trajectory limit and required precision before confirmatory analysis; cases not meeting this precision will be identified. Resampling will use complete trajectories or bitstring counts, preserving dependencies between the means and products composing each cumulant.

2.5. Circuits, mitigation, and assessment of H2

Circuits will be constructed using a symmetric second-order Trotter formula. Trotterization approximates the full evolution through operations associated with Hamiltonian terms over small time steps. The proposed initial step will be $\Delta\tau = 0.05$, with $\tau = \bar{\omega}t$,

followed by successive halvings. Ideal comparison with the exact reference must reduce this error to a predefined fraction of the PSA–exact discrepancy to be measured. This control will precede noise assessment.

Measurements in the flavor basis will simultaneously provide the products Z_i, Z_iZ_j, and $Z_iZ_jZ_k$. Each cumulant will be reconstructed from the same counts, with intervals obtained by joint resampling. The initial budget scan will include 1,024, 2,048, 4,096, and 8,192 measurements, with possible expansion during the pilot study. The total budget will include measurements used in calibration and auxiliary circuits; mitigated and unmitigated results will be compared under the same limit.

The main assessment of H2 will consider independent symmetric readout error, parametrized by the bit-flip probability $0 \leq p < 1/2$, and mitigation through a readout calibration matrix. Intervals will include calibration uncertainty. The effects of depolarizing gate noise and zero-noise extrapolation may be examined as extensions, with resources and results reported separately. Thresholds obtained under the readout model will not be presented as general hardware-noise limits.

A difference will be considered resolved when the confidence interval for the circuit–exact deviation lies within a predefined equivalence margin and the interval for the circuit–PSA difference excludes zero. PSA uncertainties and reference tolerances will be considered. For a total budget M, the threshold $p_{\max}(M)$ will be the largest assessed noise level satisfying this criterion with a specified minimum success frequency across independent repetitions. The smallest assessed budget allowing distinction at each noise level will also be reported. If the transition is not reached, bounds will be reported without assigning a precise threshold; nonmonotonic behavior will be shown as resolution regions.

H2 will be contradicted if the mitigated threshold is smaller than or equivalent to the unmitigated threshold within the specified comparison margin and uncertainties. An unresolved difference will be considered inconclusive. Execution on IBM hardware will depend on availability and will be reported with device conditions, without replacing the controlled noise-model scan.

2.6. Preregistration and assessment limits

Before confirmatory comparisons, the Hamiltonian, initial configurations, interaction and size grid, time windows, particle sets, tolerances, equivalence margins, confidence level, and treatment of multiple comparisons will be fixed. Trajectory limits, seeds, Trotter steps, noise levels, total budgets, and their allocation between calibration and execution will also be recorded. Initial values proposed in this section may be adjusted during the pilot, with justification recorded before confirmatory cases are analyzed.

Quantum scenarios will be selected through a rule defined from the classical comparison, requiring a PSA–exact discrepancy resolved by the uncertainties. Selection of triples and times will precede noisy runs. The assessment will thus distinguish physical limitations of the approximation, insufficient sampling, and a circuit's inability to resolve a known difference.

3. Planned presentation of results

Results will be presented in separate panels for probabilities, pair cumulants, and triple cumulants. Line and symbol meanings will remain consistent: exact evolution as a solid black line, mean field as a dotted gray line, PSA as a dashed blue line, and circuit estimates as symbols with uncertainty bars. Circles and squares will distinguish unmitigated and mitigated estimates without relying on color alone. Captions will specify the initial state, N, $\mu/\bar{\omega}$, dimensionless time $\bar{\omega}t$, sampling resources, and the statistical meaning of the bars. Deviation and normalized-error panels will complement the curves according to the established criteria; error tables or maps will summarize dependence on system size and interaction strength when sufficient data are available. Gate diagrams will show the implemented circuit and its Trotter order.

Comparisons will be organized by initial state and time window. Noise thresholds and the smallest assessed budgets will be reported with uncertainties and scan limits. Inconclusive cases will remain in the presentation, without selecting only configurations favorable to the hypotheses.

Use of generative artificial intelligence

Generative artificial intelligence assistance was used to support writing and consistency revision, translation, and the production of code for the conceptual diagrams.

References

Abi, B. et al. (2020). Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume II: DUNE Physics. Technical Report FERMILAB-PUB-20-025-ND, DUNE Collaboration. https://arxiv.org/abs/2002.03005.

Akaho, R., Nagakura, H., Iwakami, W., Furusawa, S., Harada, A., Okawa, H., Matsufuru, H., Sumiyoshi, K., and Yamada, S. (2026). Bifurcated Impact of Neutrino Fast Flavor Conversion on Core-Collapse Supernovae Informed by Multiangle Neutrino Radiation Hydrodynamics. *Physical Review Letters*, 136(19):191002. https://doi.org/10.1103/fksy-1jtw.

Baum, S., Capozzi, F., and Horiuchi, S. (2022). Rocks, water, and noble liquids: Unfolding the flavor contents of supernova neutrinos. *Physical Review D*, 106(12):123008. https://doi.org/10.1103/PhysRevD.106.123008.

Cervia, M. J., Patwardhan, A. V., Balantekin, A. B., Coppersmith, S. N., and Johnson, C. W. (2019). Entanglement and collective flavor oscillations in a dense neutrino gas. *Physical Review D*, 100(8):083001. https://doi.org/10.1103/PhysRevD.100.083001.

Cowan, J. J., Sneden, C., Lawler, J. E., Aprahamian, A., Wiescher, M., Langanke, K., Martínez-Pinedo, G., and Thielemann, F.-K. (2021). Origin of the heaviest elements: The rapid neutron-capture process. *Reviews of Modern Physics*, 93(1):015002. https://doi.org/10.1103/RevModPhys.93.015002.

Duan, H., Fuller, G. M., and Qian, Y.-Z. (2010). Collective Neutrino Oscillations. *Annual Review of Nuclear and Particle Science*, 60(1):569–594. https://doi.org/10.1146/annurev.nucl.012809.104524.

Hall, B., Roggero, A., Baroni, A., and Carlson, J. (2021). Simulation of collective neutrino oscillations on a quantum computer. *Physical Review D*, 104(6):063009. https://doi.org/10.1103/PhysRevD.104.063009.

Illa, M. and Savage, M. J. (2023). Multi-Neutrino Entanglement and Correlations in Dense Neutrino Systems. *Physical Review Letters*, 130:221003. https://doi.org/10.1103/PhysRevLett.130.221003.

Janka, H.-T. (2012). Explosion Mechanisms of Core-Collapse Supernovae. *Annual Review of Nuclear and Particle Science*, 62:407–451. https://doi.org/10.1146/annurev-nucl-102711-094901.

Kiss, O., Tavernelli, I., Tacchino, F., Lacroix, D., and Roggero, A. (2026). Neutrino thermalization via randomization on a quantum processor. *Communications Physics*, 9(1):282. https://doi.org/10.1038/s42005-026-02697-2.

Lacroix, D., Balantekin, A. B., Cervia, M. J., Patwardhan, A. V., and Siwach, P. (2022). Role of non-Gaussian quantum fluctuations in neutrino entanglement. *Physical Review D*, 106(12):123006. https://doi.org/10.1103/PhysRevD.106.123006.

Lacroix, D., Bauge, A., Yilmaz, B., Mangin-Brinet, M., Roggero, A., and Balantekin, A. B. (2024). Phase-space methods for neutrino oscillations: Extension to multibeams. *Physical Review D*, 110:103027. https://doi.org/10.1103/PhysRevD.110.103027.

Mangin-Brinet, M., Bauge, A., and Lacroix, D. (2026). Three-flavor neutrino oscillations using the phase space approach. *Physical Review D*, 113:036026. https://doi.org/10.1103/yr2g-1g2g.

Nagakura, H. (2023). Roles of Fast Neutrino-Flavor Conversion on the Neutrino-Heating Mechanism of Core-Collapse Supernova. *Physical Review Letters*, 130(21):211401. https://doi.org/10.1103/PhysRevLett.130.211401.

Riccio, M. and Balantekin, A. B. (2026). Improved Approximations for Collective Neutrino Oscillations. Preprint. https://doi.org/10.48550/arXiv.2607.28619.

Sarmah, P., Chakraborty, S., Medhi, A., Bose, D., and Devi, M. M. (2026). Neutrino mass ordering from the next Galactic supernova at DUNE, HK, and JUNO. Preprint. https://arxiv.org/abs/2606.06580.

Vissani, F., Costantini, M. L., Fulgione, W., Ianni, A., and Pagliaroli, G. (2010). What is the issue with SN1987A neutrinos? Preprint. https://arxiv.org/abs/1008.4726.