

# Methodology, Statistics and Pitfalls

## Data Science and Society Statistics Lecture 1

---

Matthieu Brinkhuis

October 4, 2018

Utrecht University, Information and Computing Sciences  
m.j.s.brinkhuis@uu.nl



# Introduction

---

Topics we will *discuss* today:

- Methodological issues in data analysis
- Some pitfalls

## Intended learning outcomes

By the end of this lecture, you'll be able to:

- Recognize that methodology for data science is crucial (1)
- Understand different types of pitfalls (2)
- Apply the principles in your own research (3)
- Analyze potential traps (4,5)

Bloom's Taxonomy:

1. Remember
2. Understand
3. Apply
4. Analyze
5. Evaluate
6. Create

# Table of Contents


1. Introduction
2. Hypothesis testing
3. Prediction
4. Inference and Causality
5. Data quality
6. Closing

# Structured method for conducting analyses

Structured method helpful in preventing methodological errors.

CRISP-DM (Chapman et al., 2000, p. 12):

1. Business understanding
2. Data understanding
3. Data preparation
4. Modeling
5. Evaluation
6. Deployment



**Figure 1:** CRISP-DM Process Diagram

# Hypothesis testing


---

## Null Hypothesis Significance Testing (NHST)

Traditional steps (I Miller and M Miller, 2014):

1. Formulate  $H_0$  and  $H_a$  (e.g.,  $H_0 : \mu = \mu_0$ ,  $H_a : \mu \neq \mu_0$ ).
2. Using the sampling distribution of the (appropriate) test statistics, determine critical region of size  $\alpha$ .
3. Determine the value of the test statistics from the sample data.
4. Check if it falls in the critical region (reject  $H_0$ ) or outside (retain  $H_0$ ).





**Figure 2:** Critical region for two-tailed test (I Miller and M Miller, 2014, p. 360).

## What is a p-value?

Depends on who you ask.

- Frequentist: limiting relative frequency, if you could repeat the experiment.
- Bayesian: subjective, degree of belief, personally defined.

Back to the example: if  $p \leq \alpha$ , the observed data is inconsistent with the null hypothesis, so the null hypothesis must be rejected. Does not prove that the tested hypothesis is true. Guarantees that the Type I error (false positive) rate is at most  $\alpha$ .

**Problems with  $p \leq .05$** 

Traditionally  $\alpha = .05$ , but there are calls for change.

“We propose to change the default P-value threshold for statistical significance from 0.05 to 0.005 for claims of new discoveries.” (Benjamin et al., 2018).

Why do you think that is?

## The problem of multiple testing

“When pursuing multiple inferences, researchers tend to select the (statistically) significant ones for emphasis, discussion and support for conclusions. An unguarded use of single-inference procedures results in a greatly increased false positive (significance) rate.” (Benjamini and Hochberg, 1995, p. 289)

How to deal with false discoveries? Be aware! Correction possible, e.g. Bonferroni and Hochberg methods, see R function `p.adjust{stats}`.

# Prediction

---

## Statistical learning

$$Y = f(X) + \epsilon \quad (1)$$

where  $f$  is unknown function of  $X_1, X_2, \dots, X_p$  and  $\epsilon$  is a random error term, independent of  $X$  (James et al., 2013).

Estimate  $\hat{f}$  for:

- prediction (black box)
- inference (interest in associations, what is the type of relationship, etc.)

There is a trade-off between prediction accuracy and model interpretability (Waa et al., 2018; James et al., 2013).

In prediction (Kaggle?), what do you predict against?

- role of testing data
- how much is there to gain? baseline construction? (burglary)
- from 90% to 100% can be a long way (reducible vs irreducible error)
- how to quantify prediction quality? (model accuracy)
- dynamic prediction (feedback loops, fraud prediction and Netflix challenge)

## Validation set approach

Basic idea is to split your data set in two parts (e.g., James et al., 2013, p. 176):

- training set (to fit the model)
- validation set (to evaluate the model)

Only a subset used, test error can be variable. Different cross-validation approaches possible (leave-one-out, k-fold, bootstrap, etc.).



## Measuring the quality of fit

- mean squared error =  $\frac{1}{n} \sum_{i=1}^n (y_i - \hat{f}(x_i))^2$
- error rate =  $\frac{1}{n} \sum_{i=1}^n (y_i \neq \hat{y}_i)$


However, there are many more ways, especially in classification.

# Classification prediction quality: discrete

		True class		Measures
		Positive	Negative	
Predicted class	Positive	True positive <i>TP</i>	False positive <i>FP</i>	Positive predictive value (PPV) $\frac{TP}{TP+FP}$
	Negative	False negative <i>FN</i>	True negative <i>TN</i>	Negative predictive value (NPV) $\frac{TN}{FN+TN}$
Measures		Sensitivity $\frac{TP}{TP+FN}$	Specificity $\frac{TN}{FP+TN}$	Accuracy $\frac{TP+TN}{TP+FP+FN+TN}$

**Figure 3:** Confusion matrix (see Wikipedia for more).

## Classification prediction quality: discrete



**Figure 4:** ROC analysis (from Wikipedia ROC).

# Inference and Causality

---

## Exploratory vs. confirmatory analysis


Important distinction in statistics. Why would you do exploratory analysis? Why confirmatory analysis?

Using this distinction: what is data mining?

Related concepts:

- Spurious relations
- Coincidence
- Causal relationships (or lack thereof)
- Fishing

# Spurious relations



**Figure 5:** Spurious Correlation example, see [tylervigen.com](http://tylervigen.com)

## Related problems

So beware of causal statements, here it seems easy, but if you relate 'obvious' data, you easily fall for it!

Pitfalls include:

- Simpson's paradox
- 3rd variable
- Anscombe's quartet (next)
- Lack of theory
- Lack of experiment


Many things interesting, without making causal statements.

# Graphs are essential

“Graphs are essential to good statistical analysis” (Anscombe, 1973)

Four data sets:


- $n = 11$
- $\bar{x} = 9.0$
- $\bar{y} = 7.5$
- $y = 3 + .5x$
- Multiple  $R^2 = .667$



**Figure 6:** Anscombe's Quartet, obtained by `example(anscombe)`



# Simpson's Paradox i




**Figure 7:** A clear positive relation...

# Simpson's Paradox ii

How is this possible?

- Graphs from Paul van der Laken ([rpubs.com/lakenp](http://rpubs.com/lakenp))
- Further reading: Simpson (1951) and Kievit, Frankenhuis, et al. (2013)
- R Package: Simpsons (Kievit and Epskamp, 2012)



**Figure 8:** ...until conditioning.

## Data quality

---


# Get a grip on the quality of your data

Data quality is a topic by itself, here some thoughts:


- Garbage-in garbage-out principle
- Examples of measures: completeness, validity, accuracy, consistency, availability and timeliness
- Beware of missing data:
  - what deleting or disregarding data can do to your research
  - but what to do then?
  - explicit assumptions
  - model when needed

# Sometimes you cannot do without models

Sometimes your data is model generated, example:



**Figure 9:** Math garden landing page.



**Figure 10:** Math garden practice item.

## Closing

---


### Practical remarks:

- Read article *and* the response letters on the Google Flu (Lazer et al., 2014a; Broniatowski, Paul, and Dredze, 2014; Lazer et al., 2014b)
- Read article on p-values (Benjamin et al., 2018)
- Tutorial on data analysis in R and Spark

# Intended learning outcomes recap

By the end of this lecture, you'll be able to:

- Recognize that methodology for data science is crucial (1)
- Understand different types of pitfalls (2)
- Apply the principles in your own research (3)
- Next lecture: *Analyze potential traps* (4,5)






**Figure 11:** Bloom's Taxonomy (image from Wikipedia).






**Thank you.**




## References i

-  Anscombe, FJ (1973). "Graphs in Statistical Analysis". In: *The American Statistician* 27.1, pp. 17–21. DOI: 10.2307/2682899. JSTOR: 2682899 (cit. on p. 24).
-  Benjamin, DJ et al. (2018). "Redefine statistical significance". In: *Nature Human Behaviour* 2.1, p. 6. DOI: 10.1038/s41562-017-0189-z (cit. on pp. 11, 31).
-  Benjamini, Y et al. (1995). "Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing". English. In: *Journal of the Royal Statistical Society. Series B (Methodological)* 57.1, pp. 289–300. URL: <http://www.jstor.org/stable/2346101> (cit. on p. 12).

## References ii

-  Broniatowski, DA et al. (2014). "Twitter: Big data opportunities". In: *Science* 345.6193, p. 148. DOI: 10.1126/science.345.6193.148-a. eprint: <http://www.sciencemag.org/content/345/6193/148.1.full.pdf> (cit. on p. 31).
-  Chapman, P et al. (2000). *CRISP-DM 1.0. Step-by-step data mining guide*. Tech. rep. The CRISP-DM consortium. URL: <ftp://ftp.software.ibm.com/software/analytics/spss/support/Modeler/Documentation/14/UserManual/CRISP-DM.pdf> (cit. on p. 6).
-  James, G et al. (2013). *An introduction to statistical learning*. Springer Texts in Statistics. Springer. DOI: 10.1007/978-1-4614-7138-7 (cit. on pp. 14, 16).

## References iii

-  Kievit, RA, WE Frankenhuis, et al. (2013). “Simpson’s paradox in psychological science: a practical guide”. In: *Frontiers in Psychology* 4. DOI: 10.3389/fpsyg.2013.00513 (cit. on p. 26).
-  Kievit, RA et al. (2012). *Simpsons: Detecting Simpson’s Paradox*. R package version 0.1.0. URL: <https://CRAN.R-project.org/package=Simpsons> (cit. on p. 26).
-  Lazer, D et al. (2014a). “The Parable of Google Flu: Traps in Big Data Analysis”. In: *Science* 343.6176, pp. 1203–1205. DOI: 10.1126/science.1248506 (cit. on p. 31).

## References iv



Lazer, D et al. (2014b). “Twitter: Big data opportunities - Response”. In: *Science* 345.6193, pp. 148–149. DOI: 10.1126/science.345.6193.148-b. eprint: <http://www.sciencemag.org/content/345/6193/148.2.full.pdf> (cit. on p. 31).



Miller, I et al. (2014). *John E. Freund's Mathematical Statistics with Applications*. 8th ed. Pearson Education Limited (cit. on pp. 8, 9).



Simpson, EH (1951). “The Interpretation of Interaction in Contingency Tables”. In: *Journal of the Royal Statistical Society. Series B (Methodological)* 13.2, pp. 238–241. JSTOR: 2984065 (cit. on p. 26).



Waa, J van der et al. (2018). “Contrastive Explanations with Local Foil Trees”. In: *2018 Workshop on Human Interpretability in Machine Learning (WHI)*. (July 14, 2018). Stockholm, Sweden. arXiv: <http://arxiv.org/abs/1806.07470v1> [stat.ML] (cit. on p. 14).