

# CS189/CS289A

## Introduction to Machine Learning

### Lecture 9: Regression

Peter Bartlett

February 17, 2015



- Review: Decision theory.

# Outline

- Review: Decision theory.
- Empirical risk minimization.

# Outline

- Review: Decision theory.
- Empirical risk minimization.
  - Least squares.

- Review: Decision theory.
- Empirical risk minimization.
  - Least squares.
  - Normal equations.

- Review: Decision theory.
- Empirical risk minimization.
  - Least squares.
  - Normal equations.
- Linear model with additive Gaussian noise.

- Review: Decision theory.
- Empirical risk minimization.
  - Least squares.
  - Normal equations.
- Linear model with additive Gaussian noise.
  - Maximum likelihood is least squares.



- Review: Decision theory.
- Empirical risk minimization.
  - Least squares.
  - Normal equations.
- Linear model with additive Gaussian noise.
  - Maximum likelihood is least squares.
  - Distributions of parameter estimates.

- **Review: Decision theory.**
- Empirical risk minimization.
  - Least squares.
  - Normal equations.
- Linear model with additive Gaussian noise.
  - Maximum likelihood is least squares.
  - Distributions of parameter estimates.

# Review: Quadratic loss

## Regression with quadratic loss

Outcomes are in  $\mathcal{Y} = \mathbb{R}$ .

We consider the quadratic loss function,  $\ell(\hat{y}, y) = (\hat{y} - y)^2$ .

Risk is expected squared error:

$$R(f) = \mathbb{E}\ell(f(X), Y) = \mathbb{E}(f(X) - Y)^2.$$

# Review: Quadratic loss


## Regression with quadratic loss

Outcomes are in  $\mathcal{Y} = \mathbb{R}$ .

We consider the quadratic loss function,  $\ell(\hat{y}, y) = (\hat{y} - y)^2$ .

Risk is expected squared error:

$$R(f) = \mathbb{E}\ell(f(X), Y) = \mathbb{E}(f(X) - Y)^2.$$



## Review: Quadratic loss

Risk:

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2$$

## Review: Quadratic loss

Risk:

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2 = \mathbb{E} \mathbb{E} [(f(X) - Y)^2 | X] .$$

# Review: Quadratic loss

Risk:

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2 = \mathbb{E} \mathbb{E} [(f(X) - Y)^2 | X] .$$

For each  $X$ , we minimize the conditional expectation of the loss,

$$\mathbb{E} [(f(X) - Y)^2 | X] .$$

# Review: Quadratic loss

Risk:

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2 = \mathbb{E} \mathbb{E} [(f(X) - Y)^2 | X] .$$

For each  $X$ , we minimize the conditional expectation of the loss,

$$\mathbb{E} [(f(X) - Y)^2 | X] .$$

The minimizer is  $f^*(X) = \mathbb{E}[Y|X]$ .



# Review: Quadratic loss

## Bias-variance decomposition

$$R(f) = \underbrace{\mathbb{E} \left[ (f(X) - \mathbb{E}[Y|X])^2 \right]}_{\text{bias}^2} + \underbrace{\mathbb{E} \left[ (\mathbb{E}[Y|X] - Y)^2 \right]}_{\text{variance}}$$

The minimizer is  $f^*(X) = \mathbb{E}[Y|X]$ .

# Review: Quadratic loss

## Bias-variance decomposition

$$\begin{aligned} R(f) &= \underbrace{\mathbb{E} \left[ (f(X) - \mathbb{E}[Y|X])^2 \right]}_{\text{bias}^2} + \underbrace{\mathbb{E} \left[ (\mathbb{E}[Y|X] - Y)^2 \right]}_{\text{variance}} \\ &= \mathbb{E} \left[ (f(X) - f^*(X))^2 \right] + \mathbb{E} \left[ (f^*(X) - Y)^2 \right] \end{aligned}$$

The minimizer is  $f^*(X) = \mathbb{E}[Y|X]$ .

# Review: Quadratic loss

## Bias-variance decomposition

$$\begin{aligned} R(f) &= \underbrace{\mathbb{E} \left[ (f(X) - \mathbb{E}[Y|X])^2 \right]}_{\text{bias}^2} + \underbrace{\mathbb{E} \left[ (\mathbb{E}[Y|X] - Y)^2 \right]}_{\text{variance}} \\ &= \mathbb{E} \left[ (f(X) - f^*(X))^2 \right] + \mathbb{E} \left[ (f^*(X) - Y)^2 \right] \\ &= \mathbb{E} \left[ (f(X) - f^*(X))^2 \right] + R(f^*). \end{aligned}$$

The minimizer is  $f^*(X) = \mathbb{E}[Y|X]$ .

# Review: Quadratic loss

## Bias-variance decomposition

$$\begin{aligned} R(f) &= \underbrace{\mathbb{E} \left[ (f(X) - \mathbb{E}[Y|X])^2 \right]}_{\text{bias}^2} + \underbrace{\mathbb{E} \left[ (\mathbb{E}[Y|X] - Y)^2 \right]}_{\text{variance}} \\ &= \mathbb{E} \left[ (f(X) - f^*(X))^2 \right] + \mathbb{E} \left[ (f^*(X) - Y)^2 \right] \\ &= \mathbb{E} \left[ (f(X) - f^*(X))^2 \right] + R(f^*). \end{aligned}$$

$$R(f) - R^* = \mathbb{E} \left[ (f(X) - f^*(X))^2 \right].$$

The minimizer is  $f^*(X) = \mathbb{E}[Y|X]$ .

# Linear regression

Consider  $X \in \mathbb{R}^p$ ,  $Y \in \mathbb{R}$ , and consider linear (affine) prediction rules

# Linear regression

Consider  $X \in \mathbb{R}^p$ ,  $Y \in \mathbb{R}$ , and consider linear (affine) prediction rules,

$$F_{lin} := \{x \mapsto x'\beta + \beta_0 : \beta \in \mathbb{R}^p, \beta_0 \in \mathbb{R}\}.$$

# Linear regression

Consider  $X \in \mathbb{R}^p$ ,  $Y \in \mathbb{R}$ , and consider linear (affine) prediction rules,

$$F_{lin} := \{x \mapsto x'\beta + \beta_0 : \beta \in \mathbb{R}^p, \beta_0 \in \mathbb{R}\}.$$

Two ways to motivate least squares:

# Linear regression

Consider  $X \in \mathbb{R}^p$ ,  $Y \in \mathbb{R}$ , and consider linear (affine) prediction rules,

$$F_{lin} := \{x \mapsto x'\beta + \beta_0 : \beta \in \mathbb{R}^p, \beta_0 \in \mathbb{R}\}.$$

Two ways to motivate least squares:

- 1 Consider the class of linear prediction rules. Minimize *empirical risk* over the class of linear prediction rules.



# Linear regression

Consider  $X \in \mathbb{R}^p$ ,  $Y \in \mathbb{R}$ , and consider linear (affine) prediction rules,

$$F_{lin} := \{x \mapsto x'\beta + \beta_0 : \beta \in \mathbb{R}^p, \beta_0 \in \mathbb{R}\}.$$

Two ways to motivate least squares:

- 1 Consider the class of linear prediction rules. Minimize *empirical risk* over the class of linear prediction rules.
- 2 Model the process generating the  $Y_i$ s as a linear function of the  $X_i$ s, plus additive Gaussian noise.  
Compute the maximum likelihood estimate for the linear coefficients.

# Linear regression

Consider  $X \in \mathbb{R}^p$ ,  $Y \in \mathbb{R}$ , and consider linear (affine) prediction rules,

$$F_{lin} := \{x \mapsto x'\beta + \beta_0 : \beta \in \mathbb{R}^p, \beta_0 \in \mathbb{R}\}.$$

Two ways to motivate least squares:

- 1 Consider the class of linear prediction rules. Minimize *empirical risk* over the class of linear prediction rules.
- 2 Model the process generating the  $Y_i$ s as a linear function of the  $X_i$ s, plus additive Gaussian noise.  
Compute the maximum likelihood estimate for the linear coefficients.

In both cases, we arrive at the *normal equations*: the choice of  $\beta$  corresponds to a projection on to a linear sub-space.

- Review: Decision theory.
- **Empirical risk minimization.**
  - Least squares.
  - Normal equations.
- Linear model with additive Gaussian noise.
  - Maximum likelihood is least squares.
  - Distributions of parameter estimates.

## Risk and empirical risk

Risk is the expected squared error:

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2.$$

## Risk and empirical risk

Risk is the expected squared error:

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2.$$

*Empirical risk* is the sample average of squared error:

## Risk and empirical risk

Risk is the expected squared error:

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2.$$

*Empirical risk* is the sample average of squared error:

$$\hat{R}(f) = \hat{\mathbb{E}}_n \ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

# Linear regression: Least squares

## Risk and empirical risk

Risk is the expected squared error:

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2.$$

*Empirical risk* is the sample average of squared error:

$$\hat{R}(f) = \hat{\mathbb{E}}_n \ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

Here,  $\hat{\mathbb{E}}_n$  means expectation under the *empirical distribution*, which puts mass  $1/n$  at each  $(X_i, Y_i)$  pair in the sample.

# Linear regression: Least squares

We want to choose a linear prediction rule  $f \in F_{lin}$  to minimize risk.



# Linear regression: Least squares

We want to choose a linear prediction rule  $f \in F_{lin}$  to minimize risk. One approach is to choose the linear prediction rule that minimizes empirical risk:


$$\hat{f} := \arg \min_{f \in F_{lin}} \hat{\mathbb{E}}_n \ell(f(X), Y)$$

# Linear regression: Least squares

We want to choose a linear prediction rule  $f \in F_{lin}$  to minimize risk. One approach is to choose the linear prediction rule that minimizes empirical risk:

$$\begin{aligned}\hat{f} &:= \arg \min_{f \in F_{lin}} \hat{\mathbb{E}}_n \ell(f(X), Y) \\ &= \arg \min_{f \in F_{lin}} \sum_{i=1}^n (f(X_i) - Y_i)^2.\end{aligned}$$

# Linear regression: Least squares



**FIGURE 3.1.** *Linear least squares fitting with  $X \in \mathbb{R}^2$ . We seek the linear function of  $X$  that minimizes the sum of squared residuals from  $Y$ .*

# Linear regression: Least squares

Just as we did when we were considering linear classifiers, we'll simplify notation by bundling the offset term ( $\beta_0$ ) into the parameter vector  $\beta$  and assuming that the covariates  $X_i$  include a constant 1 component.

# Linear regression: Least squares

Just as we did when we were considering linear classifiers, we'll simplify notation by bundling the offset term ( $\beta_0$ ) into the parameter vector  $\beta$  and assuming that the covariates  $X_i$  include a constant 1 component. Then  $f \in F_{lin}$  is of the form  $f(x) = x'\beta$ .

# Linear regression: Least squares

We wish to find  $\hat{f} : x \mapsto x' \hat{\beta}$

# Linear regression: Least squares

We wish to find  $\hat{f} : x \mapsto x' \hat{\beta}$ , where

$$\hat{\beta} = \arg \min_{\beta \in \mathbb{R}^p} \sum_{i=1}^n (X_i' \beta - Y_i)^2$$

# Linear regression: Least squares

We wish to find  $\hat{f} : x \mapsto x' \hat{\beta}$ , where

$$\begin{aligned}\hat{\beta} &= \arg \min_{\beta \in \mathbb{R}^p} \sum_{i=1}^n (X_i' \beta - Y_i)^2 \\ &= \arg \min_{\beta \in \mathbb{R}^p} \underbrace{\|X\beta - y\|^2}_{\text{RSS}},\end{aligned}$$



where the *design matrix*  $X \in \mathbb{R}^{n \times p}$  and response vector  $y \in \mathbb{R}^n$  are

$$X = \begin{pmatrix} X_1' \\ X_2' \\ \vdots \\ X_n' \end{pmatrix}, \quad y = \begin{pmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{pmatrix}.$$

(Think of  $n \gg p$ , so  $X$  is tall.)



# Linear regression: Least squares



# Linear regression: Least squares

Defining

$$RSS(\beta) = \frac{1}{2} \|X\beta - y\|^2$$

# Linear regression: Least squares

Defining

$$\begin{aligned}RSS(\beta) &= \frac{1}{2} \|X\beta - y\|^2 \\ &= \frac{1}{2} (X\beta - y)' (X\beta - y)\end{aligned}$$

# Linear regression: Least squares

Defining

$$\begin{aligned}RSS(\beta) &= \frac{1}{2} \|X\beta - y\|^2 \\&= \frac{1}{2} (X\beta - y)' (X\beta - y) \\&= \frac{1}{2} \beta' X' X \beta - y' X \beta + \frac{1}{2} y' y,\end{aligned}$$

# Linear regression: Least squares

Defining

$$\begin{aligned}RSS(\beta) &= \frac{1}{2} \|X\beta - y\|^2 \\&= \frac{1}{2} (X\beta - y)' (X\beta - y) \\&= \frac{1}{2} \beta' X' X \beta - y' X \beta + \frac{1}{2} y' y,\end{aligned}$$

we can differentiate wrt  $\beta$ :

$$\nabla_{\beta} RSS(\beta) = X' X \beta - X' y, \quad \nabla_{\beta}^2 RSS(\beta) = X' X.$$

# Linear regression: Least squares

Defining

$$\begin{aligned}RSS(\beta) &= \frac{1}{2} \|X\beta - y\|^2 \\&= \frac{1}{2} (X\beta - y)' (X\beta - y) \\&= \frac{1}{2} \beta' X' X \beta - y' X \beta + \frac{1}{2} y' y,\end{aligned}$$

we can differentiate wrt  $\beta$ :

$$\nabla_{\beta} RSS(\beta) = X' X \beta - X' y, \quad \nabla_{\beta}^2 RSS(\beta) = X' X.$$

Now,  $X' X \succeq 0$

# Linear regression: Least squares

Defining

$$\begin{aligned}RSS(\beta) &= \frac{1}{2} \|X\beta - y\|^2 \\&= \frac{1}{2} (X\beta - y)' (X\beta - y) \\&= \frac{1}{2} \beta' X' X \beta - y' X \beta + \frac{1}{2} y' y,\end{aligned}$$

we can differentiate wrt  $\beta$ :

$$\nabla_{\beta} RSS(\beta) = X' X \beta - X' y, \quad \nabla_{\beta}^2 RSS(\beta) = X' X.$$

Now,  $X' X \succeq 0$ , so setting  $\nabla_{\beta} RSS(\beta) = 0$  gives a minimum of  $RSS$

# Linear regression: Least squares

Defining

$$\begin{aligned}RSS(\beta) &= \frac{1}{2} \|X\beta - y\|^2 \\&= \frac{1}{2} (X\beta - y)' (X\beta - y) \\&= \frac{1}{2} \beta' X' X \beta - y' X \beta + \frac{1}{2} y' y,\end{aligned}$$

we can differentiate wrt  $\beta$ :

$$\nabla_{\beta} RSS(\beta) = X' X \beta - X' y, \quad \nabla_{\beta}^2 RSS(\beta) = X' X.$$

Now,  $X' X \succeq 0$ , so setting  $\nabla_{\beta} RSS(\beta) = 0$  gives a minimum of  $RSS$  when

$$X' X \beta = X' y.$$



# Linear regression: Least squares

## Normal equations

$$X'X\beta = X'y.$$

## Normal equations

$$X'X\beta = X'y.$$

$$\hat{\beta} = (X'X)^{-1}X'y.$$

# Linear regression: Least squares

## A projection viewpoint

We are aiming to find  $\beta$  to minimize  $\|y - X\beta\|$ .

# Linear regression: Least squares

## A projection viewpoint

We are aiming to find  $\beta$  to minimize  $\|y - X\beta\|$ .

Writing

$$X = \begin{pmatrix} x_1 & x_2 & \cdots & x_p \end{pmatrix},$$

# Linear regression: Least squares

## A projection viewpoint

We are aiming to find  $\beta$  to minimize  $\|y - X\beta\|$ .

Writing

$$X = (x_1 \quad x_2 \quad \cdots \quad x_p),$$

we have

$$y - X\beta = y - \sum_{j=1}^p \beta_j x_j.$$

# Linear regression: Least squares

## A projection viewpoint

We are aiming to find  $\beta$  to minimize  $\|y - X\beta\|$ .

Writing


$$X = (x_1 \quad x_2 \quad \cdots \quad x_p),$$

we have

$$y - X\beta = y - \sum_{j=1}^p \beta_j x_j.$$

That is, we want to find a linear combination of the columns  $x_j \in \mathbb{R}^n$  of  $X$  that minimizes Euclidean distance to  $y \in \mathbb{R}^n$ .

# Linear regression: Least squares



**FIGURE 3.2.** The  $N$ -dimensional geometry of least squares regression with two predictors. The outcome vector  $\mathbf{y}$  is orthogonally projected onto the hyperplane spanned by the input vectors  $\mathbf{x}_1$  and  $\mathbf{x}_2$ . The projection  $\hat{\mathbf{y}}$  represents the vector of the least squares predictions

# Linear regression: Least squares

## Projection Theorem

The optimal approximation  $\hat{y}$  in the space spanned by the columns  $x_j$  of  $X$  has an error  $y - \hat{y}$  that is orthogonal to that column space.



# Linear regression: Least squares

## Projection Theorem

The optimal approximation  $\hat{y}$  in the space spanned by the columns  $x_j$  of  $X$  has an error  $y - \hat{y}$  that is orthogonal to that column space.

That is,

$$(y - \hat{y})'X = 0 \quad \Leftrightarrow \quad X'(y - X\beta) = 0 \quad \Leftrightarrow \quad X'y = X'X\beta.$$

# Linear regression: Least squares

## Projection Theorem

The optimal approximation  $\hat{y}$  in the space spanned by the columns  $x_j$  of  $X$  has an error  $y - \hat{y}$  that is orthogonal to that column space.

That is,

$$(y - \hat{y})'X = 0 \quad \Leftrightarrow \quad X'(y - X\beta) = 0 \quad \Leftrightarrow \quad X'y = X'X\beta.$$

## Normal equations

$$X'X\beta = X'y.$$

# Linear regression: Least squares

## Projection Theorem

The optimal approximation  $\hat{y}$  in the space spanned by the columns  $x_j$  of  $X$  has an error  $y - \hat{y}$  that is orthogonal to that column space.

That is,

$$(y - \hat{y})'X = 0 \quad \Leftrightarrow \quad X'(y - X\beta) = 0 \quad \Leftrightarrow \quad X'y = X'X\beta.$$

## Normal equations

$$X'X\beta = X'y.$$

$$\hat{\beta} = (X'X)^{-1}X'y.$$

# Linear regression: Least squares

## Risk versus empirical risk

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2,$$

# Linear regression: Least squares

## Risk versus empirical risk

$$R(f) = \mathbb{E}\ell(f(X), Y) = \mathbb{E}(f(X) - Y)^2,$$

$$\hat{R}(f) = \hat{\mathbb{E}}_n\ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

# Linear regression: Least squares

## Risk versus empirical risk

$$R(f) = \mathbb{E}\ell(f(X), Y) = \mathbb{E}(f(X) - Y)^2,$$

$$\hat{R}(f) = \hat{\mathbb{E}}_n\ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

When is the risk of the empirical risk minimizer  $\hat{f}$  close to the minimal risk?

# Linear regression: Least squares

## Risk versus empirical risk

$$R(f) = \mathbb{E}\ell(f(X), Y) = \mathbb{E}(f(X) - Y)^2,$$

$$\hat{R}(f) = \hat{\mathbb{E}}_n\ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

When is the risk of the empirical risk minimizer  $\hat{f}$  close to the minimal risk?

It suffices if

# Linear regression: Least squares

## Risk versus empirical risk

$$R(f) = \mathbb{E}\ell(f(X), Y) = \mathbb{E}(f(X) - Y)^2,$$

$$\hat{R}(f) = \hat{\mathbb{E}}_n\ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

When is the risk of the empirical risk minimizer  $\hat{f}$  close to the minimal risk?

It suffices if

- 1 the  $X$  come from a compact set,



# Linear regression: Least squares

## Risk versus empirical risk

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2,$$

$$\hat{R}(f) = \hat{\mathbb{E}}_n \ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

When is the risk of the empirical risk minimizer  $\hat{f}$  close to the minimal risk?

It suffices if

- 1 the  $X$  come from a compact set,
- 2 the  $Y_i$ s have tails that are not too heavy (e.g., sub-Gaussian),

# Linear regression: Least squares

## Risk versus empirical risk

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2,$$

$$\hat{R}(f) = \hat{\mathbb{E}}_n \ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

When is the risk of the empirical risk minimizer  $\hat{f}$  close to the minimal risk?

It suffices if

- 1 the  $X$  come from a compact set,
- 2 the  $Y_i$ s have tails that are not too heavy (e.g., sub-Gaussian),
- 3  $\|\hat{\theta}\|$  is not too large, and

# Linear regression: Least squares

## Risk versus empirical risk

$$R(f) = \mathbb{E} \ell(f(X), Y) = \mathbb{E} (f(X) - Y)^2,$$

$$\hat{R}(f) = \hat{\mathbb{E}}_n \ell(f(X), Y) = \frac{1}{n} \sum_{i=1}^n (f(X_i) - Y_i)^2.$$

When is the risk of the empirical risk minimizer  $\hat{f}$  close to the minimal risk?

It suffices if

- 1 the  $X$  come from a compact set,
- 2 the  $Y_i$ s have tails that are not too heavy (e.g., sub-Gaussian),
- 3  $\|\hat{\theta}\|$  is not too large, and
- 4  $n \gg p$ .

- Review: Decision theory.
- Empirical risk minimization.
  - Least squares.
  - Normal equations.
- **Linear model with additive Gaussian noise.**
  - Maximum likelihood is least squares.
  - Distributions of parameter estimates.

# Linear regression: probability models

## Linear model

# Linear regression: probability models

## Linear model

Model the conditional distribution of  $Y$  given  $X = x$  as

$$P(Y|X = x) = \mathcal{N}(x'\beta, \sigma^2).$$

# Linear regression: probability models

## Linear model

Model the conditional distribution of  $Y$  given  $X = x$  as

$$P(Y|X = x) = \mathcal{N}(x'\beta, \sigma^2).$$

Equivalently:  $Y = x'\beta + \epsilon$ , where  $\epsilon \sim \mathcal{N}(0, \sigma^2)$ .


# Linear regression: probability models

## Linear model

Model the conditional distribution of  $Y$  given  $X = x$  as

$$P(Y|X = x) = \mathcal{N}(x'\beta, \sigma^2).$$

Equivalently:  $Y = x'\beta + \epsilon$ , where  $\epsilon \sim \mathcal{N}(0, \sigma^2)$ .





# Linear models

How to estimate  $\beta$ ?

# Linear models

How to estimate  $\beta$ ?

Maximum likelihood

# Linear models

How to estimate  $\beta$ ?

Maximum likelihood

Conditional likelihood:

$$L(\beta) = \prod_{i=1}^n p(Y_i|X_i, \beta) = \frac{1}{(2\pi\sigma^2)^{n/2}} \exp \left( -\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - X_i'\beta)^2 \right).$$

# Linear models

How to estimate  $\beta$ ?

## Maximum likelihood

Conditional likelihood:

$$L(\beta) = \prod_{i=1}^n p(Y_i|X_i, \beta) = \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - X_i'\beta)^2\right).$$

Log likelihood:

$$\ell(\beta) = (\text{function of } \sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - X_i'\beta)^2.$$

# Linear models

How to estimate  $\beta$ ?

## Maximum likelihood

Conditional likelihood:

$$L(\beta) = \prod_{i=1}^n p(Y_i|X_i, \beta) = \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - X_i'\beta)^2\right).$$

Log likelihood:

$$\ell(\beta) = (\text{function of } \sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - X_i'\beta)^2.$$

Maximum likelihood is least squares.

## Bias and variance of $\hat{\beta}$

## Bias and variance of $\hat{\beta}$

Fix  $X$ .

# Linear models

## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,



## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\mathbb{E}\hat{\beta}$$

## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\mathbb{E}\hat{\beta} = \mathbb{E}[(X'X)^{-1}X'y]$$

## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\begin{aligned}\mathbb{E}\hat{\beta} &= \mathbb{E}[(X'X)^{-1}X'y] \\ &= (X'X)^{-1}X'\mathbb{E}y\end{aligned}$$

## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\begin{aligned}\mathbb{E}\hat{\beta} &= \mathbb{E}[(X'X)^{-1}X'y] \\ &= (X'X)^{-1}X'\mathbb{E}y \\ &= (X'X)^{-1}X'X\beta\end{aligned}$$

## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\begin{aligned}\mathbb{E}\hat{\beta} &= \mathbb{E}[(X'X)^{-1}X'y] \\ &= (X'X)^{-1}X'\mathbb{E}y \\ &= (X'X)^{-1}X'X\beta \\ &= \beta.\end{aligned}$$

## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\begin{aligned}\mathbb{E}\hat{\beta} &= \mathbb{E}[(X'X)^{-1}X'y] \\ &= (X'X)^{-1}X'\mathbb{E}y \\ &= (X'X)^{-1}X'X\beta \\ &= \beta.\end{aligned}$$

$$\text{Cov}(\hat{\beta})$$

## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\begin{aligned}\mathbb{E}\hat{\beta} &= \mathbb{E}[(X'X)^{-1}X'y] \\ &= (X'X)^{-1}X'\mathbb{E}y \\ &= (X'X)^{-1}X'X\beta \\ &= \beta.\end{aligned}$$

$$\text{Cov}(\hat{\beta}) = \mathbb{E}[(\hat{\beta} - \beta)(\hat{\beta} - \beta)']$$

## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\begin{aligned}\mathbb{E}\hat{\beta} &= \mathbb{E}[(X'X)^{-1}X'y] \\ &= (X'X)^{-1}X'\mathbb{E}y \\ &= (X'X)^{-1}X'X\beta \\ &= \beta.\end{aligned}$$

$$\begin{aligned}\text{Cov}(\hat{\beta}) &= \mathbb{E}\left[(\hat{\beta} - \beta)(\hat{\beta} - \beta)'\right] \\ &= \mathbb{E}\left[((X'X)^{-1}X'y - \beta)((X'X)^{-1}X'y - \beta)'\right]\end{aligned}$$



## Bias and variance of $\hat{\beta}$

Fix  $X$ . Provided  $\mathbb{E}y = X\beta$  and  $\text{Cov}(y) = \sigma^2 I$ ,

$$\begin{aligned}\mathbb{E}\hat{\beta} &= \mathbb{E}[(X'X)^{-1}X'y] \\ &= (X'X)^{-1}X'\mathbb{E}y \\ &= (X'X)^{-1}X'X\beta \\ &= \beta.\end{aligned}$$

$$\begin{aligned}\text{Cov}(\hat{\beta}) &= \mathbb{E}[(\hat{\beta} - \beta)(\hat{\beta} - \beta)'] \\ &= \mathbb{E}\left[((X'X)^{-1}X'y - \beta)((X'X)^{-1}X'y - \beta)'\right] \\ &\vdots \\ &= \sigma^2(X'X)^{-1}.\end{aligned}$$

## Statistical tests

## Statistical tests

If the data is generated by a linear model with  $\mathcal{N}(0, \sigma^2)$  noise,

## Statistical tests

If the data is generated by a linear model with  $\mathcal{N}(0, \sigma^2)$  noise, then:

## Statistical tests

If the data is generated by a linear model with  $\mathcal{N}(0, \sigma^2)$  noise, then:

- We can compute distributions of parameter estimates:

$$\hat{\beta} \sim \mathcal{N}(\beta, (X'X)^{-1}\sigma^2)$$

## Statistical tests

If the data is generated by a linear model with  $\mathcal{N}(0, \sigma^2)$  noise, then:

- We can compute distributions of parameter estimates:

$$\hat{\beta} \sim \mathcal{N}(\beta, (X'X)^{-1}\sigma^2)$$

- We can calculate approximate confidence sets for the parameters: the standardized coefficient is

$$z_j = \frac{\hat{\beta}_j}{\sigma \sqrt{v_j}},$$

which is normal (here,  $v_j$  is the  $j$ th diagonal entry of  $(X'X)^{-1}$ ).

## Statistical tests

If the data is generated by a linear model with  $\mathcal{N}(0, \sigma^2)$  noise, then:

- We can compute distributions of parameter estimates:

$$\hat{\beta} \sim \mathcal{N}(\beta, (X'X)^{-1}\sigma^2)$$

- We can calculate approximate confidence sets for the parameters: the standardized coefficient is

$$z_j = \frac{\hat{\beta}_j}{\sigma\sqrt{v_j}},$$

which is normal (here,  $v_j$  is the  $j$ th diagonal entry of  $(X'X)^{-1}$ ).

- In particular, we can design tests for non-zero values of parameters.

## Statistical tests

If the data is generated by a linear model with  $\mathcal{N}(0, \sigma^2)$  noise, then:

- We can compute distributions of parameter estimates:

$$\hat{\beta} \sim \mathcal{N}(\beta, (X'X)^{-1}\sigma^2)$$

- We can calculate approximate confidence sets for the parameters: the standardized coefficient is

$$z_j = \frac{\hat{\beta}_j}{\sigma\sqrt{v_j}},$$

which is normal (here,  $v_j$  is the  $j$ th diagonal entry of  $(X'X)^{-1}$ ).

- In particular, we can design tests for non-zero values of parameters.

(see text)



- Review: Decision theory.
- Empirical risk minimization.
  - Least squares.
  - Normal equations.
- Linear model with additive Gaussian noise.
  - Maximum likelihood is least squares.
  - Distributions of parameter estimates.