

CS189/CS289A
Introduction to Machine Learning
Lecture 7: The Multivariate Normal Distribution

Peter Bartlett

February 10, 2015

- Probability density function.

Outline

- Probability density function.
- 2-D Gaussian

Outline

- Probability density function.
- 2-D Gaussian
- Covariance matrix.

- Probability density function.
- 2-D Gaussian
- Covariance matrix.
- Diagonal covariance.

- Probability density function.
- 2-D Gaussian
- Covariance matrix.
- Diagonal covariance.
- Non-diagonal covariance and diagonalization.

- Probability density function.
- 2-D Gaussian
- Covariance matrix.
- Diagonal covariance.
- Non-diagonal covariance and diagonalization.
- Properties of multivariate Gaussians.

- **Probability density function**
- 2-D Gaussian
- Covariance matrix.
- Diagonal covariance.
- Non-diagonal covariance and diagonalization.
- Properties of multivariate Gaussians.

Gaussian probability density function

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right).$$

Gaussian probability density function

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right).$$

- two parameters: $\mu \in \mathbb{R}$, $\sigma^2 \in \mathbb{R}_+$.

Gaussian probability density function

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right).$$

- two parameters: $\mu \in \mathbb{R}$, $\sigma^2 \in \mathbb{R}_+$.
- normalization: $\frac{1}{\sqrt{2\pi\sigma^2}}$.

Gaussian probability density function

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right).$$

- two parameters: $\mu \in \mathbb{R}$, $\sigma^2 \in \mathbb{R}_+$.

- normalization: $\frac{1}{\sqrt{2\pi\sigma^2}}$.

- argument of \exp : $-\frac{(x-\mu)^2}{2\sigma^2} \leq 0$.

Concave, maximum (of 0) at $x = \mu$.

Gaussian probability density function

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right).$$

- two parameters: $\mu \in \mathbb{R}$, $\sigma^2 \in \mathbb{R}_+$.
- normalization: $\frac{1}{\sqrt{2\pi\sigma^2}}$.
- argument of \exp : $-\frac{(x - \mu)^2}{2\sigma^2} \leq 0$.
Concave, maximum (of 0) at $x = \mu$.
- location parameter $\mu \in \mathbb{R}$: $(x - \mu)^2$.

Gaussian probability density function

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right).$$

- two parameters: $\mu \in \mathbb{R}$, $\sigma^2 \in \mathbb{R}_+$.

- normalization: $\frac{1}{\sqrt{2\pi\sigma^2}}$.

- argument of \exp : $-\frac{(x-\mu)^2}{2\sigma^2} \leq 0$.

Concave, maximum (of 0) at $x = \mu$.

- location parameter $\mu \in \mathbb{R}$: $(x-\mu)^2$.

- scale parameter $\sigma^2 \in \mathbb{R}_+$: $\frac{(x-\mu)^2}{2\sigma^2}$.

Gaussian probability density function

$$x \in \mathbb{R} : \quad p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right).$$

$$x \in \mathbb{R}^d : \quad p(x) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left(-\frac{1}{2}(x - \mu)' \Sigma^{-1} (x - \mu)\right).$$

Gaussian probability density function

$$x \in \mathbb{R} : \quad p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right).$$

$$x \in \mathbb{R}^d : \quad p(x) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left(-\frac{1}{2}(x - \mu)' \Sigma^{-1} (x - \mu)\right).$$

- parameters: $\mu \in \mathbb{R}^d$, $\sigma^2 \in \mathbb{R}^{d \times d}$.

Gaussian probability density function

$$x \in \mathbb{R} : \quad p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right).$$

$$x \in \mathbb{R}^d : \quad p(x) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left(-\frac{1}{2}(x - \mu)' \Sigma^{-1} (x - \mu)\right).$$

- parameters: $\mu \in \mathbb{R}^d$, $\sigma^2 \in \mathbb{R}^{d \times d}$.
- normalization: $\frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}}$

Gaussian probability density function

$$x \in \mathbb{R} : \quad p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right).$$

$$x \in \mathbb{R}^d : \quad p(x) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left(-\frac{1}{2}(x - \mu)' \Sigma^{-1} (x - \mu)\right).$$

- parameters: $\mu \in \mathbb{R}^d$, $\sigma^2 \in \mathbb{R}^{d \times d}$.
- normalization: $\frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}}$
- location parameter $\mu \in \mathbb{R}^d$: $(x - \mu)$.

Gaussian probability density function

$$x \in \mathbb{R} : \quad p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right).$$

$$x \in \mathbb{R}^d : \quad p(x) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left(-\frac{1}{2}(x - \mu)' \Sigma^{-1} (x - \mu)\right).$$

- parameters: $\mu \in \mathbb{R}^d$, $\sigma^2 \in \mathbb{R}^{d \times d}$.
- normalization: $\frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}}$
- location parameter $\mu \in \mathbb{R}^d$: $(x - \mu)$.
- scale parameter $\Sigma \in \mathbf{S}_+^d$: $(x - \mu)' \Sigma^{-1} (x - \mu)$.


Gaussian probability density function

$$x \in \mathbb{R} : \quad p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right).$$


$$x \in \mathbb{R}^d : \quad p(x) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left(-\frac{1}{2}(x - \mu)' \Sigma^{-1} (x - \mu)\right).$$

- parameters: $\mu \in \mathbb{R}^d$, $\sigma^2 \in \mathbb{R}^{d \times d}$.
- normalization: $\frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}}$
- location parameter $\mu \in \mathbb{R}^d$: $(x - \mu)$.
- scale parameter $\Sigma \in \mathbf{S}_+^d$: $(x - \mu)' \Sigma^{-1} (x - \mu)$.
- argument of \exp : $-\frac{1}{2}(x - \mu)' \Sigma^{-1} (x - \mu) \leq 0$.
Concave, maximum (of 0) at $x = \mu$.

Gaussian probability density function



Gaussian probability density function



Gaussian probability density function

Gaussian probability density function

- What is the mean height? mean weight?

Gaussian probability density function

- What is the mean height? mean weight?
- Where is the mean vector μ ?

Gaussian probability density function

- What is the mean height? mean weight?
- Where is the mean vector μ ?
- Are height and weight correlated?

Gaussian probability density function

- What is the mean height? mean weight?
- Where is the mean vector μ ?
- Are height and weight correlated?
- Contours are level sets of the pdf.

Gaussian probability density function

- What is the mean height? mean weight?
- Where is the mean vector μ ?
- Are height and weight correlated?
- Contours are level sets of the pdf.
- What is the pdf of the marginal distribution of height, p_h ? p_w ?

Gaussian probability density function

- What is the mean height? mean weight?
- Where is the mean vector μ ?
- Are height and weight correlated?
- Contours are level sets of the pdf.
- What is the pdf of the marginal distribution of height, p_h ? p_w ?
- Is the pdf $p = p_h \cdot p_w$?

Gaussian probability density function

- What is the mean height? mean weight?
- Where is the mean vector μ ?
- Are height and weight correlated?
- Contours are level sets of the pdf.
- What is the pdf of the marginal distribution of height, p_h ? p_w ?
- Is the pdf $p = p_h \cdot p_w$?
- Suppose we project height and weight in a particular direction. What is the direction of maximum variance? minimum variance?

2-D Gaussian distribution

Consider independent Gaussian random variables X and Y :

$$p_X(x) = \frac{1}{\sqrt{2\pi\sigma_X^2}} \exp\left(-\frac{(x - \mu_X)^2}{2\sigma_X^2}\right).$$

$$p_Y(y) = \frac{1}{\sqrt{2\pi\sigma_Y^2}} \exp\left(-\frac{(y - \mu_Y)^2}{2\sigma_Y^2}\right).$$

2-D Gaussian distribution

Consider independent Gaussian random variables X and Y :

$$p_X(x) = \frac{1}{\sqrt{2\pi\sigma_X^2}} \exp\left(-\frac{(x - \mu_X)^2}{2\sigma_X^2}\right).$$

$$p_Y(y) = \frac{1}{\sqrt{2\pi\sigma_Y^2}} \exp\left(-\frac{(y - \mu_Y)^2}{2\sigma_Y^2}\right).$$

What is their joint density?

2-D Gaussian distribution

Consider independent Gaussian random variables X and Y :

$$p_X(x) = \frac{1}{\sqrt{2\pi\sigma_X^2}} \exp\left(-\frac{(x - \mu_X)^2}{2\sigma_X^2}\right).$$

$$p_Y(y) = \frac{1}{\sqrt{2\pi\sigma_Y^2}} \exp\left(-\frac{(y - \mu_Y)^2}{2\sigma_Y^2}\right).$$

What is their joint density?

$$p(x, y) = p_X(x)p_Y(y)$$

$$= \frac{1}{\sqrt{(2\pi)^2\sigma_X^2\sigma_Y^2}} \exp\left(-\frac{(x - \mu_X)^2}{2\sigma_X^2} - \frac{(y - \mu_Y)^2}{2\sigma_Y^2}\right)$$


$$= \frac{1}{\sqrt{(2\pi)^2|\Sigma|}} \exp\left(-\frac{1}{2} \begin{pmatrix} x - \mu_X & y - \mu_Y \end{pmatrix} \begin{bmatrix} \sigma_X^2 & 0 \\ 0 & \sigma_Y^2 \end{bmatrix}^{-1} \begin{pmatrix} x - \mu_X \\ y - \mu_Y \end{pmatrix}\right).$$

(Check!)


2-D Gaussian distribution

$$\mu = \begin{bmatrix} 5.0 \\ 5.0 \end{bmatrix}$$

$$\Sigma = \begin{bmatrix} 0.8 & 0.0 \\ 0.0 & 0.6 \end{bmatrix}$$



2-D Gaussian distribution



Gaussian probability density function

- Are X and Y correlated?

Gaussian probability density function

- Are X and Y correlated?
- Suppose we project height and weight in a particular direction. What is the direction of maximum variance? minimum variance?

- Probability density function.
- 2-D Gaussian
- **Covariance matrix.**
- Diagonal covariance.
- Non-diagonal covariance and diagonalization.
- Properties of multivariate Gaussians.

Covariance matrices

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix}$$

Covariance matrices

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix}$$

$$\mathbb{E}X = \begin{bmatrix} \mathbb{E}X_1 \\ \mathbb{E}X_2 \\ \mathbb{E}X_3 \end{bmatrix}$$

Covariance matrices

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix}$$

$$\mathbb{E}X = \begin{bmatrix} \mathbb{E}X_1 \\ \mathbb{E}X_2 \\ \mathbb{E}X_3 \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \end{bmatrix}$$

Covariance matrices

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix}$$

$$\mathbb{E}X = \begin{bmatrix} \mathbb{E}X_1 \\ \mathbb{E}X_2 \\ \mathbb{E}X_3 \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \end{bmatrix} = \mu,$$

Covariance matrices

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix} \quad \mathbb{E}X = \begin{bmatrix} \mathbb{E}X_1 \\ \mathbb{E}X_2 \\ \mathbb{E}X_3 \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \end{bmatrix} = \mu,$$

$$\text{Cov}(X_i, X_j) = \mathbb{E}(X_i - \mu_i)(X_j - \mu_j),$$

Covariance matrices

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix} \quad \mathbb{E}X = \begin{bmatrix} \mathbb{E}X_1 \\ \mathbb{E}X_2 \\ \mathbb{E}X_3 \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \end{bmatrix} = \mu,$$

$$\text{Cov}(X_i, X_j) = \mathbb{E}(X_i - \mu_i)(X_j - \mu_j),$$

$$\Sigma = \mathbb{E}(X - \mu)(X - \mu)'$$

Covariance matrices

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix} \quad \mathbb{E}X = \begin{bmatrix} \mathbb{E}X_1 \\ \mathbb{E}X_2 \\ \mathbb{E}X_3 \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \end{bmatrix} = \mu,$$

$$\text{Cov}(X_i, X_j) = \mathbb{E}(X_i - \mu_i)(X_j - \mu_j),$$

$$\begin{aligned} \Sigma &= \mathbb{E}(X - \mu)(X - \mu)' \\ &= \begin{bmatrix} \text{Cov}(X_1, X_1) & \text{Cov}(X_1, X_2) & \text{Cov}(X_1, X_3) \\ \text{Cov}(X_2, X_1) & \text{Cov}(X_2, X_2) & \text{Cov}(X_2, X_3) \\ \text{Cov}(X_3, X_1) & \text{Cov}(X_3, X_2) & \text{Cov}(X_3, X_3) \end{bmatrix} \end{aligned}$$

Covariance matrices

$$X = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix} \quad \mathbb{E}X = \begin{bmatrix} \mathbb{E}X_1 \\ \mathbb{E}X_2 \\ \mathbb{E}X_3 \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \end{bmatrix} = \mu,$$

$$\text{Cov}(X_i, X_j) = \mathbb{E}(X_i - \mu_i)(X_j - \mu_j),$$

$$\begin{aligned} \Sigma &= \mathbb{E}(X - \mu)(X - \mu)' \\ &= \begin{bmatrix} \text{Cov}(X_1, X_1) & \text{Cov}(X_1, X_2) & \text{Cov}(X_1, X_3) \\ \text{Cov}(X_2, X_1) & \text{Cov}(X_2, X_2) & \text{Cov}(X_2, X_3) \\ \text{Cov}(X_3, X_1) & \text{Cov}(X_3, X_2) & \text{Cov}(X_3, X_3) \end{bmatrix} \\ &= \begin{bmatrix} \text{Var}(X_1) & \text{Cov}(X_1, X_2) & \text{Cov}(X_1, X_3) \\ \text{Cov}(X_2, X_1) & \text{Var}(X_2) & \text{Cov}(X_2, X_3) \\ \text{Cov}(X_3, X_1) & \text{Cov}(X_3, X_2) & \text{Var}(X_3) \end{bmatrix}. \end{aligned}$$

Properties of Σ

Properties of Σ

- Symmetric: $\Sigma_{i,j} = \Sigma_{j,i}$.

Properties of Σ

- Symmetric: $\Sigma_{i,j} = \Sigma_{j,i}$.
- Non-negative diagonal entries: $\Sigma_{i,i} \geq 0$.

Properties of Σ

- Symmetric: $\Sigma_{i,j} = \Sigma_{j,i}$.
- Non-negative diagonal entries: $\Sigma_{i,i} \geq 0$.
- Positive semidefinite: for all $v \in \mathbb{R}^d$, $v' \Sigma v \geq 0$.

Properties of Σ

- Symmetric: $\Sigma_{i,j} = \Sigma_{j,i}$.
- Non-negative diagonal entries: $\Sigma_{i,i} \geq 0$.
- Positive semidefinite: for all $v \in \mathbb{R}^d$, $v' \Sigma v \geq 0$.

What is $\mathbb{E} [(v'(X - \mu))^2]$?

- Probability density function.
- 2-D Gaussian
- Covariance matrix.
- **Diagonal covariance.**
- Non-diagonal covariance and diagonalization.
- Properties of multivariate Gaussians.

Diagonal covariance

Consider $X \sim N(\mu, \Sigma)$ where Σ is diagonal:

$$\Sigma = \begin{bmatrix} \sigma_1^2 & 0 & \cdots & 0 \\ 0 & \sigma_2^2 & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & \sigma_d^2 \end{bmatrix}.$$

Diagonal covariance

Consider $X \sim N(\mu, \Sigma)$ where Σ is diagonal:

$$\Sigma = \begin{bmatrix} \sigma_1^2 & 0 & \cdots & 0 \\ 0 & \sigma_2^2 & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & \sigma_d^2 \end{bmatrix}.$$

The components of X are uncorrelated:

$$\text{Cov}(X_i, X_j) = 0 \quad \text{for} \quad i \neq j.$$

Diagonal covariance

Consider $X \sim N(\mu, \Sigma)$ where Σ is diagonal:

$$\Sigma = \begin{bmatrix} \sigma_1^2 & 0 & \cdots & 0 \\ 0 & \sigma_2^2 & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & \sigma_d^2 \end{bmatrix}.$$

The components of X are uncorrelated:

$$\text{Cov}(X_i, X_j) = 0 \quad \text{for} \quad i \neq j.$$

Consider the super-level sets of the pdf:

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

Diagonal covariance

$$\begin{aligned}(x - \mu)' \Sigma^{-1} (x - \mu) &\leq r^2 \\ \Leftrightarrow \sum_{i=1}^d \frac{(x_i - \mu_i)^2}{\sigma_i^2} &\leq r^2.\end{aligned}$$

Diagonal covariance

$$\begin{aligned}(x - \mu)' \Sigma^{-1} (x - \mu) &\leq r^2 \\ \Leftrightarrow \sum_{i=1}^d \frac{(x_i - \mu_i)^2}{\sigma_i^2} &\leq r^2.\end{aligned}$$

Thus, the \mathcal{E}_r correspond to *axis-aligned ellipsoids* in \mathbb{R}^d .

Diagonal covariance

$$\begin{aligned}(x - \mu)' \Sigma^{-1} (x - \mu) &\leq r^2 \\ \Leftrightarrow \quad \sum_{i=1}^d \frac{(x_i - \mu_i)^2}{\sigma_i^2} &\leq r^2.\end{aligned}$$


Thus, the \mathcal{E}_r correspond to *axis-aligned ellipsoids* in \mathbb{R}^d .
For instance, if $x_2 = \mu_2, \dots, x_d = \mu_d$, then $x \in \mathcal{E}_r$ when

$$(x_1 - \mu_1)^2 \leq r^2 \sigma_1^2 \quad \Leftrightarrow \quad |x_1 - \mu_1| \leq r \sigma_1.$$


Diagonal covariance

$$\mu = \begin{bmatrix} 5.0 \\ 5.0 \end{bmatrix}$$

$$\Sigma = \begin{bmatrix} 0.8 & 0.0 \\ 0.0 & 0.6 \end{bmatrix}$$



2-D Gaussian distribution



Diagonal covariance

The volume of the ellipsoid \mathcal{E}_r is proportional to

$$\prod_{i=1}^d \sigma_i = \sqrt{|\Sigma|}.$$

(Think about the volume of the box with sides scaled by the σ_i .)

Diagonal covariance

The volume of the ellipsoid \mathcal{E}_r is proportional to

$$\prod_{i=1}^d \sigma_i = \sqrt{|\Sigma|}.$$

(Think about the volume of the box with sides scaled by the σ_i .)

- This explains the appearance of $1/\sqrt{|\Sigma|}$ in the normalization factor.

Diagonal covariance

The volume of the ellipsoid \mathcal{E}_r is proportional to

$$\prod_{i=1}^d \sigma_i = \sqrt{|\Sigma|}.$$

(Think about the volume of the box with sides scaled by the σ_i .)

- This explains the appearance of $1/\sqrt{|\Sigma|}$ in the normalization factor.
- It also shows that, for a fixed r , $\Pr(\mathcal{E}_r)$ is the same, for all μ and Σ (but depends on d).

- Probability density function.
- 2-D Gaussian
- Covariance matrix.
- Diagonal covariance.
- **Non-diagonal covariance and diagonalization.**
- Properties of multivariate Gaussians.

Non-diagonal covariance

What if Σ is not diagonal?

Non-diagonal covariance

What if Σ is not diagonal?

Suppose we start with a diagonal Σ (so the contours of the pdf are axis-aligned ellipsoids)

Non-diagonal covariance

What if Σ is not diagonal?

Suppose we start with a diagonal Σ (so the contours of the pdf are axis-aligned ellipsoids), and then consider the transformed vector

$$Y = \mu_X + Q(X - \mu_X),$$

where Q is a matrix that corresponds to a rotation about the origin.

Non-diagonal covariance

What if Σ is not diagonal?

Suppose we start with a diagonal Σ (so the contours of the pdf are axis-aligned ellipsoids), and then consider the transformed vector

$$Y = \mu_X + Q(X - \mu_X),$$

where Q is a matrix that corresponds to a rotation about the origin.

e.g. $Q = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$

Non-diagonal covariance

This leaves the mean unchanged:

$$\mathbb{E}Y = \mu_X + \mathbb{E}[Q(X - \mu_X)] = \mu_X + Q(\mathbb{E}[X] - \mu_X) = \mu_X.$$

Non-diagonal covariance

This leaves the mean unchanged:

$$\mathbb{E}Y = \mu_X + \mathbb{E}[Q(X - \mu_X)] = \mu_X + Q(\mathbb{E}[X] - \mu_X) = \mu_X.$$

But the covariance is now transformed

Non-diagonal covariance

This leaves the mean unchanged:

$$\mathbb{E}Y = \mu_X + \mathbb{E}[Q(X - \mu_X)] = \mu_X + Q(\mathbb{E}[X] - \mu_X) = \mu_X.$$

But the covariance is now transformed:

$$\Sigma_Y = \mathbb{E}(Y - \mu_X)(Y - \mu_X)'$$

Non-diagonal covariance

This leaves the mean unchanged:

$$\mathbb{E}Y = \mu_X + \mathbb{E}[Q(X - \mu_X)] = \mu_X + Q(\mathbb{E}[X] - \mu_X) = \mu_X.$$

But the covariance is now transformed:

$$\begin{aligned}\Sigma_Y &= \mathbb{E}(Y - \mu_X)(Y - \mu_X)' \\ &= \mathbb{E}[Q(X - \mu_X)(X - \mu_X)'Q']\end{aligned}$$

Non-diagonal covariance

This leaves the mean unchanged:

$$\mathbb{E}Y = \mu_X + \mathbb{E}[Q(X - \mu_X)] = \mu_X + Q(\mathbb{E}[X] - \mu_X) = \mu_X.$$

But the covariance is now transformed:

$$\begin{aligned}\Sigma_Y &= \mathbb{E}(Y - \mu_X)(Y - \mu_X)' \\ &= \mathbb{E}[Q(X - \mu_X)(X - \mu_X)'Q'] \\ &= Q\mathbb{E}[(X - \mu_X)(X - \mu_X)']Q'\end{aligned}$$

Non-diagonal covariance

This leaves the mean unchanged:

$$\mathbb{E}Y = \mu_X + \mathbb{E}[Q(X - \mu_X)] = \mu_X + Q(\mathbb{E}[X] - \mu_X) = \mu_X.$$

But the covariance is now transformed:

$$\begin{aligned}\Sigma_Y &= \mathbb{E}(Y - \mu_X)(Y - \mu_X)' \\ &= \mathbb{E}[Q(X - \mu_X)(X - \mu_X)'Q'] \\ &= Q\mathbb{E}[(X - \mu_X)(X - \mu_X)']Q' \\ &= Q\Sigma_XQ' .\end{aligned}$$

Non-diagonal covariance

This leaves the mean unchanged:





$$\mathbb{E}Y = \mu_X + \mathbb{E}[Q(X - \mu_X)] = \mu_X + Q(\mathbb{E}[X] - \mu_X) = \mu_X.$$

But the covariance is now transformed:

$$\begin{aligned}\Sigma_Y &= \mathbb{E}(Y - \mu_X)(Y - \mu_X)' \\ &= \mathbb{E}[Q(X - \mu_X)(X - \mu_X)'Q'] \\ &= Q\mathbb{E}[(X - \mu_X)(X - \mu_X)']Q' \\ &= Q\Sigma_XQ' .\end{aligned}$$

It is no longer diagonal.

Non-diagonal covariance



Non-diagonal covariance

We can always view a covariance matrix in this way: as a rotated version of a diagonal covariance.

Non-diagonal covariance

We can always view a covariance matrix in this way: as a rotated version of a diagonal covariance.

Eigenvalues and eigenvectors

For a square matrix A , λ is an *eigenvalue* and x is an *eigenvector* if

$$Ax = \lambda x.$$

Non-diagonal covariance

We can always view a covariance matrix in this way: as a rotated version of a diagonal covariance.

Eigenvalues and eigenvectors

For a square matrix A , λ is an *eigenvalue* and x is an *eigenvector* if

$$Ax = \lambda x.$$

That is, the transformation A corresponds to a rescaling in the x direction.

Non-diagonal covariance

We can always view a covariance matrix in this way: as a rotated version of a diagonal covariance.

Eigenvalues and eigenvectors

For a square matrix A , λ is an *eigenvalue* and x is an *eigenvector* if

$$Ax = \lambda x.$$

That is, the transformation A corresponds to a rescaling in the x direction.

(Eigenvalues are roots of the *characteristic polynomial*, $\det(A - \lambda I)$.)

Spectral Theorem

Spectral Theorem

For a symmetric real matrix $A \in \mathbb{R}^{n \times n}$, we can find n orthonormal eigenvectors of A (v_1, \dots, v_n), and the eigenvalues ($\lambda_1, \dots, \lambda_n$) are real.

Spectral Theorem

For a symmetric real matrix $A \in \mathbb{R}^{n \times n}$, we can find n orthonormal eigenvectors of A (v_1, \dots, v_n), and the eigenvalues ($\lambda_1, \dots, \lambda_n$) are real. Hence, we can write

$$AU = U\Lambda, \quad A = U\Lambda U',$$

where

$$U = [v_1 \quad v_2 \quad \cdots \quad v_n], \quad \Lambda = \text{diag} \left(\begin{bmatrix} \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_n \end{bmatrix} \right).$$

Non-diagonal covariance

Spectral Theorem

For a symmetric real matrix $A \in \mathbb{R}^{n \times n}$, we can find n orthonormal eigenvectors of A (v_1, \dots, v_n), and the eigenvalues ($\lambda_1, \dots, \lambda_n$) are real. Hence, we can write

$$AU = U\Lambda, \quad A = U\Lambda U',$$

where

$$U = [v_1 \quad v_2 \quad \cdots \quad v_n], \quad \Lambda = \text{diag} \left(\begin{bmatrix} \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_n \end{bmatrix} \right).$$

(Why “hence”? $U'U = I$, so $UU' = I$.)

Non-diagonal covariance

A covariance matrix Σ is symmetric, so we can write

$$\Sigma = U\Lambda U'.$$

Non-diagonal covariance

A covariance matrix Σ is symmetric, so we can write

$$\Sigma = U\Lambda U'.$$

It is also positive semidefinite, so we must have all of the $\lambda_i \geq 0$

Non-diagonal covariance

A covariance matrix Σ is symmetric, so we can write

$$\Sigma = U\Lambda U'.$$

It is also positive semidefinite, so we must have all of the $\lambda_i \geq 0$:

$$v_i' \Sigma v_i = v_i' U \Lambda U' v_i = \lambda_i \geq 0.$$

Non-diagonal covariance

A covariance matrix Σ is symmetric, so we can write

$$\Sigma = U\Lambda U'.$$

It is also positive semidefinite, so we must have all of the $\lambda_i \geq 0$:

$$v_i' \Sigma v_i = v_i' U \Lambda U' v_i = \lambda_i \geq 0.$$

Also, the *precision matrix* can be written

$$\Sigma^{-1} = U\Lambda^{-1}U'.$$

(Check!)

Non-diagonal covariance

Gaussian pdf

$$p(x) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right)$$

Non-diagonal covariance

Gaussian pdf

$$\begin{aligned} p(x) &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right) \\ &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' U \Lambda^{-1} U' (x - \mu) \right). \end{aligned}$$

Non-diagonal covariance

Gaussian pdf

$$\begin{aligned} p(x) &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right) \\ &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' U \Lambda^{-1} U' (x - \mu) \right). \end{aligned}$$

$U'(x - \mu)$ computes the representation of $x - \mu$ in the basis of the eigenvectors of Σ :

Non-diagonal covariance

Gaussian pdf

$$\begin{aligned} p(x) &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right) \\ &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' U \Lambda^{-1} U' (x - \mu) \right). \end{aligned}$$

$U'(x - \mu)$ computes the representation of $x - \mu$ in the basis of the eigenvectors of Σ :

$$U'(x - \mu) = \begin{bmatrix} v'_1 \\ v'_2 \\ \vdots \\ v'_d \end{bmatrix} (x - \mu)$$

Non-diagonal covariance

Gaussian pdf

$$\begin{aligned} p(x) &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right) \\ &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' U \Lambda^{-1} U' (x - \mu) \right). \end{aligned}$$

$U'(x - \mu)$ computes the representation of $x - \mu$ in the basis of the eigenvectors of Σ :

$$U'(x - \mu) = \begin{bmatrix} v_1' \\ v_2' \\ \vdots \\ v_d' \end{bmatrix} (x - \mu) = \begin{bmatrix} v_1'(x - \mu) \\ v_2'(x - \mu) \\ \vdots \\ v_d'(x - \mu) \end{bmatrix}.$$

Non-diagonal covariance

Gaussian pdf

$$\begin{aligned} p(x) &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right) \\ &= \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)' U \Lambda^{-1} U' (x - \mu) \right). \end{aligned}$$

$U'(x - \mu)$ computes the representation of $x - \mu$ in the basis of the eigenvectors of Σ :

$$U'(x - \mu) = \begin{bmatrix} v_1' \\ v_2' \\ \vdots \\ v_d' \end{bmatrix} (x - \mu) = \begin{bmatrix} v_1'(x - \mu) \\ v_2'(x - \mu) \\ \vdots \\ v_d'(x - \mu) \end{bmatrix}.$$

In this new basis, the covariance is the diagonal matrix Λ^{-1} .

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Level sets with non-diagonal covariance

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Level sets with non-diagonal covariance

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

The \mathcal{E}_r correspond to eigenvector-aligned ellipsoids in \mathbb{R}^d :

$$(x - \mu)' \Sigma^{-1} (x - \mu)$$

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Level sets with non-diagonal covariance

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

The \mathcal{E}_r correspond to eigenvector-aligned ellipsoids in \mathbb{R}^d :

$$(x - \mu)' \Sigma^{-1} (x - \mu) = (x - \mu)' U \Lambda^{-1} U' (x - \mu)$$

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Level sets with non-diagonal covariance

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

The \mathcal{E}_r correspond to eigenvector-aligned ellipsoids in \mathbb{R}^d :

$$(x - \mu)' \Sigma^{-1} (x - \mu) = (x - \mu)' U \Lambda^{-1} U' (x - \mu) = \sum_{i=1}^d \frac{(v_i'(x - \mu))^2}{\lambda_i^2}.$$

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Level sets with non-diagonal covariance

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

The \mathcal{E}_r correspond to eigenvector-aligned ellipsoids in \mathbb{R}^d :

$$(x - \mu)' \Sigma^{-1} (x - \mu) = (x - \mu)' U \Lambda^{-1} U' (x - \mu) = \sum_{i=1}^d \frac{(v_i'(x - \mu))^2}{\lambda_i^2}.$$

For instance, if $v_2'(x - \mu) = \dots = v_d'(x - \mu) = 0$, then $x \in \mathcal{E}_r$ when

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Level sets with non-diagonal covariance

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

The \mathcal{E}_r correspond to eigenvector-aligned ellipsoids in \mathbb{R}^d :

$$(x - \mu)' \Sigma^{-1} (x - \mu) = (x - \mu)' U \Lambda^{-1} U' (x - \mu) = \sum_{i=1}^d \frac{(v_i'(x - \mu))^2}{\lambda_i^2}.$$

For instance, if $v_2'(x - \mu) = \dots = v_d'(x - \mu) = 0$, then $x \in \mathcal{E}_r$ when

$$(v_1(x - \mu))^2 \leq r^2 \lambda_1^2$$

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Level sets with non-diagonal covariance

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

The \mathcal{E}_r correspond to eigenvector-aligned ellipsoids in \mathbb{R}^d :

$$(x - \mu)' \Sigma^{-1} (x - \mu) = (x - \mu)' U \Lambda^{-1} U' (x - \mu) = \sum_{i=1}^d \frac{(v_i'(x - \mu))^2}{\lambda_i^2}.$$

For instance, if $v_2'(x - \mu) = \dots = v_d'(x - \mu) = 0$, then $x \in \mathcal{E}_r$ when

$$(v_1(x - \mu))^2 \leq r^2 \lambda_1^2 \quad \Leftrightarrow \quad |v_1(x - \mu)| \leq r \lambda_1.$$

Non-diagonal covariance

The level sets are ellipsoids with axes aligned with the eigenvectors, and the eigenvalues determine the scaling of the ellipsoids:

Level sets with non-diagonal covariance

$$\mathcal{E}_r = \left\{ x \in \mathbb{R}^d : (x - \mu)' \Sigma^{-1} (x - \mu) \leq r^2 \right\}$$

The \mathcal{E}_r correspond to eigenvector-aligned ellipsoids in \mathbb{R}^d :





$$(x - \mu)' \Sigma^{-1} (x - \mu) = (x - \mu)' U \Lambda^{-1} U' (x - \mu) = \sum_{i=1}^d \frac{(v_i'(x - \mu))^2}{\lambda_i^2}.$$

For instance, if $v_2'(x - \mu) = \dots = v_d'(x - \mu) = 0$, then $x \in \mathcal{E}_r$ when

$$(v_1(x - \mu))^2 \leq r^2 \lambda_1^2 \quad \Leftrightarrow \quad |v_1(x - \mu)| \leq r \lambda_1.$$

This is the same as the diagonal case, but with a new coordinate system, defined by the eigenvectors.

Non-diagonal covariance



Non-diagonal covariance

The volume of the ellipsoid \mathcal{E}_r is proportional to

$$\prod_{i=1}^d \lambda_i = \sqrt{|\Lambda|}$$

Non-diagonal covariance

The volume of the ellipsoid \mathcal{E}_r is proportional to

$$\prod_{i=1}^d \lambda_i = \sqrt{|\Lambda|} = \sqrt{|U| \cdot |\Lambda| \cdot |U'|}$$

Non-diagonal covariance

The volume of the ellipsoid \mathcal{E}_r is proportional to

$$\prod_{i=1}^d \lambda_i = \sqrt{|\Lambda|} = \sqrt{|U| \cdot |\Lambda| \cdot |U'|} = \sqrt{|U \Lambda U'|}$$

Non-diagonal covariance

The volume of the ellipsoid \mathcal{E}_r is proportional to

$$\prod_{i=1}^d \lambda_i = \sqrt{|\Lambda|} = \sqrt{|U| \cdot |\Lambda| \cdot |U'|} = \sqrt{|U \Lambda U'|} = \sqrt{|\Sigma|}.$$

Non-diagonal covariance

The volume of the ellipsoid \mathcal{E}_r is proportional to

$$\prod_{i=1}^d \lambda_i = \sqrt{|\Lambda|} = \sqrt{|U| \cdot |\Lambda| \cdot |U'|} = \sqrt{|U \Lambda U'|} = \sqrt{|\Sigma|}.$$

(As in the diagonal case.)

- Probability density function.
- 2-D Gaussian
- Covariance matrix.
- Diagonal covariance.
- Non-diagonal covariance and diagonalization.
- **Properties of multivariate Gaussians.**

Properties of multivariate Gaussians

Properties of multivariate Gaussians

Probability measure

$$\int_{\mathbb{R}^d} p(x; \mu, \Sigma) dx = 1.$$

Properties of multivariate Gaussians

Probability measure

$$\int_{\mathbb{R}^d} p(x; \mu, \Sigma) dx = 1.$$

Expectation

$$\mathbb{E}[X] = \begin{bmatrix} \mathbb{E}[X_1] \\ \vdots \\ \mathbb{E}[X_d] \end{bmatrix} = \int_{\mathbb{R}^d} xp(x; \mu, \Sigma) dx = \mu.$$

Properties of multivariate Gaussians

Probability measure

$$\int_{\mathbb{R}^d} p(x; \mu, \Sigma) dx = 1.$$

Expectation

$$\mathbb{E}[X] = \begin{bmatrix} \mathbb{E}[X_1] \\ \vdots \\ \mathbb{E}[X_d] \end{bmatrix} = \int_{\mathbb{R}^d} x p(x; \mu, \Sigma) dx = \mu.$$

Covariance

$$\text{Cov}[X] = \mathbb{E}[(X - \mu)(X - \mu)'] = \int_{\mathbb{R}^d} (x - \mu)(x - \mu)' p(x; \mu, \Sigma) dx = \Sigma.$$

Properties of multivariate Gaussians

Sums of independent Gaussians

For independent d -dimensional Gaussians,
 $X \sim \mathcal{N}(\mu_X, \Sigma_X)$ and $Y \sim \mathcal{N}(\mu_Y, \Sigma_Y)$,

Properties of multivariate Gaussians

Sums of independent Gaussians

For independent d -dimensional Gaussians,

$$X \sim \mathcal{N}(\mu_X, \Sigma_X) \text{ and } Y \sim \mathcal{N}(\mu_Y, \Sigma_Y),$$

$$X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \Sigma_X + \Sigma_Y).$$

- NB: Adding the random variables, not the densities!
This corresponds to *convolving* the densities.
The family of Gaussian densities is closed under convolutions.

Properties of multivariate Gaussians

Sums of independent Gaussians

For independent d -dimensional Gaussians,

$$X \sim \mathcal{N}(\mu_X, \Sigma_X) \text{ and } Y \sim \mathcal{N}(\mu_Y, \Sigma_Y),$$

$$X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \Sigma_X + \Sigma_Y).$$

- NB: Adding the random variables, not the densities!
This corresponds to *convolving* the densities.
The family of Gaussian densities is closed under convolutions.
- If we accept that $X + Y$ is Gaussian, let's calculate its mean and covariance.

Properties of multivariate Gaussians

Properties of multivariate Gaussians

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$.

Properties of multivariate Gaussians

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$.
- For covariance, let's consider the mean-zero case (which has the same covariance):

$$\text{Cov}(X + Y) = \mathbb{E}[(X + Y)(X + Y)']$$

Properties of multivariate Gaussians

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$.
- For covariance, let's consider the mean-zero case (which has the same covariance):

$$\begin{aligned}\text{Cov}(X + Y) &= \mathbb{E}[(X + Y)(X + Y)'] \\ &= \mathbb{E}[XX' + XY' + YX' + YY']\end{aligned}$$

Properties of multivariate Gaussians

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$.
- For covariance, let's consider the mean-zero case (which has the same covariance):

$$\begin{aligned}\text{Cov}(X + Y) &= \mathbb{E}[(X + Y)(X + Y)'] \\ &= \mathbb{E}[XX' + XY' + YX' + YY'] \\ &= \mathbb{E}[XX'] + \mathbb{E}[XY'] + \mathbb{E}[YX'] + \mathbb{E}[YY']\end{aligned}$$

Properties of multivariate Gaussians

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$.
- For covariance, let's consider the mean-zero case (which has the same covariance):

$$\begin{aligned}\text{Cov}(X + Y) &= \mathbb{E}[(X + Y)(X + Y)'] \\ &= \mathbb{E}[XX' + XY' + YX' + YY'] \\ &= \mathbb{E}[XX'] + \mathbb{E}[XY'] + \mathbb{E}[YX'] + \mathbb{E}[YY'] \\ &= \mathbb{E}[XX'] + 0 + 0 + \mathbb{E}[YY']\end{aligned}$$

Properties of multivariate Gaussians

- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$.
- For covariance, let's consider the mean-zero case (which has the same covariance):

$$\begin{aligned}\text{Cov}(X + Y) &= \mathbb{E}[(X + Y)(X + Y)'] \\ &= \mathbb{E}[XX' + XY' + YX' + YY'] \\ &= \mathbb{E}[XX'] + \mathbb{E}[XY'] + \mathbb{E}[YX'] + \mathbb{E}[YY'] \\ &= \mathbb{E}[XX'] + 0 + 0 + \mathbb{E}[YY'] \\ &= \Sigma_X + \Sigma_Y.\end{aligned}$$

Properties of multivariate Gaussians

Sums of independent Gaussians

For independent d -dimensional Gaussians,
 $X \sim \mathcal{N}(\mu_X, \Sigma_X)$ and $Y \sim \mathcal{N}(\mu_Y, \Sigma_Y)$,
 $X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \Sigma_X + \Sigma_Y)$.

- We need X and Y independent.

Properties of multivariate Gaussians

Sums of independent Gaussians

For independent d -dimensional Gaussians,
 $X \sim \mathcal{N}(\mu_X, \Sigma_X)$ and $Y \sim \mathcal{N}(\mu_Y, \Sigma_Y)$,
 $X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \Sigma_X + \Sigma_Y)$.

- We need X and Y independent. For example: $X \sim \mathcal{N}(0, 1)$,

$$Y = \begin{cases} -X & \text{if } |X| < 1, \\ X & \text{if } |X| > 1. \end{cases}$$

Properties of multivariate Gaussians

Sums of independent Gaussians

For independent d -dimensional Gaussians,
 $X \sim \mathcal{N}(\mu_X, \Sigma_X)$ and $Y \sim \mathcal{N}(\mu_Y, \Sigma_Y)$,
 $X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \Sigma_X + \Sigma_Y)$.

- We need X and Y independent. For example: $X \sim \mathcal{N}(0, 1)$,

$$Y = \begin{cases} -X & \text{if } |X| < 1, \\ X & \text{if } |X| > 1. \end{cases}$$

Then $X \sim \mathcal{N}(0, 1)$ and $Y \sim \mathcal{N}(0, 1)$.

Properties of multivariate Gaussians

Sums of independent Gaussians

For independent d -dimensional Gaussians,
 $X \sim \mathcal{N}(\mu_X, \Sigma_X)$ and $Y \sim \mathcal{N}(\mu_Y, \Sigma_Y)$,
 $X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \Sigma_X + \Sigma_Y)$.

- We need X and Y independent. For example: $X \sim \mathcal{N}(0, 1)$,

$$Y = \begin{cases} -X & \text{if } |X| < 1, \\ X & \text{if } |X| > 1. \end{cases}$$

Then $X \sim \mathcal{N}(0, 1)$ and $Y \sim \mathcal{N}(0, 1)$.

But (X, Y) is not jointly Gaussian.

(Check!)

Properties of multivariate Gaussians

Sums of independent Gaussians

For independent d -dimensional Gaussians,
 $X \sim \mathcal{N}(\mu_X, \Sigma_X)$ and $Y \sim \mathcal{N}(\mu_Y, \Sigma_Y)$,
 $X + Y \sim \mathcal{N}(\mu_X + \mu_Y, \Sigma_X + \Sigma_Y)$.

- We need X and Y independent. For example: $X \sim \mathcal{N}(0, 1)$,

$$Y = \begin{cases} -X & \text{if } |X| < 1, \\ X & \text{if } |X| > 1. \end{cases}$$

Then $X \sim \mathcal{N}(0, 1)$ and $Y \sim \mathcal{N}(0, 1)$.

But (X, Y) is not jointly Gaussian.

(Check!)

And $X + Y$ is not Gaussian.

(Check!)

Properties of multivariate Gaussians

Marginals

Given a d -dimensional Gaussian $X \sim \mathcal{N}(\mu, \Sigma)$, write

$$X = \begin{bmatrix} Y \\ Z \end{bmatrix}, \quad \mu = \begin{bmatrix} \mu_Y \\ \mu_Z \end{bmatrix}, \quad \Sigma = \begin{bmatrix} \Sigma_{YY} & \Sigma_{YZ} \\ \Sigma_{ZY} & \Sigma_{ZZ} \end{bmatrix},$$

where $Y \in \mathbb{R}^m$ and $Z \in \mathbb{R}^{d-m}$.

Properties of multivariate Gaussians

Marginals

Given a d -dimensional Gaussian $X \sim \mathcal{N}(\mu, \Sigma)$, write

$$X = \begin{bmatrix} Y \\ Z \end{bmatrix}, \quad \mu = \begin{bmatrix} \mu_Y \\ \mu_Z \end{bmatrix}, \quad \Sigma = \begin{bmatrix} \Sigma_{YY} & \Sigma_{YZ} \\ \Sigma_{ZY} & \Sigma_{ZZ} \end{bmatrix},$$

where $Y \in \mathbb{R}^m$ and $Z \in \mathbb{R}^{d-m}$. Then $Y \sim \mathcal{N}(\mu_Y, \Sigma_{YY})$.

- The mean and covariance of Y are immediate from the definitions.

Properties of multivariate Gaussians

Marginals

Given a d -dimensional Gaussian $X \sim \mathcal{N}(\mu, \Sigma)$, write

$$X = \begin{bmatrix} Y \\ Z \end{bmatrix}, \quad \mu = \begin{bmatrix} \mu_Y \\ \mu_Z \end{bmatrix}, \quad \Sigma = \begin{bmatrix} \Sigma_{YY} & \Sigma_{YZ} \\ \Sigma_{ZY} & \Sigma_{ZZ} \end{bmatrix},$$

where $Y \in \mathbb{R}^m$ and $Z \in \mathbb{R}^{d-m}$. Then $Y \sim \mathcal{N}(\mu_Y, \Sigma_{YY})$.

- The mean and covariance of Y are immediate from the definitions.
- To show that Y has a Gaussian distribution, we write its density as an integral (integrating out the Z variables), *complete the squares* to compute the integral, and recognize the answer as a $\mathcal{N}(\mu_Y, \Sigma_{YY})$ density.

Properties of multivariate Gaussians

Affine transformations

Given a d -dimensional Gaussian $X \sim \mathcal{N}(\mu, \Sigma^2)$, matrix $A \in \mathbb{R}^{m \times d}$ and vector $b \in \mathbb{R}^m$, define

$$Y = AX + b.$$

Properties of multivariate Gaussians

Affine transformations

Given a d -dimensional Gaussian $X \sim \mathcal{N}(\mu, \Sigma^2)$, matrix $A \in \mathbb{R}^{m \times d}$ and vector $b \in \mathbb{R}^m$, define

$$Y = AX + b.$$

Then $Y \sim \mathcal{N}(A\mu + b, A\Sigma A')$.

Properties of multivariate Gaussians

Affine transformations

Given a d -dimensional Gaussian $X \sim \mathcal{N}(\mu, \Sigma^2)$, matrix $A \in \mathbb{R}^{m \times d}$ and vector $b \in \mathbb{R}^m$, define

$$Y = AX + b.$$

Then $Y \sim \mathcal{N}(A\mu + b, A\Sigma A')$.

- Why is Y Gaussian?

Suppose that $m = d$ and A has full rank. Then Y is a shifted, rescaled version of X , and hence is Gaussian.

Properties of multivariate Gaussians

Affine transformations

Given a d -dimensional Gaussian $X \sim \mathcal{N}(\mu, \Sigma^2)$, matrix $A \in \mathbb{R}^{m \times d}$ and vector $b \in \mathbb{R}^m$, define

$$Y = AX + b.$$

Then $Y \sim \mathcal{N}(A\mu + b, A\Sigma A')$.

- Why is Y Gaussian?

Suppose that $m = d$ and A has full rank. Then Y is a shifted, rescaled version of X , and hence is Gaussian.

- $\mathbb{E}Y = A\mu + b$.

Properties of multivariate Gaussians

Affine transformations

Given a d -dimensional Gaussian $X \sim \mathcal{N}(\mu, \Sigma^2)$, matrix $A \in \mathbb{R}^{m \times d}$ and vector $b \in \mathbb{R}^m$, define

$$Y = AX + b.$$

Then $Y \sim \mathcal{N}(A\mu + b, A\Sigma A')$.

- Why is Y Gaussian?

Suppose that $m = d$ and A has full rank. Then Y is a shifted, rescaled version of X , and hence is Gaussian.

- $\mathbb{E}Y = A\mu + b$.
- $\text{Cov}(Y) = \mathbb{E}[(Y - \mathbb{E}Y)(Y - \mathbb{E}Y)'] = \mathbb{E}[(A(X - \mu))(A(X - \mu))'] = A\mathbb{E}[(X - \mu)(X - \mu)']A' = A\Sigma A'$.

Properties of multivariate Gaussians

Example

Given a d -dimensional standard normal $X \sim \mathcal{N}(0, I)$, we have

$$Y = AX + \mu$$

Properties of multivariate Gaussians

Example

Given a d -dimensional standard normal $X \sim \mathcal{N}(0, I)$, we have

$$Y = AX + \mu \sim \mathcal{N}(\mu, AA').$$

Properties of multivariate Gaussians

Example

Given a d -dimensional standard normal $X \sim \mathcal{N}(0, I)$, we have

$$Y = AX + \mu \sim \mathcal{N}(\mu, AA').$$

Example

Given a d -dimensional normal $X \sim \mathcal{N}(\mu, \Sigma)$ with Σ positive definite,

$$Y = \Sigma^{-1/2}(X - \mu)$$

Properties of multivariate Gaussians

Example

Given a d -dimensional standard normal $X \sim \mathcal{N}(0, I)$, we have

$$Y = AX + \mu \sim \mathcal{N}(\mu, AA').$$

Example

Given a d -dimensional normal $X \sim \mathcal{N}(\mu, \Sigma)$ with Σ positive definite,

$$Y = \Sigma^{-1/2}(X - \mu) \sim \mathcal{N}(0, I).$$

Properties of multivariate Gaussians

Example

Given a d -dimensional standard normal $X \sim \mathcal{N}(0, I)$, we have

$$Y = AX + \mu \sim \mathcal{N}(\mu, AA').$$

Example

Given a d -dimensional normal $X \sim \mathcal{N}(\mu, \Sigma)$ with Σ positive definite,

$$Y = \Sigma^{-1/2}(X - \mu) \sim \mathcal{N}(0, I).$$

This is called *whitening* X .

Properties of multivariate Gaussians

Example

Given a d -dimensional standard normal $X \sim \mathcal{N}(0, I)$, we have

$$Y = AX + \mu \sim \mathcal{N}(\mu, AA').$$

Example

Given a d -dimensional normal $X \sim \mathcal{N}(\mu, \Sigma)$ with Σ positive definite,

$$Y = \Sigma^{-1/2}(X - \mu) \sim \mathcal{N}(0, I).$$

This is called *whitening* X .

What is $\Sigma^{-1/2}$? We can write $\Sigma = U\Lambda U'$ for U with orthonormal columns and Λ diagonal. Then defining $\Sigma^{-1/2} = U\Lambda^{-1/2}U'$, we have

$$\Sigma^{-1/2}\Sigma^{-1/2} = U\Lambda^{-1/2}U'U\Lambda^{-1/2}U' = U\Lambda^{-1}U' = \Sigma^{-1}.$$

- Probability density function.
- 2-D Gaussian
- Covariance matrix.
- Diagonal covariance.
- Non-diagonal covariance and diagonalization.
- Properties of multivariate Gaussians.