

Comment on Hallmann

Valentina Gacitua, Leon Thoma, Dominik Arend

5 1 2021

Hallmann et al. (2017) revealed a massive decline in total flying insect biomass. Dramatics being added, this decrease was found to take place on sites of nature conservation. The Authors reported a decline of “More than 75 percent” over 27 years. The majority of contributing effects to this development remains unknown. # Motivation for the Re-Analysis We aim to reanalyze this highly relevant paper. Its impact is to this day enormous, resulting in broad media coverage, high citing scores and influence on political decision makers. Therefore we want to assess the robustness of its results, mainly the decline on insect biomass.

Nevertheless we want to point out critical points about the publication, check for its relevance and discuss the paper within the context of our analysis of the data. This bulletpoints give an overview over crucial aspects of the analysis:

- Years 1989 and 2014 are over-represented
- Few locations were re-sampled
- Only one trap per location
- The exposure time varies greatly among years
- Unknown site selection procedure
- (Lack of control group ?)

The Lack of a control group and the few re-sampling sites arise the suspect of a “regression to the mean problem”. This means that of two successive measurements, the second one tends to be closer to the true underlying mean (Kelly et al. 2005). Barnett, Pols, and Dobson (2004) describe regression to the mean (RTM) as a statistical phenomenon where the normal variation of repeated measurements appears to be a real change, simply because the often the measurement is repeated, the closer it reaches to the mean. Further RTM can appear as a selection phenomenon, when there is no control group and not a random sampling of measurements sites or individuals (Blomqvist 1987). Both cases could be anticipated in the present publication. Since we could not test the results of Hallmann et al. (2017) on RTM by redesigning the sampling design, we tried to outrule the RTM effect by only including the first year of sampling per insect collection plot in our analysis.

Following Hallmann et al. (2017) the basic model was calculated to assess the temporal dynamics of flying insect biomass. Calculations of the latent expected daily mass and model specification provided the foot for our model. With 24 000 iterations we tried to provide the same quality as the original analysis. The diagnosis of the model was done nevertheless our model did only differ in being a Subset of the original Data. No problems with did appear, all variables were uncorrelated.

For the results, our analysis did not differ substantially from Hallmann et al. (2017). The hypothesis of a intensified insect biomass decline due to regression to the mean could not be verified. We report a decline of flying insect biomass by XX % within 27 years. This is a discrepancy by XX % from the original result (~ 75 % decline). A RTM effect is therefore not found. The stated results seem to be robust from this point of view.

Hence our analysis is no cure for the few re-sampled locations and the unknown plot selection procedure. As the data collection was not carried out by the authors of the paper, it remains uncertain if there has been fewer entomological interesting sites left over after years of data collection. This would introduce a decline in insect biomass, caused by plot selection.

As pointed out, there is no other way to check for RTM via data analysis. Blomqvist (1987) states that

RTM is often, like in this case possible, appearing as a selection phenomenon. To introduce control groups is often advised to control RTM effects (Kelly et al. 2005; Barnett, Pols, and Dobson 2004). Within ecological research, investigating temporal changes, this can be hard to implement. In this special case, More than one sampling procedure per location could have introduced an assured measure of temporal change.

Different trap exposure times were treated for. By modeling the biomass of the catches as the sum of expected latent biomass, the authors eliminated this problem.

- Why different model i
- Our Result
- Our Result to Hallmann
- Why is Hallmann Right/Wrong
- How is the Statistical analysis of Hallmann in general
- (Why is there a need for more research in this area) Control Group suggestion?? How could this be done? → RTM also appears as a selection phenomenon, emphasizing the need to include control groups in order to make possible adjustments for the bias caused by RTM. (Blomqvist 1987)

References

- Barnett, Adrian G, Jolieke C van der Pols, and Annette J Dobson. 2004. “Regression to the mean: what it is and how to deal with it.” *International Journal of Epidemiology* 34 (1): 215–20. <https://doi.org/10.1093/ije/dyh299>.
- Blomqvist, Nils. 1987. “On the Bias Caused by Regression Toward the Mean in Studying the Relation Between Change and Initial Value.” *Journal of Clinical Periodontology* 14 (1): 34–37. <https://doi.org/10.1111/j.1600-051X.1987.tb01510.x>.
- Hallmann, Caspar A., Martin Sorg, Eelke Jongejans, Henk Siepel, Nick Hofland, Heinz Schwan, Werner Stenmans, et al. 2017. “More Than 75 Percent Decline over 27 Years in Total Flying Insect Biomass in Protected Areas.” *PLOS ONE* 12 (10): 1–21. <https://doi.org/10.1371/journal.pone.0185809>.
- Kelly, Colleen, Trevor D. Price, Associate Editor: Stuart A. West, and Editor: Michael C. Whitlock. 2005. “Correcting for Regression to the Mean in Behavior and Ecology.” *The American Naturalist* 166 (6): 700–707. <http://www.jstor.org/stable/10.1086/497402>.