


Noise Analysis

Optomechanical Cavity

Leon Oleschko
31.12.2024

Modeling Quantum Hardware: open dynamics and control
Universität Konstanz

Problem Statement



"Cavity optomechanics", Aspelmeyer et al. 2014

Quantum Optomechanics, Bowen et al. 2015

Hamiltonian

Optical Cavity \hat{a} , $\omega_o(\hat{x}_{\text{mech}}) = \omega_o + \frac{g}{\omega_o} \hat{x}_{\text{mech}}$; mechanical oscillations \hat{b} , ω_m ; coupling g ; Drive E , ω_L

$$H = \underbrace{\omega_o a^\dagger a}_{\text{Cavity}} + \underbrace{\omega_m b^\dagger b}_{\text{Mechanical}} - \underbrace{g a^\dagger a (b + b^\dagger)}_{\text{Interaction}} + \underbrace{i(E a^\dagger e^{-i\omega_L t} - E^* a e^{i\omega_L t})}_{\text{Drive}}$$

$$\hbar = 1$$

Quantum Optomechanics, Bowen et al. 2015 (2.3)

Hamiltonian

Optical Cavity \hat{a} , $\omega_o(\hat{x}_{\text{mech}}) = \omega_o + \frac{g}{\omega_o} \hat{x}_{\text{mech}}$; mechanical oscillations \hat{b} , ω_m ; coupling g ; Drive E , ω_L

$$H = \underbrace{\omega_o a^\dagger a}_{\text{Cavity}} + \underbrace{\omega_m b^\dagger b}_{\text{Mechanical}} - \underbrace{g a^\dagger a (b + b^\dagger)}_{\text{Interaction}} + \underbrace{i(E a^\dagger e^{-i\omega_L t} - E^* a e^{i\omega_L t})}_{\text{Drive}}$$

Rotating Wave Approximation at ω_L with $\Delta = \omega_o - \omega_L$, $a \rightarrow a e^{i\omega_L t}$:

$$H_{\text{RWA}} = \Delta a^\dagger a + \omega_m b^\dagger b - g a^\dagger a (b^\dagger + b) + i(E a^\dagger - E^* a)$$

$$\hbar = 1$$

Quantum Optomechanics, Bowen et al. 2015 (2.3)

Hamiltonian Linearization

$$H_{\text{RWA}} = \Delta a^\dagger a + \omega_m b^\dagger b - g a^\dagger a (b^\dagger + b) + i(Ea^\dagger - E^* a)$$

Linearize $a = \alpha + \delta a$, $b = \beta + \delta b$; with α, β steady state.

$$\begin{aligned} H_{\text{Interaction}} &= -g a^\dagger a (b^\dagger + b) \\ &\approx -\underbrace{g|\alpha|}_G (\delta a + \delta a^\dagger + \mathcal{O}(a^2 + \delta a \delta a^\dagger)) (\delta b + \delta b^\dagger + 2\beta) \end{aligned}$$

Quantum Optomechanics, Bowen et al. 2015 (2.7)

Hamiltonian Linearization

$$H_{\text{RWA}} = \Delta a^\dagger a + \omega_m b^\dagger b - g a^\dagger a (b^\dagger + b) + i(Ea^\dagger - E^* a)$$

Linearize $a = \alpha + \delta a$, $b = \beta + \delta b$; with α, β steady state.

$$\begin{aligned} H_{\text{Interaction}} &= -g a^\dagger a (b^\dagger + b) \\ &\approx -\underbrace{g|\alpha|}_G (\delta a + \delta a^\dagger + \mathcal{O}(a^2 + \delta a \delta a^\dagger)) (\delta b + \delta b^\dagger + 2\beta) \end{aligned}$$

Therefore for small G :

$$H \approx \Delta \delta a^\dagger \delta a + \omega_m \delta b^\dagger \delta b - G(\delta a + \delta a^\dagger)(\delta b + \delta b^\dagger)$$

Quantum Optomechanics, Bowen et al. 2015 (2.7)

Dissipation

Optical decay κ :

$$\sqrt{\kappa} \delta a$$

Mechanical resonator with γ and a thermal bath at the n -th thermal state:

$$\sqrt{\gamma(n+1)} \delta b + \sqrt{\gamma n} \delta b^\dagger$$

Quantum Optomechanics, Bowen et al. 2015 (2.8)

