

CSC421: Written Homework 2

Due on Monday, Feb 11, 2019

Zhongtian Ouyang
1002341012

Problem 1

Adam

(a)

$$(\alpha_A, \beta_1, \beta_2, \epsilon_A) = (\alpha_R, 0, \gamma, \epsilon_R)$$

Since Adam is a combination of RMSprop and Momentum. If we don't do the momentum part, it will be identical to RMSprop.

By setting the hyperparameters of Adam as above,

$$m_t = \beta_1 m_{t-1} + (1 - \beta_1)g_t = 0m_{t-1} + (1 - 0)g_t = g_t,$$

$$v_t = \beta_2 v_{t-1} + (1 - \beta_2)g_t^2 = \gamma v_{t-1} + (1 - \gamma)g_t^2$$

$$\theta_t = \theta_{t-1} - \alpha_R g_t / (\sqrt{v_t} + \epsilon_R) = \theta_{t-1} - \alpha_A m_t / (\sqrt{v_t} + \epsilon_A) = \theta_{t-1} - \alpha_R g_t / (\sqrt{v_t} + \epsilon_R)$$

which is identical to RMSprop

(b)

$$(\alpha_A, \beta_1, \beta_2, \epsilon_A) = (\alpha_S, \mu, 1, 1)$$

We can make Adam and SGD with momentum equivalent by satisfying the following conditions: $m_t = -p_t$, $(\sqrt{v_t} + \epsilon_R) = 1$ and $\alpha_A = \alpha_S$

If $v_t = 0$, $\epsilon_A = 1$, $\sqrt{v_t} + \epsilon_A = 1$. We can achieve this by setting β_2 to 1. Since $v_0 = 0$, $v_1, v_2 \dots v_t = v_0 = 0$

We can prove by induction that $m_t = -p_t$ when $\beta_1 = \mu$.

$$m_0 = 0 = -p_0,$$

$$m_1 = \beta_1 m_0 + (1 - \beta_1)g_1 = (1 - \beta_1)g_1 = (1 - \mu)\nabla J(\theta_0), \quad p_1 = \mu p_0 - (1 - \mu)\nabla J(\theta_0) = -(1 - \mu)\nabla J(\theta_0). \quad m_1 = -p_1$$

$$\text{Assume } m_{t-1} = -p_{t-1}, \quad m_t = \beta_1 m_{t-1} + (1 - \beta_1)g_t = -\mu p_{t-1} + (1 - \mu)\nabla J(\theta_{t-1}) = -(\mu p_{t-1} - (1 - \mu)\nabla J(\theta_{t-1})), \quad p_t = \mu p_{t-1} - (1 - \mu)\nabla J(\theta_{t-1}). \quad m_t = -p_t$$

By setting the hyperparameters of Adam as above, $\theta_t = \theta_{t-1} - \alpha_A m_t / (\sqrt{v_t} + \epsilon_A) = \theta_{t-1} + \alpha_S p_t / 1 = \theta_{t-1} + \alpha_S p_t$, which is identical to SGD with momentum.

(c)

$$\tilde{L}(y, t) = CL(y, t), \nabla \tilde{J}(\theta_0) = C \nabla J(\theta_0), \tilde{g}_t = C g_t$$

Prove $\tilde{m}_t = C m_t$ for $t \geq 1$:

Base Case:

$$\tilde{m}_1 = \beta_1 \tilde{m}_0 + (1 - \beta_1) \tilde{g}_1 = C(1 - \beta_1) g_1, m_1 = \beta_1 m_0 + (1 - \beta_1) g_1 = (1 - \beta_1) g_1.$$

$$\tilde{m}_1 = C m_1.$$

Inductive Steps: Assume $\tilde{m}_{t-1} = C m_{t-1}$

$$\tilde{m}_t = \beta_1 \tilde{m}_{t-1} + (1 - \beta_1) \tilde{g}_t = C \beta_1 m_{t-1} + C(1 - \beta_1) g_t = C(\beta_1 m_{t-1} + (1 - \beta_1) g_t),$$

$$m_t = \beta_1 m_{t-1} + (1 - \beta_1) g_t.$$

$$\tilde{m}_t = C m_t.$$

Prove $\tilde{v}_t = C^2 v_t$ for $t \geq 1$:

Base Case:

$$\tilde{v}_1 = \beta_2 \tilde{v}_0 + (1 - \beta_2) \tilde{g}_1^2 = C^2(1 - \beta_2) g_1^2, v_1 = \beta_2 v_0 + (1 - \beta_2) g_1^2 = (1 - \beta_2) g_1^2.$$

$$\tilde{v}_1 = C^2 v_1.$$

Inductive Steps: Assume $\tilde{v}_{t-1} = C^2 v_{t-1}$

$$\tilde{v}_t = \beta_2 \tilde{v}_{t-1} + (1 - \beta_2) \tilde{g}_t^2 = C^2 \beta_2 v_{t-1} + C^2(1 - \beta_2) g_t^2 = C^2(\beta_2 v_{t-1} + (1 - \beta_2) g_t^2),$$

$$v_t = \beta_2 v_{t-1} + (1 - \beta_2) g_t^2.$$

$$\tilde{v}_t = C^2 v_t.$$

Prove $\tilde{\theta}_t = \theta_t$ for $t \geq 0$: Base Case: $\tilde{\theta}_0 = \theta_0$ because they initialize to the same value

Inductive Steps: Assume $\tilde{\theta}_{t-1} = \theta_{t-1}$

$$\tilde{\theta}_t = \tilde{\theta}_{t-1} - \alpha_A \tilde{m}_t / (\sqrt{\tilde{v}_t} + \epsilon_A) = \theta_{t-1} - \alpha_A C m_t / (\sqrt{C^2 v_t} + 0) = \theta_{t-1} - \alpha_A C m_t / C \sqrt{v_t} = \theta_{t-1} - \alpha_A m_t / \sqrt{v_t},$$

$$\theta_t = \theta_{t-1} - \alpha_A m_t / (\sqrt{v_t} + \epsilon_A) = \theta_{t-1} - \alpha_A m_t / (\sqrt{v_t} + 0) = \theta_{t-1} - \alpha_A m_t / \sqrt{v_t}$$

$$\tilde{\theta}_t = \theta_t$$