

# A fluconazole population pharmacokinetics study to improve target attainment in critically ill patients

My-Luong Vuong<sup>a</sup>, Omar Elkayal<sup>a</sup>, Ruth Van Daele<sup>a,b</sup>, Jan-Willem C. Alffenaar<sup>c,d,e</sup>, Sophie L. Stocker<sup>c,e,f,g</sup>, Jason A. Roberts<sup>h,i,j</sup>, Yves Debaveye<sup>k,l</sup>, Joost Wauters<sup>m,l</sup>, Beatrijs Mertens<sup>a,b</sup>, Jasper M. Boonstra<sup>n</sup>, Indy Sandaradura<sup>o,p,q</sup>, Deborah J.E. Marriott<sup>r,s</sup>, Roger J. Brüggemann<sup>t,u</sup>, Jeroen A. Schouten<sup>v</sup>, Raoul Bergner<sup>w</sup>, Steven Buijk<sup>x</sup>, Isabel Spriet<sup>b</sup>, Erwin Dreesen<sup>a,\*</sup>

<sup>a</sup>*KU Leuven, Department of Pharmaceutical and Pharmacological Sciences, Leuven, Belgium,*

<sup>b</sup>*University Hospitals Leuven, Pharmacy Department, Leuven, Belgium,*

<sup>c</sup>*The University of Sydney, School of Pharmacy, Faculty of Medicine and Health, Sydney, Australia,*

<sup>d</sup>*Westmead Hospital, Pharmacy Department, Sydney, Australia,*

<sup>e</sup>*The University of Sydney, Institute for Infectious Diseases (Sydney ID), Sydney, Australia,*

<sup>f</sup>*University of New South Wales, School of Clinical Medicine, Faculty of Medicine and Health, Sydney, Australia,*

<sup>g</sup>*St. Vincent's Hospital, Department of Clinical Pharmacology and Toxicology, Sydney, Australia,*

<sup>h</sup>*The University of Queensland, University of Queensland Centre for Clinical Research (UQCCR), Brisbane, Australia,*

<sup>i</sup>*Royal Brisbane and Women's Hospital, Departments of Pharmacy and Intensive Care Medicine, Brisbane, Australia,*

<sup>j</sup>*University of Montpellier, Division of Anaesthesiology Critical Care Emergency and Pain Medicine, Nîmes University Hospital, Nîmes, France,*

<sup>k</sup>*KU Leuven, Department of Cellular and Molecular Medicine, Leuven, Belgium,*

<sup>l</sup>*University Hospitals Leuven, Medical Intensive Care Unit, Leuven, Belgium,*

<sup>m</sup>*KU Leuven, Department of Microbiology, Immunology and Transplantation, Leuven, Belgium,*

<sup>n</sup>*University of Groningen, Department of Clinical Pharmacy and Pharmacology, University Medical Centre Groningen, Groningen, The Netherlands,*

<sup>o</sup>*Westmead Hospital, Centre for Infectious Diseases and Microbiology, Sydney, Australia,*

<sup>p</sup>*The University of Sydney, Faculty of Medicine and Health, Westmead Clinical School, Sydney, Australia,*

<sup>q</sup>*Westmead Hospital, Institute of Clinical Pathology and Medical Research, New South Wales Health Pathology, Sydney, Australia,*

---

\*Corresponding author

Email addresses: myluong.vuong@kuleuven.be (My-Luong Vuong), omarelkayal@hotmail.com (Omar Elkayal), isabel.spriet@uzleuven.be (Isabel Spriet), erwin.dreesen@kuleuven.be (Erwin Dreesen)

<sup>r</sup> *UNSW Sydney, School of Clinical Medicine, St Vincent's Healthcare Clinical Campus,  
Faculty of Medicine and Health, Sydney, Australia,*

<sup>s</sup> *St. Vincent's Hospital, Clinical Microbiology and Infectious Diseases  
Department, Sydney, Australia,*

<sup>t</sup> *Radboud university medical centre, Radboud Institute for Medical Innovations, Department  
of Pharmacy, Nijmegen, The Netherlands,*

<sup>u</sup> *Radboud university medical center/CWZ hospital, Centre of Expertise in  
Mycology, Nijmegen, The Netherlands,*

<sup>v</sup> *Canisius Wilhelmina Hospital, Department of Intensive Care, Nijmegen, The Netherlands,*

<sup>w</sup> *Ludwigshafen Medical Centre, Division of Rheumatology and Nephrology, Department of  
Medicine A, Ludwigshafen, Germany,*

<sup>x</sup> *Department of Surgery, IJsseland Hospital, Capelle aan den IJssel, Germany,*

---

---

## Supplementary Figures




Figure S1: Violin plot showing the distribution of 10 imputed sets of serum creatinine values and the observed one.




Figure S2: Density plot showing the distribution of 10 imputed sets of serum creatinine values and the observed one.