

constitutes a one-dimensional lattice of parallel gold ridges (50 nm height, 140 nm width) separated by a distance  $P$  defining the period of the lattice. The period  $P$  is connected to the SPP wavelength by  $\lambda_{SPP} = 2\pi/k_{SPP} < \lambda$  and to the angle of incidence reflection  $\theta_{SPP}$  of the SPP beam relatively to the (in plane) normal to the lattice by

$$P = \frac{\lambda_{SPP}}{2 \cos \theta_{SPP}}. \quad (24)$$

In the present case shown in Fig. 10 the Bragg mirror is optimized for  $\lambda = 800nm$  (i. e.,  $\lambda_{SPP} = 785nm$ ) and for  $\theta_{SPP} = 45^\circ$  incidence angle which means  $P \simeq 555nm$ . The experimental analysis of such a Bragg mirror when the resonance condition ( $\lambda, \theta_{SPP}$ ) is fulfilled reveals a very high reflectivity of  $R \simeq 95\%$  (see, for example, [34] for some earlier results on SPP Bragg mirrors studied with fluorescence microscopy). However, in the present experiment we choose an incident angle  $\theta = 65^\circ$ . As a consequence the reflectivity was much lower (see Fig. 10 A and more details in [39]). This configuration




FIG. 11: (A) Fourier space cross-cut along the short solid line in Fig. 10 B corresponding to the  $L$  beam (see inset). Experimental data (symbols) are compared with a Lorentz fit (solid curve). The intensity is normalized by setting the maximum intensity of  $L$  to 1. (B) Direct space cross-cut along the  $L$  beam in Fig. 10 A. Data points (symbols) are compared to an exponential fit (solid line). Data from [39] (©, Appl. phys. Lett., American Institute of physics, 2006).

reveals SPP interferences in the vicinity of the mirror (Fig. 10 A). In Fig. 10 B we show the