



US010189718B1

(12) **United States Patent**
Gross et al.

(10) **Patent No.:** **US 10,189,718 B1**
(45) **Date of Patent:** **Jan. 29, 2019**

(54) **METHODS FOR CHARGE-TITRATING PARTICLE ASSEMBLY, AND STRUCTURES PRODUCED THEREFROM**

(71) Applicant: **HRL Laboratories, LLC**, Malibu, CA (US)

(72) Inventors: **Adam F. Gross**, Santa Monica, CA (US); **Shanying Cui**, Calabasas, CA (US); **John J. Vajo**, West Hills, CA (US); **Christopher S. Roper**, Santa Monica, CA (US)

(73) Assignee: **HRL Laboratories, LLC**, Malibu, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 242 days.

(21) Appl. No.: **15/241,536**

(22) Filed: **Aug. 19, 2016**

Related U.S. Application Data

(60) Provisional application No. 62/212,969, filed on Sep. 1, 2015, provisional application No. 62/213,044, filed on Sep. 1, 2015.

(51) **Int. Cl.**
B01J 19/00 (2006.01)
C01G 49/02 (2006.01)
(Continued)

(52) **U.S. Cl.**
CPC **C01F 17/0062** (2013.01); **B01J 19/0046** (2013.01); **C01G 49/0045** (2013.01);
(Continued)

(58) **Field of Classification Search**
CPC B01J 19/0046; B01J 2219/00331; B01J 2219/00585; B01J 2219/00599;
(Continued)

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,534,461 A 7/1996 Kuwajima
5,591,535 A 1/1997 Hisano et al.
(Continued)

OTHER PUBLICATIONS

Barnov et al., "Chemically induced self-assembly of spherical and anisotropic inorganic nanocrystals," J. Mater. Chem. 2011, 21, 16694.

(Continued)

Primary Examiner — Jennifer Wecker

(74) *Attorney, Agent, or Firm* — O'Connor & Company

(57) **ABSTRACT**

Methods to fabricate tightly packed arrays of nanoparticles are disclosed, without relying on organic ligands or a substrate. In some variations, a method of assembling particles into an array comprises dispersing particles in a liquid solution; introducing a triggerable pH-control substance capable of generating an acid or a base; and triggering the pH-control substance to generate an acid or a base within the liquid solution, thereby titrating the pH. During pH titration, the particle-surface charge magnitude is reduced, causing the particles to assemble into a particle array. Other variations provide a device for assembling particles into particle arrays, comprising a droplet-generating microfluidic region; a first-fluid inlet port; a second-fluid inlet port; a reaction microfluidic region, disposed in fluid communication with the droplet-generating microfluidic region; and a trigger source configured to trigger generation of an acid or a base from at least one pH-control substance contained within the reaction microfluidic region.

31 Claims, 20 Drawing Sheets

