

where $B = \frac{25}{54}$ for $\Gamma_\varphi \gtrsim H_{\text{TI}}$ (Eq. (4.58)) or $B = \frac{25}{384}$ for $\Gamma_\varphi \ll H_{\text{TI}}$ (Eq. (4.62)).

Then, our m from Eq. (4.3) gives

$$g_{\text{NL}} \simeq B \left[2 + \frac{6\alpha - 3}{\alpha} \left(\frac{m_0^2 M_P^{2\alpha-2}}{2h^2 \psi^{2\alpha}} - 1 \right) + \frac{(2\alpha^2 - \alpha)(\alpha - 1)}{\alpha^3} \left(\frac{m_0^4 M_P^{4\alpha-4}}{4h^4 \psi^{4\alpha}} - \frac{m_0^2 M_P^{2\alpha-2}}{h^2 \psi^{2\alpha}} + 1 \right) \right] \quad (4.74)$$

In the parameter space available (if any), we will investigate the range of values for f_{NL} and g_{NL} .

4.4.3 Constraining the Free Parameters

4.4.3.1 Primordial Inflation Energy Scale

We want the energy scale of primordial inflation to be

$$V^{\frac{1}{4}} \lesssim 10^{14} \text{ GeV} \quad (4.75)$$

so that the inflaton contribution to the curvature perturbation is negligible.

Therefore, from the Friedmann equation $3M_P^2 H_*^2 = V$ we require

$$H_* \lesssim 10^{10} \text{ GeV} \quad (4.76)$$

4.4.3.2 Thermal Inflation Dynamics

We will consider only the case in which the inflationary trajectory is 1-dimensional, in that only the ϕ field is involved in determining the trajectory of thermal inflation in field space. We do this only to work with the simplest