

- [14] P. Jung and G. Mayer-Kress, Phys. Rev. Lett. **74**, 2130 (1995).
- [15] J. F. Lindner, B. K. Meadows, W. L. Ditto, M. E. Inchiosa, Adi R. Bulsara, Phys. Rev. Lett. **75**, 3 (1995).
- [16] L. Schimansky-Geier and U. Siewert, in *Stochastic Dynamics* (Springer, Berlin 1997) p. 245.
- [17] H. Gang, H. Haken, and X. Fagen, Phys. Rev. Lett. **77**, 1925 (1996).
- [18] A. Neiman, L. Schimansky-Geier, F. Moss, Phys. Rev. E **56**, R9 (1997).
- [19] J. M. Casado, J. Gómez-Ordóñez and M. Morillo, Phys. Rev. E **73**, 011109 (2006).
- [20] M. Morillo, J. Gómez Ordóñez and J. M. Casado, Phys. Rev. E **78**, 021109 (2008).
- [21] C. Tessone, C.R. Mirasso, R. Toral, J.D. Gunton, Phys. Rev. Lett. **97**, 194101, (2006).
- [22] R. Toral, C. J. Tessone and J. Viana Lopes, Eur. Phys. J. Special Topics **143**, 59 (2007).
- [23] H. Chen, J. Zhang, J. Liu, Phys. Rev. E **75**, 041910 (2007).
- [24] J.A. Acebrón, S. Lozano, A. Arenas, Phys. Rev. Lett. **99** 128701 (2007); Erratum: **99**, 229902(E) (2007).
- [25] M. Gosak, Biophysical Chemistry, **139**, Issue 1, 53 (2009).
- [26] R. Toral, E. Hernandez-Garcia, J. D. Gunton, International Journal of Bifurcation and Chaos, **19**, 10, 3499 (2009).
- [27] M. Perc, M. Gosak, S. Kralj, Soft Matter **4**, 1861 (2008).
- [28] T. Vaz Martins, R. Toral, M.A. Santos, Eur. Phys. J. B, **67**, N. 3, 329 (2009).
- [29] P. N. McGraw and M. Menzinger, Phys. Rev. E **77**, 031102 (2008).
- [30] *Spin glasses and random fields*, ed. A. P. Young, World Scientific, Singapore (1997).
- [31] P. Jung and P. Hänggi, Europhys. Lett. **8**, 505 (1989).
- [32] M. San Miguel and R. Toral, Stochastic effects in physical systems, in *Instabilities and Nonequilibrium Structures VI*, eds. E. Tirapegui, J. Martínez and R. Tiemann, Kluwer Academic Publishers 35-130 (2000).
- [33] V. N. Livina, F. Kwasniok, and T. M. Lenton, Clim. Past Discuss. **5**, 2223 (2009). The method detects the number of states and the shape of the potential of a dynamical system, from its recorded time series, by means of estimation of the degree of a polynomial potential. The coefficients that determine the shape of the potential are then estimated using the unscented Kalman filter (UKF). The method was developed for the stochastic model with noise component and performs polynomial fit of the histogram which is expected to be smooth (i.e. no discrete peak). To detect the number of states in this paper, we had to add a noise component to the model trajectories to make the method applicable. Yet, to minimize the effect of noise, we considered white noise of small amplitude (0.01 of the trajectory amplitude), which allowed us to "fill" the histogram and make it suitable for smooth polynomial fit. Note that adding noise can only hide a certain shallow well in the potential, but never lead to a false detection of non-existent states, and therefore adding noise cannot lead to false detection of additional states.
- [34] P. W. Anderson, Phys. Rev. **109**, 1492 (1958).
- [35] P. W. Anderson, Mater. Res. Bull. **5**, 549 (1970).
- [36] C.J. Tessone, D.H. Zanette, R. Toral, Eur. Phys. J. B **62**, 319 (2008).
- [37] I. Leyva, I. Sendiña-Nadal, J. A. Almendral, and M. A. Sanjuan, Phys. Rev. E, **74**, 056112 (2006).