

- [9] G. Liu *et al.*, *Langmuir* **24**, 7923 (2008).
- [10] U. Thiele *et al.*, *Phys. Rev. Lett.* **80**, 2869 (1998).
- [11] I. I. Smalyukh *et al.*, *Phys. Rev. Lett.* **96**, 177801 (2006).
- [12] L. Zhang *et al.*, *Langmuir* **24**, 3911 (2008).
- [13] G. Reiter, *Phys. Rev. Lett.* **68**, 75 (1992).
- [14] R. Seemann *et al.*, *Phys. Rev. Lett.* **86**, 5534 (2001).
- [15] M. Elbaum and S. G. Lipson, *Phys. Rev. Lett.* **72**, 3562 (1994).
- [16] Y. Gotkis *et al.*, *Phys. Rev. Lett.* **97**, 186101 (2006).
- [17] D. Quemada and C. Berli, *Adv. Colloid Interface Sci.* **98**, 51 (2002).
- [18] U. Thiele *et al.*, *Phys. Rev. Lett.* **87**, 016104 (2001).
- [19] J. Becker *et al.*, *Nat. Mater.* **2**, 59 (2003).
- [20] C. Redon *et al.*, *Phys. Rev. Lett.* **66**, 715 (1991).
- [21] A. Sharma and G. Reiter, *J. Colloid Interface Sci.* **178**, 383 (1996).
- [22] G. Reiter and A. Sharma, *Phys. Rev. Lett.* **87**, 166103 (2001).
- [23] A. Münch and B. Wagner, *Physica D* **209**, 178 (2005).
- [24] S. Kalliadasis and U. Thiele, eds., *Thin Films of Soft Matter* (Springer, Wien / New York, 2007).
- [25] M. Maillard *et al.*, *J. Phys. Chem. B* **104**, 11871 (2000).
- [26] U. Thiele, Ph.D. thesis, Technische Universität Dresden (1998).
- [27] A. Oron *et al.*, *Rev. Mod. Phys.* **69**, 931 (1997).
- [28] A. V. Lyushnin *et al.*, *Phys. Rev. E* **65**, 021602 (2002).
- [29] M. R. E. Warner *et al.*, *J. Colloid Interface Sci.* **267**, 92 (2003).
- [30] E. Rabani *et al.*, *Nature* **426**, 271 (2003).
- [31] C. P. Martin *et al.*, *Phys. Rev. Lett.* **99**, 116103 (2007).
- [32] I. Vancea *et al.*, *Phys. Rev. E* **78**, 041601 (2008).
- [33] U. Marconi and P. Tarazona, *J. Chem. Phys.* **110**, 8032 (1999).
- [34] A. J. Archer and M. Rauscher, *J. Phys. A: Math. Gen.* **37**, 9325 (2004).
- [35] J. F. Gouyet *et al.*, *Adv. Phys.* **52**, 523 (2003).
- [36] Note that our DDFT shares some features in common with the phase field theories used to study solidification and other such processes – see e.g. A.A. Wheeler *et al.*, *Phys. Rev. A* **45**, 7424 (1992).
- [37] P. M. Chaikin and T. C. Lubensky, *Principles of condensed matter physics* (Cambridge University Press, 1997).
- [38] We denote it a “semi-grand” free energy, because the free energy is developed in an ensemble where the liquid is treated grand canonically (it is coupled to a reservoir with chemical potential μ), whereas the fixed number of nanoparticles are treated canonically.
- [39] D. Woywod and M. Schoen, *Phys. Rev. E* **73**, 011201 (2006).
- [40] J. S. Langer, in *Solids far from Equilibrium*, edited by C. Godreche (Cambridge University Press, 1992), pp. 297–363.
- [41] G. Reiter, *Langmuir* **9**, 1344 (1993).