Solving Eq. (A.27) self-consistently with Eq. (A.21), it is easy to get the effective harmonic constant

$$f = K + 3\lambda \gamma_c^2 = \frac{K + \sqrt{K^2 + 12\lambda K_B T}}{2}.$$
 (A.29)

- [1] S. Lepri, R. Livi, and A. Politi, Phys. Rep. 377, 1 (2003).
- [2] Z. Rieder, J. L. Lebowitz, and E. Lieb, J. Math. Phys. 8, 1073 (1967).
- [3] N. Li, P. Tong, and B. Li, Europhys. Lett. **75**, 49 (2006).
- [4] N. Li and B. Li, Europhys. Lett. 78, 34001 (2007).
- [5] N. Li and B. Li, Phys. Rev. E **76**, 011108 (2007).
- [6] B. Hu, D. He, L. Yang, and Y. Zhang, Phys. Rev. E 74, 060101(R) (2006).
- [7] G. P. Srivastava, The Physics of Phonons (IOP Publishing Ltd, Bristol, 1990).
- [8] S. Lepri, Phys. Rev. E **58**, 7165 (1998).
- [9] S. Lepri, R. Livi, and A. Politi, Europhys. Lett. 43, 271 (1998).
- [10] P. Brüesch, *Phonons: Theory and Experiments I*, Lattice Dynamics and Models of Interatomic Forces (Springer-Verlag, New York, 1982).
- [11] T. Dauxois, M. Peyrard, and A. R. Bishop, Phys. Rev. E 47, 684 (1993).
- [12] C. Alabiso, M. Casartelli, and P. Marenzoni, J. Stat. Phys. 79, 451 (1995).
- [13] C. Alabiso and M. Casartelli, J. Phys. A 34, 1223 (2001).
- [14] K. Aoki and D. Kusnezov, Phys. Rev. Lett. 86, 4029 (2001).
- [15] K. Giannò, A. V. Sologubenko, H. R. Ott, A. D. Bianchi, and Z. Fisk, J. Phys.: Condens. Matter 15, 6739 (2003).
- [16] D. G. Cahill, W. K. Ford, K. E. Goodson, G. D. Mahan, A. Majumdar, H. J. Maris, R. Merlin, and S. R. Phillpot, J. Appl. Phys 93, 793 (2003).
- [17] R. E. Peierls, Quantum Theory of Solids (Oxford University Press, London, 1955).
- [18] J. Hone, M. Whitney, C. Piskoti, and A. Zettl, Phys. Rev. B 59, R2514 (1999).
- [19] M. S. Dresselhaus and P. C. Eklund, Adv. Phys. 49, 705 (2000).
- [20] D. G. Cahill, K. Goodson, and A. Majumdar, J. Heat Transfer 124, 223 (2002).
- [21] R. P. Feynman and H. Kleinert, Phys. Rev. A 34, 5080 (1986).
- [22] H. Kleinert, Path Integrals in Quantum Mechanics, Statistics, and Polymer Physics (World