- [7] Calderón, A. P. (1958). Uniqueness in the Cauchy problem for partial differential equations. American Journal of Mathematics, 80(1), 16-36.
- [8] Carleman, T. (1939). Sur un probleme, d'unicité pour les systemes d'équations aux derivées partielles deux variables independentes. Ark. Mat. Astr. Fys, 2, 1-9.
- [9] Cristofol, M., and Soccorsi, E. (2011). Stability estimate in an inverse problem for non-autonomous magnetic Schrödinger equations. Applicable Analysis, 90(10), 1499-1520.
- [10] Davey, A., and Stewartson, K. (1974). On three-dimensional packets of surface waves. In Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences, 338(1613), 101-110.
- [11] Djordjevic, V. D., and Redekopp, L. G. (1977). On two-dimensional packets of capillary-gravity waves. Journal of Fluid Mechanics, 79(4), 703-714.
- [12] Gölgeleyen, F., and Yamamoto, M. (2014). Stability of inverse problems for ultrahyperbolic equations. Chinese Annals of Mathematics, Series B, 35(4), 527-556.
- [13] Hörmander, L. (1963). Linear partial differential operators, Springer, Berlin, 285 pp.
- [14] Ichinose, W. (1990). A note on the Cauchy problem for Schrödinger type equations on the Riemannian manifold. Math. Japon, 35, 205-213.
- [15] Imanuvilov, O. Y., and Yamamoto, M. (2001). Global Lipschitz stability in an inverse hyperbolic problem by interior observations. Inverse problems, 17(4), 717-728.
- [16] Imanuvilov, O. Y., and Yamamoto, M. (2001). Global uniqueness and stability in determining coefficients of wave equations. Communications in Partial Differential Equations, 26(7-8), 1409-1425.