



13

-continued



Oxime Sulfonate


A further embodiment of the invention is an oxime sulfonate compound having the general formula (3b).



Herein RO is hydroxyl or a substituted or unsubstituted, straight, branched or cyclic $\text{C}_1\text{--C}_{20}$ organoxy group; R^1 is a divalent $\text{C}_1\text{--C}_{20}$ aliphatic group which may contain a substituent group having a heteroatom such as oxygen, nitrogen or sulfur, or may form a mono- or polycyclic structure with RO; q is 0 (zero) or 1; when q is 0, p is a substituted or unsubstituted $\text{C}_1\text{--C}_{20}$ alkyl group or a substituted or unsubstituted $\text{C}_6\text{--C}_{15}$ aryl group; when q is 1, p is a single bond, a substituted or unsubstituted $\text{C}_1\text{--C}_{20}$ alkylene group or a substituted or unsubstituted $\text{C}_6\text{--C}_{15}$ arylene group; EWG is a cyano, trifluoromethyl, perfluoroethyl, perfluoropropyl, 5H-perfluoropentyl, 6H-perfluorohexyl, nitro or methyl group, and when q is 1, two EWG's may bond together to form a ring of 6 carbon atoms with the carbon atoms to which they are attached.

In formula (3b), RO and R^1 are as defined above. When q is equal to 0, p is a substituted or unsubstituted $\text{C}_1\text{--C}_{20}$ alkyl group or a substituted or unsubstituted $\text{C}_6\text{--C}_{15}$ aryl group. When q is equal to 1, p is a single bond, a substituted or unsubstituted $\text{C}_1\text{--C}_{20}$ alkylene group or a substituted or unsubstituted $\text{C}_6\text{--C}_{15}$ arylene group. EWG is a cyano, trifluoromethyl, perfluoroethyl, perfluoropropyl, 5H-perfluoropentyl, 6H-perfluorohexyl, nitro or methyl group. When q is equal to 1, two EWG's may bond together to form a ring of 6 carbon atoms with the carbon atoms to which they are attached. The skeletons of the oxime sulfonates are described in U.S. Pat. No. 6,261,738, JP-A 9-95479, JP-A 9-208554, JP-A 9-230588, JP 2,906,999, JP-A 9-301948, JP-A 2000-314956, JP-A 2001-233842, and WO 2004/074242.

Exemplary skeletons of oxime sulfonates excluding the sulfonate moiety are given below wherein the broken line denotes a bonding site to the sulfonate moiety.



14

-continued

