- (2004).
- [119] J. M. Sage, S. Sainis, T. Bergeman, and D. DeMille, "Optical production of ultracold polar molecules", Phys. Rev. Lett. 94, 203001 (2005).
- [120] C. Haimberger, J. Kleinert, M. Bhattacharya, and N. P. Bigelow, "Formation and detection of ultracold ground-state polar molecules", Phys. Rev. A 70, 021402 (2004).
- [121] J. Kleinert, C. Haimberger, P. J. Zabawa, and N. P. Bigelow, "Trapping of ultracold polar molecules with a thin-wire electrostatic trap", Phys. Rev. Lett. 99, 143002 (2007).
- [122] M. W. Mancini, G. D. Telles, A. R. L. Caires, V. S. Bagnato, and L. G. Marcassa, "Observation of ultracold ground-state heteronuclear molecules", Phys. Rev. Lett. 92, 133203 (2004).
- [123] D. Wang, J. Qi, M. F. Stone, O. Nikolayeva, B. Hattaway, S. D. Gensemer, H. Wang, W. T. Zemke, P. L. Gould, E. E. Eyler and W. C. Stwalley, "The photoassociative spectroscopy, photoassociative molecule formation, and trapping of ultracold ³⁹K⁸⁵Rb", Eur. Phys. J. D 31, 165 (2004).
- [124] S. D. Kraft, P. Staanum, J. Lange, L. Vogel, R. Wester, and M. Weidemüller, "Formation of ultracold LiCs molecules", J. Phys. B: At. Mol. Opt. Phys. 39, S993 (2006).
- [125] U. Schlöder, C. Silber, and C. Zimmermann, "Photoassociation of heteronuclear lithium", Appl. Phys. B: Lasers Opt. 73, 801 (2001).
- [126] T.-G. Lee, N. Balakrishnan, R. C. Forrey, P. C. Stancil, D. R. Schultz, and G. J. Ferland, "State-to-state rotational transitions in H₂+H₂ collisions at low temperatures", J. Chem. Phys. 125, 114302 (2006).
- [127] R. C. Forrey, "Cooling and trapping of molecules in highly excited rotational states", Phys. Rev. A 63, 051403(R) (2001).
- [128] B. Maté, F. Thibault, G. Tejeda, J. M. Fernández, and S. Montero, "Inelastic collisions in para-H₂: Translation-rotation state-to-state rate coefficients and cross sections at low temperature and energy", J. Chem. Phys. 122, 064313 (2005).
- [129] P. Diep and J. K. Johnson, "An accurate H₂-H₂ interaction potential from first principles", J. Chem. Phys. 112, 4465 (2000).
- [130] S. Montero, F. Thibault, G. Tejeda, and J. M. Fernández, "Rotranslational state-to-state rates and spectral representation of inelastic collisions in low-temperature molecular hydrogen", J. Chem. Phys. 125, 124301 (2006).
- [131] A. I. Boothroyd, P. G. Martin, W. J. Keogh, and M. J. Peterson, "An accurate analytic H₄ potential energy surface", J. Chem. Phys. 116, 666 (2002).
- [132] B. Yang, P. C. Stancil, N. Balakrishnan, and R. C. Forrey, "Quenching of rotationally excited CO by collisions with H₂", J. Chem. Phys. 124, 104304 (2006).
- [133] B. Yang, H. Perera, N. Balakrishnan, R. C. Forrey, and P. C. Stancil, "Quenching of rotationally excited CO

- in cold and ultracold collisions with H, He and H_2 ", J. Phys. B: At. Mol. Opt. Phys. **39**, S1229 (2006).
- [134] A. V. Avdeenkov and J. L. Bohn, "Ultracold collisions of oxygen molecules", Phys. Rev. A 64, 052703 (2001).
- [135] A. V. Avdeenkov and J. L. Bohn, "Collisional dynamics of ultracold OH molecules in an electrostatic field", Phys. Rev. A 66, 052718 (2002).
- [136] A. V. Avdeenkov and J. L. Bohn, "Linking Ultracold Polar Molecules", Phys. Rev. Lett. 90, 043006 (2003).
- [137] A. V. Avdeenkov and J. L. Bohn, "Ultracold collisions of fermionic OD radicals", Phys. Rev. A 71, 022706 (2005).
- [138] C. Ticknor and J. L. Bohn, "Influence of magnetic fields on cold collisions of polar molecules", Phys. Rev. A 71, 022709 (2005).
- [139] S. K. Pogrebnya and D. C. Clary, "A full-dimensional quantum dynamical study of vibrational relaxation in H₂+H₂", Chem. Phys. Lett. 363, 523 (2002).
- [140] S. K. Pogrebnya, M. E. Mandy and D. C. Clary, "Vibrational relaxation in H₂+H₂: full-dimensional quantum dynamical study", Int. J. Mass. Spectrom. 223-224, 335 (2003).
- [141] A. N. Panda, F. Otto, F. Gatti, and H.-D. Meyer, "Rovibrational energy transfer in ortho-H₂+para-H₂ collisions", J. Chem. Phys. 127, 114310 (2007).
- [142] F. Otto, F. Gatti, and H.-D. Meyer, "Rotational excitations in para-H₂+para-H₂ collisions: Full- and reduced-dimensional quantum wave packet studies comparing different potential energy surfaces", J. Chem. Phys. 128, 064305 (2008).
- [143] G. Quéméner, N. Balakrishnan, and R. V. Krems, "Vibrational energy transfer in ultracold molecule-molecule collisions", Phys. Rev. A 77, 030704(R) (2008).
- [144] R. V. Krems, TwoBC quantum scattering program, University of British Columbia, Vancouver, Canada (2006).
- [145] A. M. Arthurs and A. Dalgarno, "The theory of scattering by a rigid rotator", Proc. Roy. Soc. A 256, 540 (1960).
- [146] K. Takayanagi, "The production of rotational and vibrational transitions in encounters between molecules", Adv. At. Mol. Phys. 1, 149 (1965).
- [147] S. Green, "Rotational excitation in H₂-H₂ collisions: Close-coupling calculations", J. Chem. Phys. 62, 2271 (1975).
- [148] M. H. Alexander and A. E. DePristo, "Symmetry considerations in the quantum treatment of collisions between two diatomic molecules", J. Chem. Phys. 66, 2166 (1977).
- [149] F. Ferlaino, S. Knoop, M. Mark, M. Berninger, H. Schöbel, H.-C. Nägerl, and R. Grimm, "Collisions between tunable halo dimers: exploring an elementary four-body process with identical bosons", Phys. Rev. Lett. 101, 023201 (2008).