

Here α and G_F , respectively, are the fine-structure constant and the Fermi constant, θ_i^2 is the mixing angle defined by

$$\theta_i^2 = \frac{(m_D^\dagger m_D)_{ii}}{m_{\chi_i}^2} = \mathcal{O} \left[\left(\frac{M_N^D}{M_N^M} \right)^2 \right] \frac{v_L^2}{v_R^2}. \quad (43)$$

With the previous parameter choice, we can determine the mixing angle to be $\theta_i^2 \simeq 10^{-22}$. Therefore the decay into the electron-positron pairs (42b) can [26] provide a natural explanation for the flux of 511 keV photons from the galactic bulge observed by INTEGRAL [27] satellite,

$$\frac{\Phi_{511\gamma}}{\Phi_{\text{exp}}} \simeq \sum_i \frac{\theta_i^2}{10^{-22}} \left(\frac{m_{\chi_i}}{1.3 \text{ MeV}} \right)^4 \frac{\Omega_{\chi_i}}{\Omega_\chi}. \quad (44)$$

It is easy to check that our scenario is consistent with other astrophysical and cosmological constraints [28]. Alternatively, we may explain the observed cosmic positron/electron excess [29–33], which is probably from continuum distribution of pulsars [34, 35], by fine tuning the parameters.

V. CONCLUSION

In summary we have shown the dark matter can be well determined by the neutrino masses and the baryon

asymmetry in the left-right symmetric model with doublet and singlet fields. In this model, the SM fermions obtain masses by integrating out charged singlet fermions. In the neutrino sector, the right(left)-handed neutral fermions, associated with the left(right)-handed Higgs doublet, can generate the left(right)-handed neutrino masses through the seesaw. The mass matrices of the left- and right-handed neutrinos have a same structure as a result of the left-right symmetry. The neutral singlets are also responsible for the baryon asymmetry and the dark matter. Specifically their 2-body decays can produce a desired lepton asymmetry in the left-handed leptons and then the observed baryon asymmetry can be realized by the sphaleron induced lepton-to-baryon conversion. At the same time, the right-handed neutrinos can serve as the dark matter as they have a right relic density from the 3-body decays of the neutral singlets. The decays of these non-thermally produced right-handed neutrinos can easily induce the observed fluxes of 511 keV photons from the Galactic bulge. The attractive feature of our scenario is that the left-right symmetry can connect the properties of the dark matter to the neutrino masses and the baryon asymmetry.

Acknowledgement: I thank Manfred Lindner for hospitality at Max-Planck-Institut für Kernphysik. This work is supported by the Alexander von Humboldt Foundation.

-
- [1] J. Dunkley *et al.*, [WMAP Collaboration], *Astrophys. J. Suppl.* **180**, 306 (2009).
 - [2] V.A. Kuzmin, arXiv: hep-ph/9701269; R. Kitano and I. Low, *Phys. Rev. D* **71**, 023510 (2005); N. Cosme, L. Lopez Honorez, and M.H.G. Tytgat, *Phys. Rev. D* **72**, 043505 (2005); D.E. Kaplan, M.A. Luty, and K.M. Zurek, *Phys. Rev. D* **79**, 115016 (2009); P.H. Gu, U. Sarkar, and X. Zhang, *Phys. Rev. D* **80**, 076003 (2009); I.M. Shoemaker and A. Kusenko, *Phys. Rev. D* **80**, 075021 (2009); P.H. Gu and U. Sarkar, arXiv:0909.5463 [hep-ph]; H. An, S.L. Chen, R.N. Mohapatra, and Y. Zhang, arXiv:0911.4463 [hep-ph].
 - [3] T. Schwetz, M.A. Tórtola, J.W.F. Valle, *New J. Phys.* **10**, 113011 (2008).
 - [4] P. Minkowski, *Phys. Lett. B* **67**, 421 (1977).
 - [5] T. Yanagida, in *Proc. of the Workshop on Unified Theory and the Baryon Number of the Universe*, ed. O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p. 95.
 - [6] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, ed. F. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979), p. 315.
 - [7] S.L. Glashow, in *Quarks and Leptons*, ed. M. Lévy *et al.* (Plenum, New York, 1980), p. 707.
 - [8] R.N. Mohapatra and G. Senjanović, *Phys. Rev. Lett.* **44**, 912 (1980).
 - [9] M. Magg and C. Wetterich, *Phys. Lett. B* **94**, 61 (1980); J. Schechter and J.W.F. Valle, *Phys. Rev. D* **22**, 2227 (1980); T.P. Cheng and L.F. Li, *Phys. Rev. D* **22**, 2860 (1980); G. Lazarides, Q. Shafi, and C. Wetterich, *Nucl. Phys. B* **181**, 287 (1981); R.N. Mohapatra and G. Senjanović, *Phys. Rev. D* **23**, 165 (1981).
 - [10] M. Fukugita and T. Yanagida, *Phys. Lett. B* **174**, 45 (1986).
 - [11] P. Langacker, R.D. Peccei, and T. Yanagida, *Mod. Phys. Lett. A* **1**, 541 (1986); M.A. Luty, *Phys. Rev. D* **45**, 455 (1992); R.N. Mohapatra and X. Zhang, *Phys. Rev. D* **46**, 5331 (1992).
 - [12] M. Flanz, E.A. Paschos, and U. Sarkar, *Phys. Lett. B* **345**, 248 (1995); M. Flanz, E.A. Paschos, U. Sarkar, and J. Weiss, *Phys. Lett. B* **389**, 693 (1996); L. Covi, E. Roulet, and F. Vissani, *Phys. Lett. B* **384**, 169 (1996).
 - [13] A. Pilaftsis, *Phys. Rev. D* **56**, 5431 (1997).
 - [14] E. Ma and U. Sarkar, *Phys. Rev. Lett.* **80**, 5716 (1998).
 - [15] W. Buchmüller, R.D. Peccei, and T. Yanagida, *Ann. Rev. Nucl. Part. Sci.* **55**, 311 (2005), and references therein.
 - [16] J.C. Pati and A. Salam, *Phys. Rev. D* **10**, 275 (1974); R.N. Mohapatra and J.C. Pati, *Phys. Rev. D* **11**, 566 (1975); R.N. Mohapatra and J.C. Pati, *Phys. Rev. D* **11**, 2558 (1975); R.N. Mohapatra and G. Senjanović, *Phys. Rev. D* **12**, 1502 (1975).
 - [17] B. Brahmachari, E. Ma, and U. Sarkar, *Phys. Rev. Lett.* **91**, 011801 (2003).
 - [18] F. Siringo, *Eur. Phys. J. C* **32**, 555 (2004).
 - [19] D. Chang, R.N. Mohapatra, and M.K. Parida, *Phys. Rev. Lett.* **52**, 1072 (1984); D. Chang, R.N. Mohapatra, and M.K. Parida, *Phys. Rev. D* **30**, 1052 (1984).