

Development of a Pixel Detector for Ultra-Cold Neutrons

S. Kawasaki^{*,a}, G. Ichikawa^a, M. Hino^d, Y. Kamiya^b, M. Kitaguchi^d,
S. Komamiya^{a,b}, T. Sanuki^c, S. Sonoda^a

^a*Graduate School of Science, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan*

^b*International Center for Elementary Particle Physics, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan*

^c*Graduate School of Science, Tohoku University, Sendai-shi, Miyagi 980-8578, Japan*

^d*Research Reactor Institute, Kyoto University, Sennan-gun, Osaka 590-0494, Japan*

Abstract

A pixel detector with high spatial resolution and temporal information for ultra-cold neutrons is developed based on a commercial CCD on which a neutron converter is attached. ^{10}B and ^6Li are tested for the neutron converter and ^{10}B is found to be more suitable based on efficiency and spatial resolution. The pixel detector has an efficiency of $44.1 \pm 1.1\%$ and a spatial resolution of $2.9 \pm 0.1 \mu\text{m}$ (1 sigma).

Key words: UCN, pixel detector

1. Introduction

When ultra-cold neutrons (UCNs) are trapped in the earth's gravitational field, their energy is quantized. In consequence, their probability density distribution exhibits their vertical modulation. The scale of this density modulation is calculated to be $(\hbar^2/2m_n^2g)^{1/3} \sim 6 \mu\text{m}$. Observations of such

^{*}Corresponding author: Tel: +81-3-5841-4201; Fax: +81-3-5841-4158

Email address: kawasaki@icepp.e.u-tokyo.ac.jp (S. Kawasaki)