

(12) United States Patent

Lewis et al.

US 10,159,746 B2 (10) **Patent No.:**

(45) Date of Patent: Dec. 25, 2018

(54) CYCLODEXTRIN COMPLEXATION METHODS FOR FORMULATING PEPTIDE PROTEASOME INHIBITORS

(71) Applicant: Onyx Therapeutics, Inc., South San

Francisco, CA (US)

Inventors: Evan Lewis, Pacifica, CA (US); Peter

Shwonek, San Francisco, CA (US); Sean Dalziel, San Francisco, CA (US); Mouhannad Jumaa, Foster City, CA

Assignee: Onyx Therapeutics, Inc., Thousand

Oaks, CA (US)

Subject to any disclaimer, the term of this (*) Notice:

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

14/399,582 (21) Appl. No.:

(22) PCT Filed: May 8, 2013

(86) PCT No.: PCT/US2013/040127

§ 371 (c)(1),

(2) Date: Nov. 7, 2014

(87) PCT Pub. No.: WO2013/169897

PCT Pub. Date: Nov. 14, 2013

(65)**Prior Publication Data**

> US 2015/0111838 A1 Apr. 23, 2015

Related U.S. Application Data

- Provisional application No. 61/777,475, filed on Mar. 12, 2013, provisional application No. 61/644,122, filed on May 8, 2012.
- (51) Int. Cl.

A61K 47/40 (2006.01)(2006.01)A61K 47/48

(Continued)

(52) U.S. Cl.

CPC A61K 47/48969 (2013.01); A61K 38/07 (2013.01); A61K 47/6951 (2017.08); B82Y 5/00 (2013.01)

(58) Field of Classification Search

See application file for complete search history.

(56)**References Cited**

U.S. PATENT DOCUMENTS

4.522.811 A 6/1985 Eppstein et al. 11/1998 Reuscher 5,831,081 A (Continued)

FOREIGN PATENT DOCUMENTS

WO WO-98/10779 A1 3/1998 WO WO-2002/30455 A2 4/2002 (Continued)

OTHER PUBLICATIONS

Whalen et al. ('Specific effects of chloride ion in epoxide hydrolysis. The pH-dependence of the rates and mechanisms for the

hydrolysis of indene oxide' Journal of the American Chemical Society v98(24) Nov. 24, 1976 pp. 7859-7861).*

(Continued)

Primary Examiner — Karlheinz R. Skowronek Assistant Examiner — Ronald T Niebauer

(74) Attorney, Agent, or Firm - Marshall, Gerstein &

Borun LLP

(57)**ABSTRACT**

This disclosure provides methods for formulating compositions comprising one or more peptide proteasome inhibitors and a cyclodextrin, particularly a substituted cyclodextrin. As well as cyclodextrin complexation methods of formulating a peptide proteasome inhibitor (e.g., a compound of formula (1)-(5) or a pharmaceutically acceptable salt thereof) with one or more cyclodextrins. Such methods substantially increase the solubility and stability of these proteasome inhibitors and facilitate both their manufacture and administration. For example, homogenous solutions of a compound of formula (5) (carfilzomib) can be obtained at a pharmaceutically useful pH (e.g., about 3.5) and at higher concentrations (e.g., about 5 mg/mL) than could be obtained without one or more cyclodextrins and the processes of complexation between the compound and one or more cyclodextrins provided herein.

$$\begin{array}{c} H_2C \\ X \\ R \end{array}$$

$$R_5 \underbrace{ \left(\begin{array}{c} O \\ H \\ R \end{array} \right) \left(\begin{array}{c} R^2 \\ H \\ O \end{array} \right) \left(\begin{array}{c} H \\ N \\ R^3 \end{array} \right) \left(\begin{array}{c} R^4 \\ H \\ O \end{array} \right) \left(\begin{array}{c} X \\ X \end{array} \right)$$

(3)

(5)

30 Claims, 8 Drawing Sheets Specification includes a Sequence Listing.