



FIG. 6: Ratio of $(1/\sigma) d\sigma/dp_T$ for top quarks in $t\bar{t}$ production (two entries per event) to the expectation from NLO pQCD. The gray band encompasses uncertainties on the scale of pQCD and parton distribution functions. Also shown are ratios relative to NLO pQCD for an approximate NNLO pQCD calculation and of predictions for several event generators. Inner and outer error bars represent statistical and total (statistical and systematic added in quadrature) uncertainties, respectively.

Culiacán, Mexico.

[g] Visitor from Universität Bern, Bern, Switzerland.

- [8] I. Bigi *et al.*, Phys. Lett. **B181**, 157 (1986).
- [9] C. T. Hill and S. J. Parke, Phys. Rev. D **49**, 4454 (1994).
- [10] D. Atwood *et al.*, Phys. Rev. D **52**, 6264 (1995).
- [11] T. Affolder *et al.*, CDF Collaboration, Phys. Rev. Lett. **87**, 102001 (2001);
B. Abbott *et al.*, D0 Collaboration, Phys. Rev. D **58**, 052001 (1998).
- [12] V. M. Abazov *et al.*, D0 Collaboration, Nucl. Instrum. Methods in Phys. Res. A **565**, 463 (2006).
- [13] G. C. Blazey *et al.*, in *Proceedings of the Workshop: QCD and Weak Boson Physics in Run II*, edited by U. Baur, R. K. Ellis, and D. Zeppenfeld, Fermilab-Pub-00/297 (2000).
- [14] V. M. Abazov *et al.*, D0 Collaboration, Nucl. Instrum. Methods in Phys. Res. A **620**, 400 (2010).
- [15] V. M. Abazov *et al.*, D0 Collaboration, Phys. Rev. D **80** (RC), 071102 (2009).
- [16] Pseudorapidity is defined as $\eta = -\ln \tan(\theta/2)$ where θ is the angle measured with respect to the proton beam coinciding with the positive z axis of a right-handed coordinate system at the center of the detector.
- [17] M. L. Mangano, *et al.*, J. High Energy Phys. **0307**, 001 (2003).
- [18] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001); R. Field and R. C. Group, arXiv:hep-ph/0510198.
- [19] J. Pumplin *et al.*, J. High Energy Phys. **0207**, 012 (2002).

- [20] E. E. Boos *et al.*, Phys. Atom. Nucl. **69**, 1317 (2006);
Yad. Fiz. **69**, 1352 (2006).
- [21] R. Brun and F. Carminati, CERN Program Library Long Writeup W5013, 1993 (unpublished).
- [22] V. M. Abazov *et al.*, D0 collaboration, Phys. Rev. D **75**, 092001 (2007);
- [23] S. Snyder, Ph.D. thesis, State University of New York at Stony Brook (1995), [Institution Report No. FERMILAB-THESIS-1995-27].
- [24] A. Hoecker and V. Kartvelishvili, Nucl. Instrum. Methods in Phys. Res. A **372**, 469 (1996).
- [25] V. Kartvelishvili, <http://www.lancs.ac.uk/users/spc/staff/kartv>
- [26] M. Mangano, P. Nason, and G. Ridolfi, Nucl. Phys. B **373**, 295 (1992).
- [27] P. Nason, S. Dawson, and R. K. Ellis, Nucl. Phys. B **327**, 49 (1989) [Erratum-ibid. B **335**, 260 (1990)].
- [28] D. Stump *et al.*, J. High Energy Phys. **0310**, 046 (2003).
- [29] N. Kidonakis and R. Vogt, Phys. Rev. D **78**, 074005 (2008).
- [30] A.D. Martin, W.J. Stirling, R.S. Thorne, G. Watt, Eur. Phys. J. C **63** 189 (2009).
- [31] S. Frixione and B. R. Webber, J. High Energy Phys. **0206**, 029 (2002); S. Frixione, P. Nason, and B. R. Webber, J. High Energy Phys. **0308**, 007 (2003).
- [32] N. D. Gagunashvili, Nucl. Instrum. Methods in Phys. Res. A **596**, 439 (2008).