

# Evidence of growing spatial correlations at the glass transition from nonlinear response experiments

C. Crauste-Thibierge<sup>1</sup>, C. Brun<sup>1</sup>, F. Ladieu<sup>1\*</sup>, D. L'Hôte<sup>1\*</sup>, G. Biroli<sup>2</sup>, and J-P. Bouchaud<sup>3\*</sup>

<sup>1</sup> *SPEC (CNRS URA 2464), DSM/IRAMIS CEA Saclay,  
Bat.772, F-91191 Gif-sur-Yvette France*

<sup>2</sup> *Institut de Physique Théorique, CEA,  
(CNRS URA 2306), 91191 Gif-sur-Yvette, France and*

<sup>3</sup> *Science & Finance, Capital Fund Management,  
6, Bd. Haussmann, 75009 Paris, France*

(Dated: November 26, 2018)

## Abstract

The ac nonlinear dielectric response  $\chi_3(\omega, T)$  of glycerol was measured close to its glass transition temperature  $T_g$  to investigate the prediction that supercooled liquids respond in an increasingly non-linear way as the dynamics slows down (as spin-glasses do). We find that  $\chi_3(\omega, T)$  indeed displays several non trivial features. It is peaked as a function of the frequency  $\omega$  and obeys scaling as a function of  $\omega\tau(T)$ , with  $\tau(T)$  the relaxation time of the liquid. The height of the peak, proportional to the number of dynamically correlated molecules  $N_{corr}(T)$ , increases as the system becomes glassy, and  $\chi_3$  decays as a power-law of  $\omega$  over several decades beyond the peak. These findings confirm the collective nature of the glassy dynamics and provide the first direct estimate of the  $T$  dependence of  $N_{corr}$ .

*Accepted for publication in Physical Review Letters*