

therefore

$$d_L^{(m)} = \sum_{m=0}^{2^L-1} h_m \quad (21)$$

for the Potts model, and

$$d_L^{S(m)} = \sum_{m=0}^{2^L-1} h_m^S \quad (22)$$

for the site percolation.

The partitions can again be coded by means of a sequence of integers. A detailed description of the coding algorithm can be found in [18]. The magnetic sectors of the transfer matrix now have the dimensions  $d_L^{(m)} \times d_L^{(m)}$  and  $d_L^{S(m)} \times d_L^{S(m)}$  for the Potts model and site percolation, respectively, and are much larger than those of the non-magnetic sectors. The magnetic sector of the transfer matrix can also be converted into a product of sparse matrices in the same way as in the case of the non-magnetic sector.

The inverse magnetic correlation length is given by

$$\frac{1}{\xi_h(v, L)} = \zeta \ln \left( \frac{\lambda_0}{\lambda'_0} \right), \quad (23)$$

where  $\lambda_0$  and  $\lambda'_0$  are the leading eigenvalues of the transfer matrix in the non-magnetic and magnetic sector respectively,  $\zeta$  is a geometrical factor which is the ratio between the unit of  $L$  and the thickness of a layer added by the transfer matrix. The magnetic scaled gap then follows.

According to finite-size scaling theory [25] and Cardy's conformal mapping [26],  $X_h(v, L)$  can be expanded as

$$X_h(v, L) = X_h + atL^{y_t} + buL^{y_u} + \dots, \quad (24)$$

where  $X_h$  is the magnetic scaling dimension,  $t$  is the deviation from the critical point, and  $u$  the irrelevant field. Here,  $y_t$  is the thermal renormalization exponent,  $y_u$  the leading irrelevant renormalization exponent, and  $a$  and  $b$  are unknown constants.

We substitute (24) into the finite-size scaling equation connecting lattices of sizes  $L$  and  $L - 1$ ,

$$X_h(v, L) = X_h(v, L - 1), \quad (25)$$

and denote the solution of (25) by  $v_c(L)$ , which has the expansion

$$v_c(L) = v_c + a'uL^{y_u - y_t} + \dots, \quad (26)$$