

Thus  $a, b \in A_i$  satisfy the conditions in step 2 for construction B. This contradicts our assumption that we used construction A, so there is an arrow formed by  $x_i$  and another vertex of  $D_{i+1}$ .

In construction B we have  $b_i y_i \in A(D_i)$  so  $b_i z \notin T_i$  for all  $z \in V(D_i) \setminus \{x_i\}$ . This gives  $b_i x_i \in \sigma_i$  and  $b_i x_i \in A(D_{i+1})$  so there is an arrow formed by  $x_i$  with another vertex of  $D_{i+1}$ .

**Acknowledgements** The author wishes to thank Martin Erickson of Truman State University and Steven Szydluk of the University of Wisconsin Oshkosh for encouragement and suggesting improvements to the writing in earlier drafts.

## References

- [1] G. Abrams, Groups Gradings and Recovery Results for Generalized Incidence Rings, *J. Alg.* **164** (1994) 859-875.
- [2] K. Kuratowski, Sur le problème des courbes gauches en topologie, *Fund. Math.* **15** (1930) 271–283.
- [3] K. Price, Good Gradings of Generalized Incidence Rings, submitted
- [4] E. Spiegel and C. O'Donnell, "Incidence Algebras," Marcel Dekker, New York, NY (1997).